

# FREE RESOLUTIONS OVER A COMPLETE HYPERSURFACE (AND FRIENDS)

BASED ON “HOMOLOGICAL ALGEBRA ON A COMPLETE  
INTERSECTION, WITH AN APPLICATION TO GROUP  
REPRESENTATIONS.” BY DAVID EISENBUD

David DeMark

MATH 8212 University of Minnesota

2 May 2018

**Throughout this talk, we let...**

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular sequence of finite length

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Such a  $B$  is a *complete intersection of codimension  $n$* .

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Such a  $B$  is a *complete intersection of codimension  $n$* .

We shall study the structure of  $B$ -free resolutions of  $B$ -modules, relating these to their liftings to  $A$ .

# CLARIFICATION: $B$ -FREE

We do **NOT** mean...



# CLARIFICATION: $B$ -FREE




FIGURE: Another “ $B$ -Free” Object.

Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Let  $\tilde{\partial}_i$  denote an arbitrary lifting of