

FREE RESOLUTIONS OVER A COMPLETE HYPERSURFACE (AND FRIENDS)

BASED ON “HOMOLOGICAL ALGEBRA ON A COMPLETE
INTERSECTION, WITH AN APPLICATION TO GROUP
REPRESENTATIONS.” BY DAVID EISENBUD

David DeMark

MATH 8212 University of Minnesota

2 May 2018

Throughout this talk, we let...

Throughout this talk, we let...

- (A, \mathfrak{m}) be a regular local ring

Throughout this talk, we let...

- (A, \mathfrak{m}) be a regular local ring
- x_1, \dots, x_n a regular A -sequence of finite length

Throughout this talk, we let...

- (A, \mathfrak{m}) be a regular local ring
- x_1, \dots, x_n a regular A -sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Throughout this talk, we let...

- (A, \mathfrak{m}) be a regular local ring
- x_1, \dots, x_n a regular A -sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Such a B is a *complete intersection of codimension n* .

Throughout this talk, we let...

- (A, \mathfrak{m}) be a regular local ring
- x_1, \dots, x_n a regular A -sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Such a B is a *complete intersection of codimension n* .

We shall study the structure of B -free resolutions of B -modules, relating these to their liftings to A .

CLARIFICATION: B -FREE

We do **NOT** mean. . .

CLARIFICATION: B -FREE

We do **NOT** mean...




FIGURE: Another “ B -Free” Object.

WHY DO WE CARE?

WHY DO WE CARE?

- Classifying maximal CM modules over complete hypersurfaces! (case $n = 1$)

THEOREM (6.1)

Let $x \in A$, $d = \dim A$, and $B := A/\langle x \rangle$. Then,

- For any B -module M and minimal free resolution \mathbf{F} , the truncation at F_{d+1} is periodic with period 2.
- \mathbf{F} periodic $\iff M$ is a maximal CM module w/o a free summand.
- If so, \mathbf{F} is induced a matrix factorization.

WHY DO WE CARE?

- Expanding upon a familiar characterization!

THEOREM (AUSLANDER-BUCHSBAUM-SERRE)

For (R, \mathfrak{m}) local, R is regular $\iff \text{gl dim}(R) < \infty$.

WHY DO WE CARE?

- Expanding upon a familiar characterization!

THEOREM (6.1)

For (R, \mathfrak{m}) local with $\dim R = d$, TFAE:

- *For some $x_1, \dots, x_{d+1} \in R$, $\mathfrak{m} = \langle x_1, \dots, x_{d+1} \rangle$ and \hat{O} is unmixed in $\hat{R}^{\mathfrak{m}}$ (i.e. all associated primes of \hat{O} are minimal).*
- *For any f.g. R -module M with minimal free resolution \mathbf{F} , the truncation of \mathbf{F} at degree $d + 1$ is periodic of period 2.*
- *There exists a free resolution $\mathbf{F} : \dots \rightarrow F_1 \rightarrow F_0 \rightarrow 0$ of $R_{\mathfrak{m}}/\mathfrak{m}R_{\mathfrak{m}}$ where for some n , $\text{rank} F_n < n$.*

We call such an R an abstract hypersurface.

WHY DO WE CARE?

- Expanding upon a familiar characterization!

THEOREM (6.1)

For (R, \mathfrak{m}) local with $\dim R = d$, TFAE:

- *For some $x_1, \dots, x_{d+1} \in R$, $\mathfrak{m} = \langle x_1, \dots, x_{d+1} \rangle$ and \hat{O} is unmixed in $\hat{R}^{\mathfrak{m}}$ (i.e. all associated primes of \hat{O} are minimal).*
- *For any f.g. R -module M with minimal free resolution \mathbf{F} , the truncation of \mathbf{F} at degree $d + 1$ is periodic of period 2.*
- *There exists a free resolution $\mathbf{F} : \dots \rightarrow F_1 \rightarrow F_0 \rightarrow 0$ of $R_{\mathfrak{m}}/\mathfrak{m}R_{\mathfrak{m}}$ where for some n , $\text{rank} F_n < n$.*

We call such an R an abstract hypersurface.

CODIM n : SET-UP

Recall that $B := A/\langle x_1, \dots, x_n \rangle$. Let M be a B -module, and \mathbf{F} a free resolution of B .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

CODIM n : SET-UP

Recall that $B := A/\langle x_1, \dots, x_n \rangle$. Let M be a B -module, and \mathbf{F} a free resolution of B .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Now let...

CODIM n : SET-UP

Recall that $B := A/\langle x_1, \dots, x_n \rangle$. Let M be a B -module, and \mathbf{F} a free resolution of B .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Now let...

- \tilde{F}_i be a free A -module with $\text{rank}_A \tilde{F}_i = \text{rank}_B F_i$

CODIM n : SET-UP

Recall that $B := A/\langle x_1, \dots, x_n \rangle$. Let M be a B -module, and \mathbf{F} a free resolution of B .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Now let...

- \tilde{F}_i be a free A -module with $\text{rank}_A \tilde{F}_i = \text{rank}_B F_i$
- $\tilde{\partial}_i$ denote an arbitrary lifting of (the entries of) ∂_i to A

Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

- $\tilde{\mathbf{F}}$ is *not* necessarily a complex, but
 $\text{Im}(\tilde{\partial}_i \circ \tilde{\partial}_{i+1}) \subset \langle x_1, \dots, x_n \rangle F_{i-1}$

Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

- $\tilde{\mathbf{F}}$ is *not* necessarily a complex, but
 $\text{Im}(\tilde{\partial}_i \circ \tilde{\partial}_{i+1}) \subset \langle x_1, \dots, x_n \rangle F_{i-1}$
- Write $\tilde{\partial}_i \otimes \tilde{\partial}_{i+1} = \sum_{j=1}^n x_j \tilde{t}_j$ for some transformation
 $\tilde{t}_j := \tilde{t}_j(A, \{x_i\}, \mathbf{F}) : \tilde{\mathbf{F}} \rightarrow \tilde{\mathbf{F}}.$

Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

- $\tilde{\mathbf{F}}$ is *not* necessarily a complex, but $\text{Im}(\tilde{\partial}_i \circ \tilde{\partial}_{i+1}) \subset \langle x_1, \dots, x_n \rangle F_{i-1}$
- Write $\tilde{\partial}_i \otimes \tilde{\partial}_{i+1} = \sum_{j=1}^n x_j \tilde{t}_j$ for some transformation $\tilde{t}_j := \tilde{t}_j(A, \{x_i\}, \mathbf{F}) : \tilde{\mathbf{F}} \rightarrow \tilde{\mathbf{F}}$.
- Finally, let $t_j = t_j(A, \{x_i\}, \mathbf{F}) : \mathbf{F} \rightarrow \mathbf{F}$ be defined by $t_j = B \otimes \tilde{t}_j$.

THE t_j HAVE MANY NICE PROPERTIES!

THEOREM (SUMMARY OF §1)

The t_j have many nice properties! These include:

THE t_j HAVE MANY NICE PROPERTIES!

THEOREM (SUMMARY OF §1)

The t_j have many nice properties! These include:

- *The t_j are unique (w/r/t fixed $A, \mathbf{F}, \{x_i\}$) up to homotopy.*

THE t_j HAVE MANY NICE PROPERTIES!

THEOREM (SUMMARY OF §1)

The t_j have many nice properties! These include:

- *The t_j are unique (w/r/t fixed $A, \mathbf{F}, \{x_i\}$) up to homotopy.*
- *The t_j are natural in the sense that they commute with morphisms of complexes up to homotopy.*

THE t_j HAVE MANY NICE PROPERTIES!

THEOREM (SUMMARY OF §1)

The t_j have many nice properties! These include:

- *The t_j are unique (w/r/t fixed $A, \mathbf{F}, \{x_i\}$) up to homotopy.*
- *The t_j are natural in the sense that they commute with morphisms of complexes up to homotopy.*
- *The t_j commute with each other up to homotopy.*

THE t_j HAVE MANY NICE PROPERTIES!

THEOREM (SUMMARY OF §1)

The t_j have many nice properties! These include:

- *The t_j are unique (w/r/t fixed $A, \mathbf{F}, \{x_i\}$) up to homotopy.*
- *The t_j are natural in the sense that they commute with morphisms of complexes up to homotopy.*
- *The t_j commute with each other up to homotopy.*
- *If M, N are B -modules, with B -free resolutions $\mathbf{F} \rightarrow M$, $\mathbf{G} \rightarrow N$ $t_j(\mathbf{F})$ and $t_j(\mathbf{G})$ induce the same map on $\mathrm{Tor}_{\bullet}^B(M, N)$.*

THE t_j HAVE MANY NICE PROPERTIES!

THEOREM (SUMMARY OF §1)

The t_j have many nice properties! These include:

- *The t_j are unique (w/r/t fixed $A, \mathbf{F}, \{x_i\}$) up to homotopy.*
- *The t_j are natural in the sense that they commute with morphisms of complexes up to homotopy.*
- *The t_j commute with each other up to homotopy.*
- *If M, N are B -modules, with B -free resolutions $\mathbf{F} \rightarrow M$, $\mathbf{G} \rightarrow N$ $t_j(\mathbf{F})$ and $t_j(\mathbf{G})$ induce the same map on $\mathrm{Tor}_\bullet^B(M, N)$.*
- *The t_j play only a little less well with ring morphisms $\alpha : A \rightarrow A'$.*

INTERLUDE: MATRIX FACTORIZATIONS

Let $x \in A$ and $\phi : F \rightarrow G$ a morphism of A -modules.

INTERLUDE: MATRIX FACTORIZATIONS

Let $x \in A$ and $\phi : F \rightarrow G$ a morphism of A -modules.

INTERLUDE: MATRIX FACTORIZATIONS

Let $x \in A$ and $\phi : F \rightarrow G$ a morphism of A -modules.

DEFINITION

The ordered pair (ψ, ϕ) is a *matrix factorization* of x if