

# MATH 8302 Take-Home Final

David DeMark

9 May 2018

1.)

We let  $f : S^2 \rightarrow T^2$  be any smooth map.

a.)

We recall that by de Rahm's theorem, we have an isomorphism of rings  $\Psi : H_{\text{dR}}^*(T^2) \xrightarrow{\sim} H^*(T^2, \mathbb{R})$ . As we showed in class last semester (or can be found in Hatcher),  $H^*(T^2, \mathbb{R})$  is generated in degree one by  $[\alpha], [\beta] \in H^1(T^2)$  under the relations  $\langle [\alpha] \cup [\beta] + [\beta] \cup [\alpha], [\alpha]^2, [\beta]^2 \rangle$ . In particular, we have that  $H^2(T^2, \mathbb{R}) \cong \mathbb{R}\{[\alpha] \cup [\beta]\}$ . We let  $[\omega_\alpha] = \Psi^{-1}([\alpha])$ , and  $[\omega_\beta] = \Psi^{-1}([\beta])$ . We recall as well that  $H^*(S^2, \mathbb{R})$  is generated in degree two by a single element  $[\gamma] \in H^2(S^2, \mathbb{R})$  modulo the relation  $\langle [\gamma] \cup [\gamma] \rangle$ . We let  $[\omega_\gamma] \in H_{\text{dR}}^2(S^2)$  be its image under de Rahm's theorem. We finally also recall that thanks to de Rahm's isomorphism of rings, the map  $f^*$  induces a graded morphism of  $\mathbb{R}$ -algebras.

**Proposition.** *For any closed  $\omega \in \Omega^2(T^2)$ ,  $f^*(\omega) \in \Omega^2(S^2)$  is exact.*

*Proof.* We let  $\omega \in \Omega^2(T^2)$  be an arbitrary closed 2-form on  $T^2$ . We suppose for the sake of contradiction that  $f^*(\omega)$  is not exact, that is  $f^*([\omega]) = r[\omega_\gamma]$  for some  $0 \neq r \in \mathbb{R}$ . Then, we have two cases to consider. If  $\omega$  is exact, we have that  $[\omega] = [0]$  in  $H^2(T^2)$ . Then,  $f^*([0]) \neq [0]$  contradicting that  $f^*$  is a ring homomorphism. On the other hand, if  $\omega$  is not exact, then  $[\omega] = q[\omega_\alpha] \wedge [\omega_\beta]$  for some  $0 \neq q \in \mathbb{R}$ . Then, we have that  $f^*(q[\omega_\alpha]) \wedge f^*([\omega_\beta]) = r[\omega_\gamma]$ , but as  $H_{\text{dR}}^1(S^2) = 0$  and  $f^*$  is a *graded* homomorphism, we then have that  $[0] \wedge [0] = r[\omega_\gamma]$ , contradicting that  $r \neq 0$ . This completes our proof.  $\square$

b.)

**Proposition.**  $\deg f = 0$ .

*Proof.* As a corollary of de Rahm's theorem that we showed in class, for smooth path-connected  $q$ -manifolds  $M, N$  with smooth  $f : M \rightarrow N$ ,  $\deg f$  can be calculated as the determinant of  $f^* : H_{\text{dR}}^q(N) \rightarrow H_{\text{dR}}^q(M)$ . As we showed above,  $f^*H_{\text{dR}}^2(T^2) \rightarrow H_{\text{dR}}^2(S^2)$  is the zero map, and hence  $\deg f = 0$ .  $\square$

2.)

We let  $f : \mathbb{C}^2 \rightarrow \mathbb{C}$  be defined by  $f(z, w) = w^2 - z^3$ , and let  $X \subseteq \mathbb{C}^2$  be defined by  $X := \{(z, w) \mid f(w, z) = c\}$ .

a.)

We use the diffeomorphism  $\mathbb{C}^2 \xrightarrow{\sim} \mathbb{R}^4$  by  $(r + is, x + iy) \mapsto (r, s, x, y)$  and identify  $\mathbb{C}^2$  as  $\mathbb{R}^4$  as such (and similarly identify  $\mathbb{C}$  with  $\mathbb{R}^2$ ). Then,  $f : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  can be given new coordinates as

$$f(r, s, x, y) = (x^2 - y^2 - r^3 + 3rs^2, 2xy - 3r^2s + s^3).$$

Then, letting  $c = a + ib$ ,  $X$  is defined as  $X = X_c := \{(r, s, x, y) \mid f(r, s, x, y) = (a, b)\}$

**Proposition.** *For any  $(a, b) \neq (0, 0)$ ,  $X_c$  is a smooth manifold.*

As  $f$  is a smooth map with Euclidean domain and codomain, we may compute  $df_P$  as the Jacobian of  $f$  at  $P = (r, s, x, y)$ . Doing so yields the following:

$$df_P = \begin{bmatrix} 3(s^2 - r^2) & 6rs & 2x & -2y \\ -6rs & 3(s^2 - r^2) & 2y & 2x \end{bmatrix}$$

We have that  $c = (a, b) \in \mathbb{R}^2$  is a critical value if for some  $P = (r, s, x, y)$  with  $f(P) = c$ , the derivative  $df_P$  fails to be surjective, i.e. its maximal minors are identically zero. In particular, if  $P$  is a critical point, this forces the relations

$\Delta_{12}(\mathrm{d}f_P) = 9(s^2 - r^2)^2 + 36r^2s^2 = 9(s^2 + r^2)^2 = 0$  and  $\Delta_{34}(\mathrm{d}f_P) = 4(x^2 + y^2) = 0$ . As  $r, s, x, y$  are **real** coordinates, these two relations force  $(r, s, x, y) = \vec{0}$ . Hence,  $\vec{0} \in \mathbb{R}^4$  is the only critical point of  $f$ , and  $\vec{0} \in \mathbb{R}^2$  its corresponding critical value. By the regular value theorem, for  $c = (a, b) \neq (0, 0)$ , we have that  $X_c$  is a smooth manifold of dimension  $\dim \mathbb{R}^4 - \dim \mathbb{R}^2 = 2$ .

**b.)**

We let<sup>1</sup>  $g : \mathbb{C}^2 \rightarrow \mathbb{C}$  be defined by  $g(z, w) = w$ . Under our change of coordinates above, this is the map  $g : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $g(r, s, x, y) = (x, y)$ .

**Proposition.** *The critical points of  $g$  are exactly the plane  $r = s = 0$ .*

*Proof.* We have by a proposition on page 24 of Guillemin-Pollack that  $T_P X = \ker \mathrm{d}f_P$ . Assuming that  $P \neq \vec{0}$  (justified by part a), we compute manually that

$$\ker \mathrm{d}f_P \supseteq \text{Span} \left\{ v_1 = \begin{bmatrix} 2x \\ 2y \\ -3(s^2 - r^2) \\ 6rs \end{bmatrix}, v_2 = \begin{bmatrix} 2y \\ -2x \\ 6rs \\ 3(s^2 - r^2) \end{bmatrix} \right\}$$

We note that the linear independence of  $v_1$  and  $v_2$  can be checked by computing the rank of the matrix  $[v_1 \ v_2]$ . Doing so reveals the same maximal minors as  $\mathrm{d}f_P$ , implying that  $v_1$  and  $v_2$  span the tangent space at  $P$  for any  $P \in X$  by part a. We take  $v_1$  and  $v_2$  to be a basis for  $T_P X$ . Then, the inclusion map  $i : T_P X \rightarrow T_P \mathbb{R}^4$  is given by  $i(v) = [v_1 \ v_2]v$ . We let  $h$  be the extension of  $g$  to all of  $\mathbb{R}^4$  given coordinate-wise by the same formula as  $g$ . Then, we may compute  $\mathrm{d}h_P$  as

$$\mathrm{d}h_P = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Furthermore, we have that  $g = h \circ i$  and thus  $\mathrm{d}g_P = \mathrm{d}h_{i(P)} \mathrm{d}i_P$ . We compute

$$\mathrm{d}g_P = (\mathrm{d}h_{i(P)} \circ \mathrm{d}i_P)(v) = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2x & 2y \\ 2y & -2x \\ -3(s^2 - r^2) & 6rs \\ 6rs & 3(s^2 - r^2) \end{bmatrix} v = \begin{bmatrix} -3(s^2 - r^2) & 6rs \\ 6rs & 3(s^2 - r^2) \end{bmatrix} v$$

We have that  $P$  is a critical point of  $g$  if  $\mathrm{d}g_P$  fails to be surjective, i.e. fails to be an isomorphism. We compute  $\det \mathrm{d}g_P = 9(s^2 + r^2)^2$  and have that  $\det \mathrm{d}g_P = 0 \iff (s, r) = (0, 0)$ . As  $\det \mathrm{d}g_P = 0$  if and only if  $P$  is a critical point, this proves our claim.  $\square$

**3.)**

**Proposition.** *If  $\omega$  is a closed 2-form on  $S^4$ , then  $\omega \wedge \omega$  vanishes somewhere on  $S^4$ .*

*Proof.* We break our proof into a series of claims.

**Claim 3.A.**  $\omega \wedge \omega$  is closed

*Proof.* We write  $d(\omega \wedge \omega) = d\omega \wedge \omega + \omega \wedge d\omega = 0 \wedge \omega + \omega \wedge 0 = 0$ .  $\blacksquare$

**Claim 3.B.**  $\omega \wedge \omega$  is exact.

*Proof.* We recall that de Rahm's isomorphism  $\Psi : H_{\mathrm{dR}}^*(S^4) \xrightarrow{\sim} H^*(S^4, \mathbb{R})$  induces a ring structure via the wedge product on  $H_{\mathrm{dR}}^*(S^4)$ . We have that  $\omega$  is necessarily exact as it is closed and  $H^2(S^4) = 0$ . Thus,  $[\omega] = [0] \in H^2(S^4)$ , so if  $[\omega \wedge \omega] \neq [0]$ , we have that  $0 \neq [\omega \wedge \omega] = [\omega] \wedge [\omega] = [0] \wedge [0] = [0]$ , a contradiction. Thus,  $[\omega \wedge \omega] = [0]$  indeed, i.e.  $\omega \wedge \omega$  is exact.  $\blacksquare$

As an immediate corollary, as de Rahm's isomorphism works by integration on  $H_{\mathrm{dR}}^4(S^4)$ , we have that  $\int_{S^4} \omega \wedge \omega = 0$ .

**Lemma 3.C** (Mean value theorem for smooth manifolds). *If  $M$  is a  $d$ -manifold and  $\sigma \in \Omega^d(M)$  is exact, then there is some point  $p \in M$  such that where  $\omega = f(x) \mathrm{d}x_1 \wedge \cdots \wedge \mathrm{d}x_d$ ,  $f(p) = 0$ .*

*Proof.*   $\square$

use the  
mean va  
theorem  
integrati

<sup>1</sup>Oops, I'm changing names on you—my bad.

**4.)**

**a.)**

**b.)**

**5.)**