

Math 1271 - Lectures 010 and 030

Name (Print): Jessica Carlson

Fall 2017

Quiz 8C

11/07/17

Time Limit: 25 Minutes

Teaching Assistant \_\_\_\_\_

You may *not* use your books, notes, graphing calculator, phones or any other internet devices on this exam.

You are required to show your work on each problem on this quiz.

Problem	Points	Score
1	3	2
2	4	1.5
3	3	1.5
Total:	10	4

1. (3 points) Starting with the initial guess  $x_1 = -2$ , use Newton's method to approximate a root to the equation  $e^x + x^2 - 3 = 0$  to eight decimal places.

$$-2 - \frac{e^x + x^2 - 3}{e^x + 2x} = -1.706$$

+2

This doesn't work if you only record yr  $x_n$ 's to 3 decimal places.

$$-1.706 - \frac{e(-1.706) + (-1.706)^2 - 3}{e(-1.706) + 2(-1.706)} = -1.6775$$

$$-1.6775 - \frac{e(-1.6775) + (-1.6775)^2 - 3}{e(-1.6775) + 2(-1.6775)} = -1.6772$$

$$-1.6772 - \frac{e(-1.6772) + (-1.6772)^2 - 3}{e(-1.6772) + 2(-1.6772)} = -1.6772778$$


2. (4 points) If  $600\pi \text{ cm}^2$  material is available to make a cylinder with an open top, find the largest possible volume of the cylinder.

Hint: The surface area of a cylinder with an open top is  $\pi r^2 + 2\pi rh$ , where  $r$  is the base radius,  $h$  is the height.

\* Surface Area =  $\pi r^2 + 2\pi rh = 600\pi \text{ cm}^2$

\* Volume =  $\pi r^2 h$

+1.5



$$\pi r^2 + 2\pi rh = 600\pi \text{ cm}^2$$

$$r(\pi r + 2\pi h) = 600\pi \text{ cm}^2$$

3. (3 points) Show that the curve  $y = \sqrt{x^2 + 5} + 2x$  has one slant asymptote at  $y = 3x$  and one horizontal asymptote at  $y = 0$ .

\*  $\sqrt{x^2 + 5} + 2x \xrightarrow{\text{that doesn't work}} x + 5 + 2x \rightarrow 3x + 5$

+1.5

$$\lim_{x \rightarrow \infty} 3x + 5 = \infty$$

\*  $\sqrt{x^2 + 5} + 2x$