Math 1271 - Lectures 010 and 030 **Name (Print):** Ted Sorensen
Fall 2017
Quiz 8C
11/07/17
Time Limit: 25 Minutes Teaching Assistant David Dennerlein

You may *not* use your books, notes, graphing calculator, phones or any other internet devices on this exam.

You are required to show your work on each problem on this quiz.

Problem	Points	Score
1	3	3
2	4	1.75
3	3	.75
Total:	10	5.5

1. (3 points) Starting with the initial guess $x_1 = -2$, use Newton's method to approximate a root to the equation $e^x + x^2 - 3 = 0$ to eight decimal places.

$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = -2 - \frac{e^{-2} + 4 - 3}{e^{-2} - 4} = -2 - \frac{1.13533528}{-3.86466471} = -1.70622670$

$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)} = -2 - \frac{e^{-1.70622670} + 2.91120955 - 3}{e^{-1.70622670} - 3.41245340} = -1.70622670 - \frac{.09275909}{-3.23890386}$

$x_4 = x_3 - \frac{f(x_3)}{f'(x_3)} = -1.67751675 - \frac{.18683736 + 2.81406245 - 3}{.18683736 - 3.35503350} = \frac{.00089981}{-3.16819614} = -1.67723274 = -1.67751675$

$x_5 = -1.67723274 - \frac{.18689044 + 2.81310966 - 3}{.18689044 - 3.35446555} = \frac{.00000001}{-3.16757506} = -1.67723271$

$x_6 = -1.67723271$ + 3 stop at x_6

2. (4 points) If 600π cm^2 material is available to make a cylinder with an open top, find the largest possible volume of the cylinder.
Hint: The surface area of a cylinder with an open top is $\pi r^2 + 2\pi rh$, where r is the base radius, h is the height.

$h = \frac{\pi r^2}{2\pi r}$? $SA = \pi r^2 + 2\pi r\left(\frac{\pi r^2}{2\pi r}\right)$

$600\pi = \pi r^2 + 2\pi r\left(\frac{\pi r^2}{2\pi r}\right)$

+1.75



find r using 600π cm^2 in the equation I found then when you find r you use $h = \frac{\pi r^2}{2\pi r}$ to get h & you have height & radius.

3. (3 points) Show that the curve $y = \sqrt{x^2+5} + 2x$ has one slant asymptote at $y = 3x$ and one horizontal asymptote at $y = 0$.

$y = \frac{2x}{x+\sqrt{5}} \sim 2x$ assumption

$\lim_{x\to\infty} f(x) - 2x = y$-int

+.75