$\begin{array}{ll} \alpha,\,\beta & \text{type variables} \\ n,\,m,\,i,\,j & \text{index variables} \end{array}$

```
[\sigma]N
                                    Μ
                                          positive variable list
                                             empty list
                                             a variable
                                             concatenate lists
                                          negative variables
                                             empty list
                                             a variable
                                             concatenate lists
P, Q
                                          multi-quantified positive types
                       \alpha^+
                       {\downarrow}N
                       \exists \alpha^{-}.P
                                             P \neq \exists \dots
                       [\sigma]P
                                    Μ
                       [\mu]P
                                    Μ
                                    S
                       (P)
                       P_1 \vee P_2
                                    Μ
                       \mathbf{nf}(P')
                                    Μ
N, M
                                          multi-quantified negative types
                       \alpha^{-}
                       \uparrow P
                                            N \neq \forall \dots
                       [\sigma]N
                                    Μ
                       [\mu]N
                                    Μ
                       (N)
                       \mathbf{nf}(N')
                                    Μ
\overrightarrow{P}
                                          list of positive types
                                             empty list
                                             a singel type
                                             concatenate lists
\overrightarrow{N}
                                          list of negative types
                                             empty list
                                             a singel type
                                             concatenate lists
Γ
                                          declarative type context
                                             empty context
                                             list of variables
                                             list of variables
                       vars
                                             concatenate contexts
```

S

 (Γ)

```
\vec{\alpha}, \vec{\beta}
                                                     ordered positive or negative variables
                                                        empty list
                                                        list of variables
                                                        list of variables
                       \overrightarrow{\alpha}_1 \backslash vars
                                                        setminus
                                                        context
                       vars
                       \overline{\overrightarrow{\alpha}_i}^i
                                                        concatenate contexts
                                              S
                       (\vec{\alpha})
                                                        parenthesis
                       [\mu]\vec{\alpha}
                                              Μ
                                                        apply moving to list
                       ord vars in P
                                              Μ
                       \mathbf{ord}\ vars \mathbf{in}\ N
                                              Μ
                       ord varsin P
                                              Μ
                       \operatorname{\mathbf{ord}} \operatorname{\mathbf{\mathit{vars}}} \operatorname{\mathbf{in}} N
                                              Μ
                                                     set of variables
vars
                       Ø
                                                        empty set
                       \mathbf{fv} P
                                                        free variables
                       \mathbf{fv} N
                                                        free variables
                       \mathbf{fv} P
                                                        free variables
                                                        free variables
                       \mathbf{fv} N
                       vars_1 \cap vars_2
                                                        set intersection
                                                        set union
                       vars_1 \cup vars_2
                                                        set complement
                       vars_1 \backslash vars_2
                       \mathbf{mv} P
                                                        movable variables
                                                        movable variables
                       \mathbf{mv} N
                       \mathbf{u}\mathbf{v} N
                                                        unification variables
                       \mathbf{uv} P
                                                        unification variables
                       \mathbf{fv} N
                                                        free variables
                       \mathbf{fv} P
                                                        free variables
                       (vars)
                                              S
                                                        parenthesis
                       \{\vec{\alpha}\}
                                                        ordered list of variables
                       [\mu]vars
                                              Μ
                                                         apply moving to varset
                                                        empty moving
                                                         Positive unit substitution
                                                        Positive unit substitution
                                              Μ
                                                        Set-like union of movings
                                                        concatenate movings
                                              Μ
                       \mu|_{vars}
                                                        restriction on a set
                                              Μ
                                                        inversion
                                                     cohort index
n
                       0
                       n+1
\tilde{\alpha}^+
                                                     positive movable variable
```

```
negative movable variable
                     \tilde{\alpha}^{-n}
                                      positive movable variable list
                                         empty list
                                         a variable
                                         from a non-movable variable
                                         concatenate lists
                                      negatiive movable variable list
                                         empty list
                                         a variable
                                         from a non-movable variable
                                         concatenate lists
P, Q
                                      multi-quantified positive types with movable variables
                     \alpha^+
                     \tilde{\alpha}^+
                     \downarrow N
                      [\sigma]P
                                 Μ
                                 Μ
N, M
                                      multi-quantified negative types with movable variables
                     \alpha^{-}
                     \tilde{\alpha}^-
                     \uparrow P
                                 Μ
                      [\mu]N
                                 Μ
\hat{\alpha}^+
                                      positive unification variable
               ::=
                     \hat{\alpha}^+
\hat{\alpha}^-
                                      negative unification variable
                                      positive unification variable list
                                         empty list
                                         a variable
                                         from a normal variable
                                         from a normal variable, context unspecified
                                         concatenate lists
                                      negative unification variable list
                                         empty list
                                         a variable
                                                        5
```

```
from a normal variable
                                                        from a normal variable, context unspecified
                                                        concatenate lists
P, Q
                                                    a positive algorithmic type (potentially with metavariables)
                             \alpha^+
                             \widetilde{\alpha}^+
                            \hat{\alpha}^+ \{vars\}
                             [\sigma]P
                                             Μ
                             [\mu]P
                                             Μ
                             \mathbf{nf}(P')
                                             Μ
N, M
                                                    a negative algorithmic type (potentially with metavariables)
                             \alpha^{-}
                            \tilde{\alpha}^-
                             \hat{\alpha}^- \{vars\}
                             \uparrow P
                             P \rightarrow N
                            \forall \overrightarrow{\alpha^+}.N
                             [\sigma]N
                                             Μ
                             [\mu]N
                                             Μ
                             \mathbf{nf}(N')
                                             Μ
terminals
                             \exists
                             \forall
                             \geqslant
                             Ø
```

```
\neq
                                  :≥
formula
                                  judgement
                                  formula_1 .. formula_n
                                  \mu : vars_1 \leftrightarrow vars_2
                                  \mu is bijective
                                  \hat{\sigma} is functional
                                  \hat{\sigma}_1 \in \hat{\sigma}_2
                                  \mathit{vars}_1 \subseteq \mathit{vars}_2
                                  vars_1 = vars_2
                                  vars is fresh
                                  \alpha^- \not\in \mathit{vars}
                                  \alpha^+ \notin vars
                                  \alpha^- \in vars
                                  \alpha^+ \in vars
                                  if any other rule is not applicable
                                  N \neq M
                                  P \neq Q
E1A
                                 n \vDash N \simeq_1^A M \rightrightarrows \mun \vDash P \simeq_1^A Q \rightrightarrows \mu
                                                                                                                Negative multi-quantified type equivalence (algorit
                                                                                                                Positive multi-quantified type equivalence (algorith
A
                                  \begin{array}{l} \Gamma \vDash N \leqslant M \dashv \widehat{\sigma} \\ \Gamma \vDash P \geqslant Q \dashv \widehat{\sigma} \end{array}
                                                                                                                Negative subtyping
                                                                                                                Positive supertyping
E1
                                 N \simeq_1^D MP \simeq_1^D Q
                                                                                                                Negative multi-quantified type equivalence
                                                                                                                Positive multi-quantified type equivalence
D1
                               \Gamma \vdash N \simeq_1^{\varsigma} M
\Gamma \vdash P \simeq_1^{\varsigma} Q
\Gamma \vdash N \leqslant_1 M
\Gamma \vdash P \geqslant_1 Q
                                                                                                                Negative equivalence on MQ types
                                                                                                                Positive equivalence on MQ types
                                                                                                                Negative subtyping
                                                                                                                Positive supertyping
D\theta
                        ::=
                                \begin{array}{l} \Gamma \vdash N \simeq_0^{\scriptscriptstyle \leqslant} M \\ \Gamma \vdash P \simeq_0^{\scriptscriptstyle \leqslant} Q \\ \Gamma \vdash N \leqslant_0 M \\ \Gamma \vdash P \geqslant_0 Q \end{array}
                                                                                                                Negative equivalence
                                                                                                                Positive equivalence
                                                                                                                Negative subtyping
                                                                                                                Positive supertyping
```

A

E1D1 $D\theta$ LUBAUNrmOrderSMUWF

 $user_syntax$

 α

n α^+

 α^{-}

P $\overrightarrow{\alpha^{+}}$ $\overrightarrow{\alpha^{-}}$ P \overrightarrow{N} \overrightarrow{P} \overrightarrow{N}

Γ

vars

 μ

P

terminals

formula

 $n \models N \simeq^A_1 M \dashv \mu$

Negative multi-quantified type equivalence (algorithmic)

$$\frac{n \vDash \alpha^{-} \simeq_{1}^{A} \alpha^{-} \dashv}{n \vDash P \simeq_{1}^{A} Q \dashv \mu} \quad \text{E1ANVAR}$$

$$\frac{n \vDash P \simeq_{1}^{A} Q \dashv \mu}{n \vDash \uparrow P \simeq_{1}^{A} \uparrow Q \dashv \mu} \quad \text{E1ASHIFTU}$$

$$\frac{n \vDash P \simeq_{1}^{A} Q \dashv \mu_{1} \quad n \vDash N \simeq_{1}^{A} M \dashv \mu_{2} \quad \mu_{1} \cup \mu_{2} \text{ is bijective}}{n \vDash P \to N \simeq_{1}^{A} Q \to M \dashv \mu_{1} \cup \mu_{2}} \quad \text{E1AARROW}$$

$$\frac{n+1 \vDash [\overrightarrow{\alpha^{+n}}/\overrightarrow{\alpha^{+}}] N \simeq_{1}^{A} [\overrightarrow{\beta^{+n}}/\overrightarrow{\beta^{+}}] M \dashv \mu}{n \vDash \forall \overrightarrow{\alpha^{+}}. N \simeq_{1}^{A} \forall \overrightarrow{\beta^{+}}. M \dashv \mu|_{\mathbf{mv} M}} \quad \text{E1ANMVAR}$$

$$\frac{1}{n \models \widetilde{\alpha}^{-n} \simeq_1^A \widetilde{\beta}^{-n} \rightrightarrows \widetilde{\beta}^{-n} \mapsto \widetilde{\alpha}^{-n}} \quad \text{E1ANMVAR}$$

 $n \models P \simeq_1^A Q = \mu$ Positive multi-quantified type equivalence (algorithmic)

$$\frac{n \vDash \alpha^{+} \simeq_{1}^{A} \alpha^{+} \dashv \cdot}{n \vDash \sqrt{N} \simeq_{1}^{A} \sqrt{M} \dashv \mu} \quad \text{E1ASHIFTD}$$

$$\frac{n \vDash N \simeq_{1}^{A} M \dashv \mu}{n \vDash \sqrt{N} \simeq_{1}^{A} \sqrt{M} \dashv \mu} \quad \text{E1ASHIFTD}$$

$$\frac{n+1 \vDash [\overrightarrow{\alpha^{-n}}/\overrightarrow{\alpha^{-}}]P \simeq_{1}^{A} [\overrightarrow{\beta^{-n}}/\overrightarrow{\beta^{-}}]Q \dashv \mu}{n \vDash \overrightarrow{\alpha^{-}} \cdot P \simeq_{1}^{A} \exists \overrightarrow{\beta^{-}} \cdot Q \dashv \mu|_{\mathbf{mv} Q}} \quad \text{E1AEXISTS}$$

$$\frac{n+1 \vDash [\overrightarrow{\alpha^{-n}}/\overrightarrow{\alpha^{-}}]P \simeq_{1}^{A} (\overrightarrow{\beta^{-n}}/\overrightarrow{\beta^{-}})Q \dashv \mu|_{\mathbf{mv} Q}}{n \vDash \overrightarrow{\alpha^{+n}} \simeq_{1}^{A} \widetilde{\beta^{+n}} \dashv \widetilde{\beta^{+n}} \mapsto \widetilde{\alpha^{+n}}} \quad \text{E1APMVAR}$$

 $\Gamma \models N \leqslant M \dashv \widehat{\sigma}$ Negative subtyping

$$\frac{\mathbf{nf}(P) \stackrel{u}{\simeq} \mathbf{nf}(Q) \dashv \widehat{\sigma}}{\Gamma \vDash \Lambda} \quad \text{ANVAR}$$

$$\frac{\mathbf{nf}(P) \stackrel{u}{\simeq} \mathbf{nf}(Q) \dashv \widehat{\sigma}}{\Gamma \vDash \Lambda} \quad \text{ASHIFTU}$$

$$\frac{\Gamma \vDash P \geqslant Q \dashv \widehat{\sigma}_{1} \quad \Gamma \vDash N \leqslant M \dashv \widehat{\sigma}_{2}}{\Gamma \vDash P \rightarrow N \leqslant Q \rightarrow M \dashv \widehat{\sigma}_{1} \& \widehat{\sigma}_{2}} \quad \text{AARROW}$$

$$\frac{\Gamma, \overrightarrow{\beta^{+}} \vDash [\widehat{\alpha}^{+} \{\Gamma, \overrightarrow{\beta^{+}}\} / \overrightarrow{\alpha^{+}}] N \leqslant M \dashv \widehat{\sigma}}{\Gamma \vDash \forall \alpha^{+} . N \leqslant \forall \overrightarrow{\beta^{+}} . M \dashv \widehat{\sigma} \backslash \widehat{\alpha^{+}}} \quad \text{AFORALL}$$

 $\Gamma \vDash P \geqslant Q \dashv \widehat{\sigma}$ Positive supertyping

$$\frac{\mathbf{nf}(N) \stackrel{u}{\simeq} \mathbf{nf}(M) \dashv \widehat{\sigma}}{\Gamma \vDash \downarrow N \geqslant \downarrow M \dashv \widehat{\sigma}} \quad \text{ASHIFTD}$$

$$\frac{\mathbf{nf}(N) \stackrel{u}{\simeq} \mathbf{nf}(M) \dashv \widehat{\sigma}}{\Gamma \vDash \downarrow N \geqslant \downarrow M \dashv \widehat{\sigma}} \quad \text{ASHIFTD}$$

$$\frac{\Gamma, \overrightarrow{\beta^-} \vDash [\widehat{\alpha}^- \{\Gamma, \overrightarrow{\beta^-}\} / \overrightarrow{\alpha^-}] P \geqslant Q \dashv \widehat{\sigma}}{\Gamma \vDash \exists \overrightarrow{\alpha^-} . P \geqslant \exists \overrightarrow{\beta^-} . Q \dashv \widehat{\sigma}} \quad \text{AEXISTS}$$

$$\frac{\mathbf{nf}(P) = P' \quad vars_1 = \mathbf{fv} P' \setminus vars \quad vars_2 \mathbf{is} \mathbf{fresh}}{\Gamma \vDash \widehat{\alpha}^+ \{vars\} \geqslant P \dashv (\widehat{\alpha}^+ : \geqslant P' \vee [vars_2 / vars_1] P')} \quad \text{APUVAR}$$

 $|N \simeq_1^D M|$ Negative multi-quantified type equivalence

$$\frac{\overline{\alpha^{-}} \simeq_{1}^{D} \alpha^{-}}{\alpha^{-}} \quad \text{E1NVAR}$$

$$\frac{P \simeq_{1}^{D} Q}{\uparrow P \simeq_{1}^{D} \uparrow Q} \quad \text{E1SHIFTU}$$

$$\frac{P \simeq_{1}^{D} Q \quad N \simeq_{1}^{D} M}{P \to N \simeq_{1}^{D} Q \to M} \quad \text{E1ARROW}$$

$$\frac{\{\overrightarrow{\alpha^{+}}\} \cap \text{fv } M = \varnothing \quad \mu : (\{\overrightarrow{\beta^{+}}\} \cap \text{fv } M) \leftrightarrow (\{\overrightarrow{\alpha^{+}}\} \cap \text{fv } N) \quad N \simeq_{1}^{D} [\mu] M}{\forall \overrightarrow{\alpha^{+}} . N \simeq_{1}^{D} \forall \overrightarrow{\beta^{+}} . M}$$

$$\text{E1FORALL}$$

 $P \simeq_{1}^{D} Q$ Positive multi-quantified type equivalence

$$\frac{\alpha^{+} \simeq_{1}^{D} \alpha^{+}}{\alpha^{+} \simeq_{1}^{D} M} \quad \text{E1ShiftD}$$

$$\frac{N \simeq_{1}^{D} M}{\downarrow N \simeq_{1}^{D} \downarrow M} \quad \text{E1ShiftD}$$

$$\frac{\{\overrightarrow{\alpha^{-}}\} \cap \text{fv } Q = \varnothing \quad \mu : (\{\overrightarrow{\beta^{-}}\} \cap \text{fv } Q) \leftrightarrow (\{\overrightarrow{\alpha^{-}}\} \cap \text{fv } P) \quad P \simeq_{1}^{D} [\mu]Q}{\exists \overrightarrow{\alpha^{-}}.P \simeq_{1}^{D} \exists \overrightarrow{\beta^{-}}.Q} \quad \text{E1Exist}$$

 $\overline{|\Gamma \vdash N \simeq_1^{\leq} M|}$ Negative equivalence on MQ types

$$\frac{\Gamma \vdash N \leqslant_{1} M \quad \Gamma \vdash M \leqslant_{1} N}{\Gamma \vdash N \simeq_{1}^{s} M} \quad \text{D1NDEF}$$

 $\Gamma \vdash P \simeq_1^{\varsigma} Q$ Positive equivalence on MQ types

$$\frac{\Gamma \vdash P \geqslant_1 Q \quad \Gamma \vdash Q \geqslant_1 P}{\Gamma \vdash P \simeq_1^{\varsigma} Q} \quad \text{D1PDEF}$$

 $\Gamma \vdash N \leq_1 M$ Negative subtyping

$$\frac{\Gamma \vdash \alpha^{-} \leqslant_{1} \alpha^{-}}{\Gamma \vdash P \rightleftharpoons_{1}^{\circ} Q} \quad D1SHIFTU$$

$$\frac{\Gamma \vdash P \geqslant_{1} Q \quad \Gamma \vdash N \leqslant_{1} M}{\Gamma \vdash P \rightarrow N \leqslant_{1} Q \rightarrow M} \quad D1ARROW$$

$$\frac{\Gamma, \overrightarrow{\beta^{+}} \vdash P_{i} \quad \Gamma, \overrightarrow{\beta^{+}} \vdash [\overrightarrow{P}/\alpha^{+}]N \leqslant_{1} M}{\Gamma \vdash \forall \alpha^{+}, N \leqslant_{1} \forall \overrightarrow{\beta^{+}}, M} \quad D1FORALL$$

 $\Gamma \vdash P \geqslant_1 Q$ Positive supertyping

$$\frac{\Gamma \vdash \alpha^{+} \geqslant_{1} \alpha^{+}}{\Gamma \vdash N \cong_{1}^{\leq} M} \quad D1PVAR$$

$$\frac{\Gamma \vdash N \cong_{1}^{\leq} M}{\Gamma \vdash \downarrow N \geqslant_{1} \downarrow M} \quad D1SHIFTD$$

$$\frac{\Gamma, \overrightarrow{\beta^{-}} \vdash N_{i} \quad \Gamma, \overrightarrow{\beta^{-}} \vdash [\overrightarrow{N}/\overrightarrow{\alpha^{-}}]P \geqslant_{1} Q'}{\Gamma \vdash \exists \overrightarrow{\alpha^{-}}.P \geqslant_{1} \exists \overrightarrow{\beta^{-}}.Q} \quad D1EXISTSL$$

 $\Gamma \vdash N \simeq_0^{\leq} M$ Negative equivalence

$$\frac{\Gamma \vdash N \leqslant_0 M \quad \Gamma \vdash M \leqslant_0 N}{\Gamma \vdash N \simeq_0^{\varsigma} M} \quad \text{D0NDEF}$$

 $\Gamma \vdash P \simeq_0^{\leqslant} Q$ Positive equivalence

$$\frac{\Gamma \vdash P \geqslant_0 Q \quad \Gamma \vdash Q \geqslant_0 P}{\Gamma \vdash P \simeq_0^{\varsigma} Q} \quad \text{D0PDeF}$$

 $\Gamma \vdash N \leq_0 M$ Negative subtyping

$$\frac{\Gamma \vdash a - \leqslant_0 a -}{\Gamma \vdash P = \circ_0^{\leqslant} Q} \quad D0\text{NVAR}$$

$$\frac{\Gamma \vdash P = \circ_0^{\leqslant} Q}{\Gamma \vdash \uparrow P \leqslant_0 \uparrow Q} \quad D0\text{SHIFTU}$$

$$\frac{\Gamma \vdash P \quad \Gamma \vdash [P/a +] N \leqslant_0 M \quad M \neq \forall \beta^+ . M'}{\Gamma \vdash \forall \alpha^+ . N \leqslant_0 M} \quad D0\text{ForallL}$$

$$\frac{\Gamma, \alpha^+ \vdash N \leqslant_0 M}{\Gamma \vdash N \leqslant_0 \forall \alpha^+ . M} \quad D0\text{ForallR}$$

$$\frac{\Gamma \vdash P \geqslant_0 Q \quad \Gamma \vdash N \leqslant_0 M}{\Gamma \vdash P \to N \leqslant_0 Q \to M} \quad D0\text{Arrow}$$

 $\Gamma \vdash P \geqslant_0 Q$ Positive supertyping

$$\frac{\Gamma \vdash a + \geqslant_0 a +}{\Gamma \vdash N \simeq_0^{\leq} M} \quad D0PVAR$$

$$\frac{\Gamma \vdash N \simeq_0^{\leq} M}{\Gamma \vdash \downarrow N \geqslant_0 \downarrow M} \quad D0SHIFTD$$

$$\frac{\Gamma \vdash N \quad \Gamma \vdash [N/a -]P \geqslant_0 Q \quad Q \neq \exists \alpha^-. Q'}{\Gamma \vdash \exists \alpha^-. P \geqslant_0 Q} \quad D0EXISTSL$$

$$\frac{\Gamma, \alpha^- \vdash P \geqslant_0 Q}{\Gamma \vdash P \geqslant_0 \exists \alpha^-. Q} \quad D0EXISTSR$$

 $P_1 \vee P_2$

ord vars in P

ord vars in N

ord vars in P

 $\mathbf{ord} \ vars \mathbf{in} \ N$

 $\mathbf{nf}(N')$

 $\mathbf{nf}(P')$

 $\mathbf{nf}(N')$

 $\mathbf{nf}(P')$

 $e_1 \& e_2$

 $\hat{\sigma}_1 \& \hat{\sigma}_2$

 $\overline{P_1 \vee P_2 = Q}$ Least Upper Bound (Least Common Supertype)

$$\frac{\alpha^{+} \vee \alpha^{+} = \alpha^{+}}{\alpha^{+} \vee \alpha^{+} = \alpha^{+}} \quad \text{LUBVAR}$$

$$\frac{(\mathbf{fv} \, N \cup \mathbf{fv} \, M) \vDash \downarrow N \overset{a}{\simeq} \downarrow M = (P, \hat{\sigma}_{1}, \hat{\sigma}_{2})}{\downarrow N \vee \downarrow M = \exists \alpha^{-}. [\alpha^{-}/(\mathbf{uv} \, P)]P} \quad \text{LUBSHIFT}$$

$$\frac{\{\alpha^{-}\} \cap \{\beta^{-}\} = \emptyset}{\exists \alpha^{-}. P_{1} \vee \exists \beta^{-}. P_{2} = P_{1} \vee P_{2}} \quad \text{LUBEXISTS}$$

 $\Gamma \vDash P_1 \stackrel{a}{\simeq} P_2 = (Q, \hat{\sigma}_1, \hat{\sigma}_2)$

$$\frac{\Gamma \vDash \alpha^{+} \stackrel{a}{\simeq} \alpha^{+} \dashv (\alpha^{+}, \cdot, \cdot)}{\Gamma \vDash N_{1} \stackrel{a}{\simeq} N_{2} \dashv (M, \widehat{\sigma}_{1}, \widehat{\sigma}_{2})} \qquad \text{AUPSHIFT}$$

$$\frac{\Gamma \vDash N_{1} \stackrel{a}{\simeq} N_{2} \dashv (M, \widehat{\sigma}_{1}, \widehat{\sigma}_{2})}{\Gamma \vDash \downarrow N_{1} \stackrel{a}{\simeq} \downarrow N_{2} \dashv (\downarrow M, \widehat{\sigma}_{1}, \widehat{\sigma}_{2})} \qquad \text{AUPSHIFT}$$

$$\frac{\{\overrightarrow{\alpha^{-}}\} \cap \{\Gamma\} = \varnothing \quad \Gamma \vDash P_{1} \stackrel{a}{\simeq} P_{2} \dashv (Q, \widehat{\sigma}_{1}, \widehat{\sigma}_{2})}{\Gamma \vDash \exists \overrightarrow{\alpha^{-}} . P_{1} \stackrel{a}{\simeq} \exists \overrightarrow{\alpha^{-}} . P_{2} \dashv (\exists \overrightarrow{\alpha^{-}} . Q, \widehat{\sigma}_{1}, \widehat{\sigma}_{2})} \qquad \text{AUPEXISTS}$$

 $\Gamma \vDash N_1 \stackrel{a}{\simeq} N_2 \dashv (M, \hat{\sigma}_1, \hat{\sigma}_2)$

$$\frac{\Gamma \vDash \alpha^{-\frac{a}{2}} \alpha^{-} \dashv (\alpha^{-}, \cdot, \cdot)}{\Gamma \vDash P_{1} \stackrel{a}{\simeq} P_{2} \dashv (Q, \hat{\sigma}_{1}, \hat{\sigma}_{2})} \quad \text{AUNSHIFT}$$

$$\frac{\Gamma \vDash P_{1} \stackrel{a}{\simeq} P_{2} \dashv (Q, \hat{\sigma}_{1}, \hat{\sigma}_{2})}{\Gamma \vDash P_{1} \stackrel{a}{\simeq} \uparrow P_{2} \dashv (\uparrow Q, \hat{\sigma}_{1}, \hat{\sigma}_{2})} \quad \text{AUNSHIFT}$$

$$\frac{\Gamma \vDash P_{1} \stackrel{a}{\simeq} P_{2} \dashv (Q, \hat{\sigma}_{1}, \hat{\sigma}_{2}) \quad \Gamma \vDash N_{1} \stackrel{a}{\simeq} N_{2} \dashv (M, \hat{\sigma}'_{1}, \hat{\sigma}'_{2})}{\Gamma \vDash P_{1} \rightarrow N_{1} \stackrel{a}{\simeq} P_{2} \rightarrow N_{2} \dashv (Q \rightarrow M, \hat{\sigma}_{1} \cup \hat{\sigma}'_{1}, \hat{\sigma}_{2} \cup \hat{\sigma}'_{2})} \quad \text{AUNARROW}$$

$$\begin{array}{c} \text{if any other rule is not applicable} \quad \Gamma \vdash N \quad \Gamma \vdash M \\ \hline \Gamma \vdash N \stackrel{\alpha}{\simeq} M \dashv (\widehat{\alpha}_{\{N,M\}}^-, (\widehat{\alpha}_{\{N,M\}}^-) \approx N), (\widehat{\alpha}_{\{N,M\}}^-) \approx M)) \end{array} \quad \text{AUNAU} \\ \hline \mathbf{nf}(N) = M \\ \hline \begin{array}{c} \mathbf{nf}(P) = \alpha \\ \hline \mathbf{nf}(P) = Q \\ \mathbf{nf}(P) = Q \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(N) = M \\ \hline \mathbf{nf}(P) = Q \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(N) = M \\ \hline \mathbf{nf}(P) = Q \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(N) = M \\ \hline \mathbf{nf}(N) = M \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(N) = M \\ \hline \mathbf{nf}(N) = M \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(N) = M \\ \hline \mathbf{nf}(N) = M \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(N) = M \\ \hline \mathbf{nf}(R) = M \\ \hline \end{array} \quad \begin{array}{c} \mathbf{nf}(R) = R \\ \hline \end{array} \quad \begin{array}{c} \mathbf{$$

 $\operatorname{ord} vars \operatorname{in} P = \overrightarrow{\alpha}$

$$\frac{\alpha^+ \in \mathit{vars}}{\mathbf{ord} \, \mathit{vars} \, \mathbf{in} \, \alpha^+ = \alpha^+} \quad \mathrm{OPVarIn}$$

$$\frac{\alpha^{+} \notin vars}{\mathbf{ord} \ vars \mathbf{in} \ \alpha^{+} = \cdot} \quad \text{OPVARNIN}$$

$$\frac{\mathbf{ord} \ vars \mathbf{in} \ N = \overrightarrow{\alpha}}{\mathbf{ord} \ vars \mathbf{in} \ \sqrt{N} = \overrightarrow{\alpha}} \quad \text{OSHIFTD}$$

$$\frac{vars \cap \{\overrightarrow{\alpha^{-}}\} = \varnothing \quad \mathbf{ord} \ vars \mathbf{in} \ P = \overrightarrow{\alpha}}{\mathbf{ord} \ vars \mathbf{in} \ \overrightarrow{\exists \alpha^{-}} . P = \overrightarrow{\alpha}} \quad \text{OEXISTS}$$

 $\mathbf{ord} \ vars \mathbf{in} \ N = \overrightarrow{\alpha}$

$$\frac{1}{\operatorname{ord} \operatorname{varsin} \widehat{\alpha}^{-} \{\operatorname{vars'}\} = \cdot}$$
 ONUVAR

 $\operatorname{ord} vars \operatorname{in} P = \overrightarrow{\alpha}$

$$\frac{1}{\operatorname{ord} \operatorname{vars} \operatorname{in} \widehat{\alpha}^{+} \{ \operatorname{vars'} \} = \cdot} \quad \operatorname{OPUVAR}$$

 $e_1 \& e_2 = e_3$ Unification Solution Entry Merge

$$\overline{(\widehat{\alpha}^{+}:\geqslant P)\ \&\ (\widehat{\alpha}^{+}:\geqslant Q) = (\widehat{\alpha}^{+}:\geqslant P\vee Q)} \quad \text{SMESUPSUP}$$

$$\frac{\mathbf{fv}\ P \cup \mathbf{fv}\ Q \vDash P \geqslant Q \dashv \widehat{\sigma}'}{(\widehat{\alpha}^{+}:\approx P)\ \&\ (\widehat{\alpha}^{+}:\geqslant Q) = (\widehat{\alpha}^{+}:\approx P)} \quad \text{SMEEQSUP}$$

$$\frac{\mathbf{fv}\ P \cup \mathbf{fv}\ Q \vDash Q \geqslant P \dashv \widehat{\sigma}'}{(\widehat{\alpha}^{+}:\geqslant P)\ \&\ (\widehat{\alpha}^{+}:\approx Q) = (\widehat{\alpha}^{+}:\approx Q)} \quad \text{SMESUPEQ}$$

$$\overline{(\widehat{\alpha}^{+}:\approx P)\ \&\ (\widehat{\alpha}^{+}:\approx P) = (\widehat{\alpha}^{+}:\approx P)} \quad \text{SMEPEQEQ}$$

$$\overline{(\widehat{\alpha}^{-}:\approx N)\ \&\ (\widehat{\alpha}^{-}:\approx N) = (\widehat{\alpha}^{-}:\approx N)} \quad \text{SMENEQEQ}$$

 $\begin{array}{|c|c|} \hline \widehat{\sigma}_1 \& \widehat{\sigma}_2 = \widehat{\sigma}_3 \\ \hline N \stackrel{u}{\simeq} M = \widehat{\sigma} \\ \hline \end{array}$ Merge unification solutions

$$\frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-}} \stackrel{\text{UNVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-}} \stackrel{\text{UNVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-}} \stackrel{\text{USHIFTU}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{USHIFTU}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UARROW}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UARROW}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UARROW}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UFORALL}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UNUVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UNUVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UNUVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UNUVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}\alpha^{-\frac{u}{2}}} \stackrel{\text{UNUVAR}}{\longrightarrow} \frac{1}{\alpha^{-\frac{u}{2}}\alpha^{-\frac{u$$

 $P \stackrel{u}{\simeq} Q = \hat{\sigma}$ Positive unification

$$\frac{-\frac{u}{\alpha^{+}} \overset{u}{\simeq} \alpha^{+} \dashv \cdot}{\frac{N \overset{u}{\simeq} M \dashv \hat{\sigma}}{\downarrow N \overset{u}{\simeq} \downarrow M \dashv \hat{\sigma}}} \quad \text{USHIFTD}$$

$$\frac{P\overset{u}{\simeq}Q \dashv \hat{\sigma}}{\exists \overrightarrow{\alpha^{-}}.P\overset{u}{\simeq}\exists \overrightarrow{\alpha^{-}}.Q \dashv \hat{\sigma}} \quad \text{UEXISTS}$$

$$\frac{\mathbf{fv}\,P \subseteq vars}{\widehat{\alpha}^{+}\{vars\}\overset{u}{\simeq}P \dashv \widehat{\alpha}^{+}:\approx P} \quad \text{UPUVAR}$$

 $\begin{array}{|c|c|c|c|}\hline \Gamma \vdash N & \text{Negative type well-formedness} \\ \hline \Gamma \vdash P & \text{Positive type well-formedness} \\ \hline \Gamma \vdash N & \text{Negative type well-formedness} \\ \hline \Gamma \vdash P & \text{Positive type well-formedness} \\ \end{array}$

Definition rules: 88 good 0 bad Definition rule clauses: 150 good 0 bad