

1 The Vanilla System

First, we present the top-level system, which is easy to understand.

1.1 Grammar

$P, Q ::=$ positive types

- α^+
- $\downarrow N$
- $\exists \alpha^-. P$

$N, M ::=$ negative types

- α^-
- $\uparrow P$
- $\forall \alpha^+. N$
- $P \rightarrow N$

1.2 Declarative Subtyping

$\boxed{\Gamma \vdash N \simeq_0^{\leq} M}$ Negative equivalence

$$\frac{\Gamma \vdash N \leq_0 M \quad \Gamma \vdash M \leq_0 N}{\Gamma \vdash N \simeq_0^{\leq} M} \text{ D0NDEF}$$

$\boxed{\Gamma \vdash P \simeq_0^{\leq} Q}$ Positive equivalence

$$\frac{\Gamma \vdash P \geq_0 Q \quad \Gamma \vdash Q \geq_0 P}{\Gamma \vdash P \simeq_0^{\leq} Q} \text{ D0PDEF}$$

$\boxed{\Gamma \vdash N \leq_0 M}$ Negative subtyping

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^- \leq_0 \alpha^-} \quad \text{D0NVAR} \\ \frac{\Gamma \vdash P \simeq_0^{\leq} Q}{\Gamma \vdash \uparrow P \leq_0 \uparrow Q} \quad \text{D0SHIFTU} \\ \frac{\Gamma \vdash P \quad \Gamma \vdash [P/\alpha^+]N \leq_0 M \quad M \neq \forall \beta^+. M'}{\Gamma \vdash \forall \alpha^+. N \leq_0 M} \quad \text{D0FORALLL} \\ \frac{\Gamma, \alpha^+ \vdash N \leq_0 M}{\Gamma \vdash N \leq_0 \forall \alpha^+. M} \quad \text{D0FORALLR} \\ \frac{\Gamma \vdash P \geq_0 Q \quad \Gamma \vdash N \leq_0 M}{\Gamma \vdash P \rightarrow N \leq_0 Q \rightarrow M} \quad \text{D0ARROW} \end{array}$$

$\boxed{\Gamma \vdash P \geq_0 Q}$ Positive supertyping

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^+ \geq_0 \alpha^+} \quad \text{D0PVAR} \\ \frac{\Gamma \vdash N \simeq_0^{\leq} M}{\Gamma \vdash \downarrow N \geq_0 \downarrow M} \quad \text{D0SHIFTD} \\ \frac{\Gamma \vdash N \quad \Gamma \vdash [N/\alpha^-]P \geq_0 Q \quad Q \neq \exists \alpha^-. Q'}{\Gamma \vdash \exists \alpha^-. P \geq_0 Q} \quad \text{D0EXISTSL} \\ \frac{\Gamma, \alpha^- \vdash P \geq_0 Q}{\Gamma \vdash P \geq_0 \exists \alpha^-. Q} \quad \text{D0EXISTSR} \end{array}$$

2 Multi-Quantified System

2.1 Grammar

P, Q	$::=$	multi-quantified positive types
	α^+	
	$\downarrow N$	
	$\exists \alpha^+. P$	$P \neq \exists \dots$
	(P)	S
N, M	$::=$	multi-quantified negative types
	α^-	
	$\uparrow P$	
	$P \rightarrow N$	
	$\forall \alpha^+. N$	$N \neq \forall \dots$
	(N)	S

2.2 Declarative Subtyping

$\boxed{\Gamma \vdash N \simeq_1^< M}$ Negative equivalence on MQ types

$$\frac{\Gamma \vdash N \leq_1 M \quad \Gamma \vdash M \leq_1 N}{\Gamma \vdash N \simeq_1^< M} \quad \text{D1NDEF}$$

$\boxed{\Gamma \vdash P \simeq_1^< Q}$ Positive equivalence on MQ types

$$\frac{\Gamma \vdash P \geq_1 Q \quad \Gamma \vdash Q \geq_1 P}{\Gamma \vdash P \simeq_1^< Q} \quad \text{D1PDEF}$$

$\boxed{\Gamma \vdash N \leq_1 M}$ Negative subtyping

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^- \leq_1 \alpha^-} \quad (\text{VAR}^{-\leq_1}) \\ \frac{\Gamma \vdash P \simeq_1^< Q}{\Gamma \vdash \uparrow P \leq_1 \uparrow Q} \quad (\uparrow^{\leq_1}) \\ \frac{\Gamma \vdash P \geq_1 Q \quad \Gamma \vdash N \leq_1 M}{\Gamma \vdash P \rightarrow N \leq_1 Q \rightarrow M} \quad (\rightarrow^{\leq_1}) \\ \frac{\text{fv } N \cap \{\vec{\beta}^+\} = \emptyset \quad \Gamma, \vec{\beta}^+ \vdash P_i \quad \Gamma, \vec{\beta}^+ \vdash [\vec{P}/\alpha^+] N \leq_1 M}{\Gamma \vdash \forall \alpha^+. N \leq_1 \forall \vec{\beta}^+. M} \quad (\forall^{\leq_1}) \end{array}$$

$\boxed{\Gamma \vdash P \geq_1 Q}$ Positive supertyping

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^+ \geq_1 \alpha^+} \quad (\text{VAR}^{+\geq_1}) \\ \frac{\Gamma \vdash N \simeq_1^< M}{\Gamma \vdash \downarrow N \geq_1 \downarrow M} \quad (\downarrow^{\geq_1}) \\ \frac{\text{fv } P \cap \{\vec{\beta}^-\} = \emptyset \quad \Gamma, \vec{\beta}^- \vdash N_i \quad \Gamma, \vec{\beta}^- \vdash [\vec{N}/\alpha^-] P \geq_1 Q}{\Gamma \vdash \exists \alpha^-. P \geq_1 \exists \vec{\beta}^-. Q} \quad (\exists^{\geq_1}) \end{array}$$

2.3 Declarative Equivalence

$\boxed{N \simeq_1^D M}$ Negative multi-quantified type equivalence

$$\begin{array}{c} \overline{\alpha^- \simeq_1^D \alpha^-} \quad (\text{VAR}^{-\simeq_1^D}) \\ \frac{P \simeq_1^D Q}{\uparrow P \simeq_1^D \uparrow Q} \quad (\uparrow^{\simeq_1^D}) \\ \frac{P \simeq_1^D Q \quad N \simeq_1^D M}{P \rightarrow N \simeq_1^D Q \rightarrow M} \quad (\rightarrow^{\simeq_1^D}) \end{array}$$

$$\frac{\{\vec{\alpha}^+\} \cap \mathbf{fv} M = \emptyset \quad \mu : (\{\vec{\beta}^+\} \cap \mathbf{fv} M) \leftrightarrow (\{\vec{\alpha}^+\} \cap \mathbf{fv} N) \quad N \simeq_1^D [\mu]M}{\forall \vec{\alpha}^+. N \simeq_1^D \forall \vec{\beta}^+. M} \quad (\forall \simeq_1^D)$$

$\boxed{P \simeq_1^D Q}$ Positive multi-quantified type equivalence

$$\frac{\overline{\alpha^+ \simeq_1^D \alpha^+} \quad (\text{VAR}^+) \quad \frac{N \simeq_1^D M}{\downarrow N \simeq_1^D \downarrow M} \quad (\downarrow \simeq_1^D)}{\frac{\{\vec{\alpha}^-\} \cap \mathbf{fv} Q = \emptyset \quad \mu : (\{\vec{\beta}^-\} \cap \mathbf{fv} Q) \leftrightarrow (\{\vec{\alpha}^-\} \cap \mathbf{fv} P) \quad P \simeq_1^D [\mu]Q}{\exists \vec{\alpha}^-. P \simeq_1^D \exists \vec{\beta}^-. Q}} \quad (\exists \simeq_1^D)$$

$\boxed{P \simeq Q}$

3 Algorithm

3.1 Normalization

3.1.1 Ordering

$\boxed{\text{ord vars in } N = \vec{\alpha}}$

$$\frac{\alpha^- \in \text{vars}}{\text{ord vars in } \alpha^- = \alpha^-} \quad (\text{VAR}_{\in}^-)$$

$$\frac{\alpha^- \notin \text{vars}}{\text{ord vars in } \alpha^- = \cdot} \quad (\text{VAR}_{\notin}^-)$$

$$\frac{\text{ord vars in } P = \vec{\alpha}}{\text{ord vars in } \uparrow P = \vec{\alpha}} \quad (\uparrow)$$

$$\frac{\text{ord vars in } P = \vec{\alpha}_1 \quad \text{ord vars in } N = \vec{\alpha}_2}{\text{ord vars in } P \rightarrow N = \vec{\alpha}_1, (\vec{\alpha}_2 \setminus \{\vec{\alpha}_1\})} \quad (\rightarrow)$$

$$\frac{\text{vars} \cap \{\vec{\alpha}^+\} = \emptyset \quad \text{ord vars in } N = \vec{\alpha}}{\text{ord vars in } \forall \vec{\alpha}^+. N = \vec{\alpha}} \quad (\forall)$$

$\boxed{\text{ord vars in } P = \vec{\alpha}}$

$$\frac{\alpha^+ \in \text{vars}}{\text{ord vars in } \alpha^+ = \alpha^+} \quad (\text{VAR}_{\in}^+)$$

$$\frac{\alpha^+ \notin \text{vars}}{\text{ord vars in } \alpha^+ = \cdot} \quad (\text{VAR}_{\notin}^+)$$

$$\frac{\text{ord vars in } N = \vec{\alpha}}{\text{ord vars in } \downarrow N = \vec{\alpha}} \quad (\downarrow)$$

$$\frac{\text{vars} \cap \{\vec{\alpha}^-\} = \emptyset \quad \text{ord vars in } P = \vec{\alpha}}{\text{ord vars in } \exists \vec{\alpha}^-. P = \vec{\alpha}} \quad (\exists)$$

$\boxed{\text{ord vars in } N = \vec{\alpha}}$

$$\frac{}{\text{ord vars in } \hat{\alpha}^- = \cdot} \quad (\text{UVar}^-)$$

$\boxed{\text{ord vars in } P = \vec{\alpha}}$

$$\frac{}{\text{ord vars in } \hat{\alpha}^+ = \cdot} \quad (\text{UVar}^+)$$

3.1.2 Quantifier Normalization

$$\boxed{\mathbf{nf}(N) = M}$$

$$\begin{array}{c} \overline{\mathbf{nf}(\alpha^-) = \alpha^-} \quad (\text{VAR}^-) \\ \frac{\mathbf{nf}(P) = Q}{\mathbf{nf}(\uparrow P) = \uparrow Q} \quad (\uparrow) \\ \frac{\mathbf{nf}(P) = Q \quad \mathbf{nf}(N) = M}{\mathbf{nf}(P \rightarrow N) = Q \rightarrow M} \quad (\rightarrow) \\ \frac{\mathbf{nf}(N) = N' \quad \mathbf{ord}\{\overrightarrow{\alpha^+}\} \text{ in } N' = \overrightarrow{\alpha^{+'}}}{\mathbf{nf}(\forall \overrightarrow{\alpha^+}.N) = \forall \overrightarrow{\alpha^{+'}}.N'} \quad (\forall) \end{array}$$

$$\boxed{\mathbf{nf}(P) = Q}$$

$$\begin{array}{c} \overline{\mathbf{nf}(\alpha^+) = \alpha^+} \quad (\text{VAR}^+) \\ \frac{\mathbf{nf}(N) = M}{\mathbf{nf}(\downarrow N) = \downarrow M} \quad (\downarrow) \\ \frac{\mathbf{nf}(P) = P' \quad \mathbf{ord}\{\overrightarrow{\alpha^-}\} \text{ in } P' = \overrightarrow{\alpha^{-'}}}{\mathbf{nf}(\exists \overrightarrow{\alpha^-}.P) = \exists \overrightarrow{\alpha^{-'}}.P'} \quad (\exists) \end{array}$$

$$\boxed{\mathbf{nf}(N) = M}$$

$$\boxed{\mathbf{nf}(P) = Q}$$

$$\overline{\mathbf{nf}(\hat{\alpha}^-) = \hat{\alpha}^-} \quad (\text{UVAR}^-)$$

$$\overline{\mathbf{nf}(\hat{\alpha}^+) = \hat{\alpha}^+} \quad (\text{UVAR}^+)$$

3.2 Unification

$$\boxed{\Theta \models N \stackrel{u}{\simeq} M \Rightarrow \hat{\sigma}} \quad \text{Negative unification}$$

$$\begin{array}{c} \overline{\Theta \models \alpha^- \stackrel{u}{\simeq} \alpha^- \Rightarrow \cdot} \quad \text{UNVAR} \\ \frac{\Theta \models P \stackrel{u}{\simeq} Q \Rightarrow \hat{\sigma}}{\Theta \models \uparrow P \stackrel{u}{\simeq} \uparrow Q \Rightarrow \hat{\sigma}} \quad \text{USHIFTU} \\ \frac{\Theta \models P \stackrel{u}{\simeq} Q \Rightarrow \hat{\sigma}_1 \quad \Theta \models N \stackrel{u}{\simeq} M \Rightarrow \hat{\sigma}_2}{\Theta \models P \rightarrow N \stackrel{u}{\simeq} Q \rightarrow M \Rightarrow \hat{\sigma}_1 \ \& \ \hat{\sigma}_2} \quad \text{UARROW} \\ \frac{\Theta \models N \stackrel{u}{\simeq} M \Rightarrow \hat{\sigma}}{\Theta \models \forall \overrightarrow{\alpha^+}.N \stackrel{u}{\simeq} \forall \overrightarrow{\alpha^+}.M \Rightarrow \hat{\sigma}} \quad \text{UFORALL} \\ \frac{\hat{\alpha}^-\{\Delta\} \in \Theta \quad \Delta \vdash N}{\Theta \models \hat{\alpha}^- \stackrel{u}{\simeq} N \Rightarrow (\Delta \vdash \hat{\alpha}^- : \approx N)} \quad \text{UNUVAR} \end{array}$$

$$\boxed{\Theta \models P \stackrel{u}{\simeq} Q \Rightarrow \hat{\sigma}} \quad \text{Positive unification}$$

$$\begin{array}{c} \overline{\Theta \models \alpha^+ \stackrel{u}{\simeq} \alpha^+ \Rightarrow \cdot} \quad \text{UPVAR} \\ \frac{\Theta \models N \stackrel{u}{\simeq} M \Rightarrow \hat{\sigma}}{\Theta \models \downarrow N \stackrel{u}{\simeq} \downarrow M \Rightarrow \hat{\sigma}} \quad \text{USHIFTD} \\ \frac{\Theta \models P \stackrel{u}{\simeq} Q \Rightarrow \hat{\sigma}}{\Theta \models \exists \overrightarrow{\alpha^-}.P \stackrel{u}{\simeq} \exists \overrightarrow{\alpha^-}.Q \Rightarrow \hat{\sigma}} \quad \text{UEXISTS} \\ \frac{\hat{\alpha}^+\{\Delta\} \in \Theta \quad \Delta \vdash P}{\Theta \models \hat{\alpha}^+ \stackrel{u}{\simeq} P \Rightarrow (\Delta \vdash \hat{\alpha}^+ : \approx P)} \quad \text{UPUVAR} \end{array}$$

3.3 Algorithmic Subtyping

$\boxed{\Gamma; \Theta \models N \leq M \Rightarrow \hat{\sigma}}$ Negative subtyping

$$\begin{array}{c}
\overline{\Gamma; \Theta \models \alpha^- \leq \alpha^- \Rightarrow \cdot} \quad \text{ANVAR} \\
\\
\frac{\Theta \models \mathbf{nf}(P) \stackrel{u}{\simeq} \mathbf{nf}(Q) \Rightarrow \hat{\sigma}}{\Gamma; \Theta \models \uparrow P \leq \uparrow Q \Rightarrow \hat{\sigma}} \quad \text{AShiftU} \\
\\
\frac{\Gamma; \Theta \models P \geq Q \Rightarrow \hat{\sigma}_1 \quad \Gamma; \Theta \models N \leq M \Rightarrow \hat{\sigma}_2}{\Gamma; \Theta \models P \rightarrow N \leq Q \rightarrow M \Rightarrow \hat{\sigma}_1 \& \hat{\sigma}_2} \quad \text{AArrow} \\
\\
\frac{\Gamma, \vec{\beta}^+; \Theta, \vec{\alpha}^+ \{ \Gamma, \vec{\beta}^+ \} \models [\vec{\alpha}^+ / \alpha^+] N \leq M \Rightarrow \hat{\sigma}}{\Gamma; \Theta \models \forall \alpha^+. N \leq \forall \beta^+. M \Rightarrow \hat{\sigma} \setminus \vec{\alpha}^+} \quad \text{AForall}
\end{array}$$

$\boxed{\Gamma; \Theta \models P \geq Q \Rightarrow \hat{\sigma}}$ Positive supertyping

$$\begin{array}{c}
\overline{\Gamma; \Theta \models \alpha^+ \geq \alpha^+ \Rightarrow \cdot} \quad \text{APVAR} \\
\\
\frac{\Theta \models \mathbf{nf}(N) \stackrel{u}{\simeq} \mathbf{nf}(M) \Rightarrow \hat{\sigma}}{\Gamma; \Theta \models \downarrow N \geq \downarrow M \Rightarrow \hat{\sigma}} \quad \text{AShiftD} \\
\\
\frac{\Gamma, \vec{\beta}^-; \Theta, \vec{\alpha}^- \{ \Gamma, \vec{\beta}^- \} \models [\vec{\alpha}^- / \alpha^-] P \geq Q \Rightarrow \hat{\sigma}}{\Gamma; \Theta \models \exists \alpha^-. P \geq \exists \beta^-. Q \Rightarrow \hat{\sigma}} \quad \text{AExists} \\
\\
\frac{\text{upgrade } \Gamma \vdash \mathbf{nf}(P) \text{ to } \Delta = Q}{\Gamma; \Theta \models \hat{\alpha}^+ \{ \Delta \} \geq P \Rightarrow (\Delta \vdash \hat{\alpha}^+ : \geq Q)} \quad \text{APUVar}
\end{array}$$

3.4 Unification Solution Merge

Unification solution is represented by a list of unification solution entries. Each entry restrict an unification variable in two possible ways: either stating that it must be equivalent to a certain type ($\Delta \vdash \hat{\alpha}^+ : \approx P$ or $\Delta \vdash \hat{\alpha}^- : \approx N$) or that it must be a (positive) supertype of a certain type ($\Delta \vdash \hat{\alpha}^+ : \geq P$).

Definition 1 (Matching Entries). *We call two entries matching if they are restricting the same unification variable.*

Two matching entries can be merged in the following way:

Definition 2.

$\boxed{e_1 \& e_2 = e_3}$ Unification Solution Entry Merge

$$\begin{array}{c}
\frac{\Gamma \vdash P_1 \vee P_2 = Q}{(\Gamma \vdash \hat{\alpha}^+ : \geq P_1) \& (\Gamma \vdash \hat{\alpha}^+ : \geq P_2) = (\Gamma \vdash \hat{\alpha}^+ : \geq Q)} \quad (\geq \& \geq) \\
\\
\frac{\Gamma; \cdot \models P \geq Q \Rightarrow \hat{\sigma}'}{(\Gamma \vdash \hat{\alpha}^+ : \approx P) \& (\Gamma \vdash \hat{\alpha}^+ : \geq Q) = (\Gamma \vdash \hat{\alpha}^+ : \approx P)} \quad (\simeq \& \geq) \\
\\
\frac{\Gamma; \cdot \models Q \geq P \Rightarrow \hat{\sigma}'}{(\Gamma \vdash \hat{\alpha}^+ : \geq P) \& (\Gamma \vdash \hat{\alpha}^+ : \approx Q) = (\Gamma \vdash \hat{\alpha}^+ : \approx Q)} \quad (\geq \& \simeq) \\
\\
\frac{}{(\Gamma \vdash \hat{\alpha}^+ : \approx P) \& (\Gamma \vdash \hat{\alpha}^+ : \approx P) = (\Gamma \vdash \hat{\alpha}^+ : \approx P)} \quad (\simeq \& \simeq^+) \\
\\
\frac{}{(\Gamma \vdash \hat{\alpha}^- : \approx N) \& (\Gamma \vdash \hat{\alpha}^- : \approx N) = (\Gamma \vdash \hat{\alpha}^- : \approx N)} \quad (\simeq \& \simeq^-)
\end{array}$$

To merge two unification solution, we merge each pair of matching entries, and unite the results.

Definition 3. $\hat{\sigma}_1 \& \hat{\sigma}_2 = \{e_1 \& e_2 \mid e_1 \in \hat{\sigma}_1, e_2 \in \hat{\sigma}_2, \text{ s.t. } e_1 \text{ matches with } e_2\}$
 $\cup \{e_1 \mid e_1 \in \hat{\sigma}_1, \text{ s.t. } \forall e_2 \in \hat{\sigma}_2, e_1 \text{ does not match with } e_2\}$
 $\cup \{e_2 \mid e_2 \in \hat{\sigma}_2, \text{ s.t. } \forall e_1 \in \hat{\sigma}_1, e_2 \text{ does not match with } e_2\}$

3.5 Least Upper Bound

$\boxed{\Gamma \vdash P_1 \vee P_2 = Q}$ Least Upper Bound (Least Common Supertype)

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^+ \vee \alpha^+ = \alpha^+} \quad (\text{VAR}^\vee) \\[10pt] \frac{\Gamma, \cdot \vdash \downarrow N \stackrel{a}{\simeq} \downarrow M \Rightarrow (\Xi, P, \hat{\tau}_1, \hat{\tau}_2)}{\Gamma \vdash \downarrow N \vee \downarrow M = \exists \alpha^-. [\overrightarrow{\alpha^-} / \Xi] P} \quad (\downarrow^\vee) \\[10pt] \frac{\Gamma, \overrightarrow{\alpha^-}, \overrightarrow{\beta^-} \vdash P_1 \vee P_2 = Q}{\Gamma \vdash \exists \overrightarrow{\alpha^-}. P_1 \vee \exists \overrightarrow{\beta^-}. P_2 = Q} \quad (\exists^\vee) \end{array}$$

$\boxed{\text{upgrade} \Gamma \vdash P \text{ to } \Delta = Q}$

3.6 Antiunification

$\boxed{\Gamma \vdash P_1 \stackrel{a}{\simeq} P_2 \Rightarrow (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2)}$

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^+ \stackrel{a}{\simeq} \alpha^+ \Rightarrow (\cdot, \alpha^+, \cdot, \cdot)} \quad \text{AUPVAR} \\[10pt] \frac{\Gamma \vdash N_1 \stackrel{a}{\simeq} N_2 \Rightarrow (\Xi, M, \hat{\tau}_1, \hat{\tau}_2)}{\Gamma \vdash \downarrow N_1 \stackrel{a}{\simeq} \downarrow N_2 \Rightarrow (\Xi, \downarrow M, \hat{\tau}_1, \hat{\tau}_2)} \quad \text{AUPSHIFT} \\[10pt] \frac{\{\overrightarrow{\alpha^-}\} \cap \{\Gamma\} = \emptyset \quad \Gamma \vdash P_1 \stackrel{a}{\simeq} P_2 \Rightarrow (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2)}{\Gamma \vdash \exists \overrightarrow{\alpha^-}. P_1 \stackrel{a}{\simeq} \exists \overrightarrow{\alpha^-}. P_2 \Rightarrow (\Xi, \exists \overrightarrow{\alpha^-}. Q, \hat{\tau}_1, \hat{\tau}_2)} \quad \text{AUPEXISTS} \end{array}$$

$\boxed{\Gamma \vdash N_1 \stackrel{a}{\simeq} N_2 \Rightarrow (\Xi, M, \hat{\tau}_1, \hat{\tau}_2)}$

$$\begin{array}{c} \overline{\Gamma \vdash \alpha^- \stackrel{a}{\simeq} \alpha^- \Rightarrow (\Xi, \alpha^-, \cdot, \cdot)} \quad \text{AUNVAR} \\[10pt] \frac{\Gamma \vdash P_1 \stackrel{a}{\simeq} P_2 \Rightarrow (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2)}{\Gamma \vdash \uparrow P_1 \stackrel{a}{\simeq} \uparrow P_2 \Rightarrow (\Xi, \uparrow Q, \hat{\tau}_1, \hat{\tau}_2)} \quad \text{AUNSHIFT} \\[10pt] \frac{\Gamma \vdash P_1 \stackrel{a}{\simeq} P_2 \Rightarrow (\Xi_1, Q, \hat{\tau}_1, \hat{\tau}_2) \quad \Gamma \vdash N_1 \stackrel{a}{\simeq} N_2 \Rightarrow (\Xi_2, M, \hat{\tau}'_1, \hat{\tau}'_2)}{\Gamma \vdash P_1 \rightarrow N_1 \stackrel{a}{\simeq} P_2 \rightarrow N_2 \Rightarrow (\Xi_1 \cup \Xi_2, Q \rightarrow M, \hat{\tau}_1 \cup \hat{\tau}'_1, \hat{\tau}_2 \cup \hat{\tau}'_2)} \quad \text{AUNARROW} \\[10pt] \frac{\text{if any other rule is not applicable} \quad \Gamma \vdash N \quad \Gamma \vdash M}{\Gamma \vdash N \stackrel{a}{\simeq} M \Rightarrow (\hat{\alpha}_{\{N,M\}}^-, \hat{\alpha}_{\{N,M\}}^-, (\hat{\alpha}_{\{N,M\}}^- : \approx N), (\hat{\alpha}_{\{N,M\}}^- : \approx M))} \quad \text{AUNAU} \end{array}$$

4 Proofs

4.1 Declarative Subtyping

Lemma 1 (Free Variable Propagation). *In the judgments of negative subtyping or positive supertyping, free variables propagate left-to-right. For a context Γ ,*

- $-$ if $\Gamma \vdash N \leqslant_1 M$ then $\mathbf{fv}(N) \subseteq \mathbf{fv}(M)$
- $+$ if $\Gamma \vdash P \geqslant_1 Q$ then $\mathbf{fv}(P) \subseteq \mathbf{fv}(Q)$

Proof. Mutual induction on $\Gamma \vdash N \leqslant_1 M$ and $\Gamma \vdash P \geqslant_1 Q$.

Case 1. $\Gamma \vdash \alpha^- \leqslant_1 \alpha^-$

It is self-evident that $\{\alpha^-\} \subseteq \{\alpha^-\}$.

Case 2. $\Gamma \vdash \uparrow P \leqslant_1 \uparrow Q$ From the inversion (and unfolding $\Gamma \vdash P \simeq_1^< Q$), we have $\Gamma \vdash P \geqslant_1 Q$. Then by the induction hypothesis, $\mathbf{fv}(P) \subseteq \mathbf{fv}(Q)$. The desired inclusion holds, since $\mathbf{fv}(\uparrow P) = \mathbf{fv}(P)$ and $\mathbf{fv}(\uparrow Q) = \mathbf{fv}(Q)$.

Case 3. $\Gamma \vdash P \rightarrow N \leqslant_1 Q \rightarrow M$ The induction hypothesis applied to the premises gives: $\mathbf{fv}(P) \subseteq \mathbf{fv}(Q)$ and $\mathbf{fv}(N) \subseteq \mathbf{fv}(M)$. Then $\mathbf{fv}(P \rightarrow N) = \mathbf{fv}(P) \cup \mathbf{fv}(N) \subseteq \mathbf{fv}(Q) \cup \mathbf{fv}(M) = \mathbf{fv}(Q \rightarrow M)$.

Case 4. $\Gamma \vdash \forall \alpha^+ . N \leq_1 \forall \beta^+ . M$
 $\mathbf{fv} \forall \alpha^+ . N \subseteq \mathbf{fv} ([\vec{P}/\alpha^+]N) \setminus \{\beta^+\}$ here $\{\beta^+\}$ is excluded by the premise $\mathbf{fv} N \cap \{\beta^+\} = \emptyset$
 $\subseteq \mathbf{fv} M \setminus \{\beta^+\}$ by the induction hypothesis, $\mathbf{fv} ([\vec{P}/\alpha^+]N) \subseteq \mathbf{fv} M$
 $\subseteq \mathbf{fv} \forall \beta^+ . M$

Case 5. The positive cases are symmetric. □

4.2 Substitution

Lemma 2 (Substitution strengthening). *Restricting the substitution to the free variables of the substitution subject does not affect the result. Suppose that $\Gamma_2 \vdash \sigma : \Gamma_1$. Then*

- + if $\Gamma_1 \vdash P$ then $[\sigma]P = [\sigma|_{\mathbf{fv} P}]P$,
- if $\Gamma_1 \vdash N$ then $[\sigma]N = [\sigma|_{\mathbf{fv} N}]N$

Proof. Ilya: todo □

4.3 Type well-formedness

Lemma 3 (Well-formedness agrees with substitution). *Suppose that $\Gamma_1 \vdash \sigma : \Gamma_2$. Then*

- + $\Gamma_1, \Gamma_2 \vdash P \Leftrightarrow \Gamma_1 \vdash [\sigma]P$
- $\Gamma_1, \Gamma_2 \vdash N \Leftrightarrow \Gamma_1 \vdash [\sigma]N$

Lemma 4 (Equivalent Contexts). *In the well-formedness judgment, only used variables matter:*

- + if $\{\Gamma_1\} \cap \mathbf{fv} P = \{\Gamma_2\} \cap \mathbf{fv} P$ then $\Gamma_1 \vdash P \iff \Gamma_2 \vdash P$,
- if $\{\Gamma_1\} \cap \mathbf{fv} N = \{\Gamma_2\} \cap \mathbf{fv} N$ then $\Gamma_1 \vdash N \iff \Gamma_2 \vdash N$.

Proof. By simple mutual induction on P and Q . □

4.4 Overview

Algorithm	Soundness	Completeness	Initiality
Ordering	$\overline{\{\mathbf{ord\ vars\ in\ } N\} \equiv \mathbf{vars} \cap \mathbf{fv\ } N}$	$\frac{N \simeq_1^D M}{\mathbf{ord\ vars\ in\ } N = \mathbf{ord\ vars\ in\ } M}$	—
Normalization	$\overline{N \simeq_1^D \mathbf{nf}(N)}$	$\frac{N \simeq_1^D M}{\mathbf{nf}(N) = \mathbf{nf}(M)}$	—
Equivalence	$\frac{\Gamma \vdash P \quad \Gamma \vdash Q \quad P \simeq_1^D Q}{\Gamma \vdash P \simeq_1^{\leq} Q}$	$\frac{\Gamma \vdash P \simeq_1^{\leq} Q}{P \simeq_1^D Q}$	—
Uppgrade	$\frac{\mathbf{upgrade}\ \Gamma \vdash P \mathbf{to}\ \Delta = Q}{Q \text{ is sound } \left\{ \begin{array}{l} \Delta \vdash Q \\ \Gamma \vdash Q \geq_1 P \end{array} \right.}$	$\frac{\exists Q \text{ s.t. } \mathbf{upgrade}\ \Gamma \vdash P \mathbf{to}\ \Delta = Q}{\exists Q \text{ s.t. } \mathbf{upgrade}\ \Gamma \vdash P \mathbf{to}\ \Delta = Q}$	$\frac{Q' \text{ is sound } \quad \mathbf{upgrade}\ \Gamma \vdash P \mathbf{to}\ \Delta = Q}{\Delta \vdash Q' \geq_1 Q}$
LUB	$\frac{\Gamma \models P_1 \vee P_2 = Q}{Q \text{ is sound } \left\{ \begin{array}{l} \Gamma \vdash Q \\ \Gamma \vdash Q \geq_1 P_1 \\ \Gamma \vdash Q \geq_1 P_2 \end{array} \right.}$	$\frac{\exists \text{ sound } Q'}{\exists Q \text{ s.t. } \Gamma \models P_1 \vee P_2 = Q}$	$\frac{Q' \text{ is sound } \quad \Gamma \models P_1 \vee P_2 = Q}{\Delta \vdash Q' \geq_1 Q}$
Anti-unification	$\frac{\Gamma \models P_1 \simeq P_2 \Rightarrow (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2)}{(\Xi, Q, \hat{\tau}_1, \hat{\tau}_2) \left\{ \begin{array}{l} \Xi \text{ is negative} \\ \Gamma; \Xi \vdash Q \\ \Gamma; \cdot \vdash \hat{\tau}_i : \Xi \\ [\hat{\tau}_i] Q = P_i \end{array} \right. \text{ is sound}}$	$\frac{\exists \text{ sound } (\Xi', Q', \hat{\tau}'_1, \hat{\tau}'_2)}{\exists (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2) \text{ s.t. } \Gamma \models P_1 \simeq P_2 \Rightarrow (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2)}$	$\frac{(\Xi', Q', \hat{\tau}'_1, \hat{\tau}'_2) \text{ is sound } \quad \Gamma \models P_1 \simeq P_2 \Rightarrow (\Xi, Q, \hat{\tau}_1, \hat{\tau}_2)}{\exists \Gamma; \Xi \vdash \hat{\tau} : \Xi' \text{ s.t. } [\hat{\tau}] Q' = Q}$
Unification (matching)	$\frac{\Theta \models P \stackrel{u}{\simeq} Q \Rightarrow \hat{\sigma}}{\hat{\sigma} \text{ is sound } \left\{ \begin{array}{l} \Theta \vdash \hat{\sigma} \\ [\hat{\sigma}] P = Q \end{array} \right.}$	$\frac{\exists \text{ sound } \hat{\sigma}'}{\exists \hat{\sigma} \text{ s.t. } \Theta \models P \stackrel{u}{\simeq} Q \Rightarrow \hat{\sigma}}$	—
Subtyping	$\frac{\Gamma; \Theta \models N \leq M \Rightarrow \hat{\sigma}}{\hat{\sigma} \text{ is sound } \left\{ \begin{array}{l} \Theta \vdash \hat{\sigma} \\ \Gamma \vdash [\hat{\sigma}] N \leq_1 M \end{array} \right.}$	$\frac{\exists \text{ sound } \hat{\sigma}'}{\exists \hat{\sigma} \text{ s.t. } \Gamma; \Theta \models N \leq M \Rightarrow \hat{\sigma}}$	—

4.5 Anti-unification

Lemma 5 (Soundness of the anti-unification algorithm).

Lemma 6 (Completeness of the anti-unification algorithm).

Lemma 7 (Initiality of the anti-unification algorithm).

4.6 Variable Ordering

Definition 4 (Collision free bijection). *We say that a bijection $\mu : A \leftrightarrow B$ between sets of variables is **collision free on sets P and Q** if and only if*

1. $\mu(P \cap A) \cap Q = \emptyset$
2. $\mu(Q \cap A) \cap P = \emptyset$

Lemma 8 (Soundness of variable ordering). *Variable ordering extracts precisely used free variables.*

- $\{\mathbf{ord\ vars\ in\ } N\} \equiv \mathbf{vars} \cap \mathbf{fv\ } N$ (as sets)
- + $\{\mathbf{ord\ vars\ in\ } P\} \equiv \mathbf{vars} \cap \mathbf{fv\ } P$ (as sets)

Proof. Straightforward mutual induction on $\mathbf{ord\ vars\ in\ } N = \vec{\alpha}$ and $\mathbf{ord\ vars\ in\ } P = \vec{\alpha}$ □

Corollary 1 (Additivity of ordering). *Variable ordering is additive (in terms of set union) with respect to its first argument.*

- $\{\mathbf{ord}(vars_1 \cup vars_2) \mathbf{in} N\} \equiv \{\mathbf{ord} vars_1 \mathbf{in} N\} \cup \{\mathbf{ord} vars_2 \mathbf{in} N\}$ (as sets)
- + $\{\mathbf{ord}(vars_1 \cup vars_2) \mathbf{in} P\} \equiv \{\mathbf{ord} vars_1 \mathbf{in} P\} \cup \{\mathbf{ord} vars_2 \mathbf{in} P\}$ (as sets)

Corollary 2 (Weakening of ordering). *Extending the first argument of the ordering with unused variables does not change the result.*

- $\mathbf{ord}(vars \cap \mathbf{fv} N) \mathbf{in} N = \mathbf{ord} vars \mathbf{in} N$
- + $\mathbf{ord}(vars \cap \mathbf{fv} P) \mathbf{in} P = \mathbf{ord} vars \mathbf{in} P$

Lemma 9 (Distributivity of renaming over variable ordering). *Suppose that μ is a bijection between two sets of variables $\mu : A \leftrightarrow B$.*

- *If μ is collision free on $vars$ and $\mathbf{fv} N$ then $[\mu](\mathbf{ord} vars \mathbf{in} N) = \mathbf{ord}([\mu]vars) \mathbf{in} [\mu]N$*
- + *If μ is collision free on $vars$ and $\mathbf{fv} P$ then $[\mu](\mathbf{ord} vars \mathbf{in} P) = \mathbf{ord}([\mu]vars) \mathbf{in} [\mu]P$*

Proof. Mutual induction on N and P .

Case 1. $N = \alpha^-$

let us consider four cases:

a. $\alpha^- \in A$ and $\alpha^- \in vars$

$$\begin{aligned}
 \text{Then } [\mu](\mathbf{ord} vars \mathbf{in} N) &= [\mu](\mathbf{ord} vars \mathbf{in} \alpha^-) \\
 &= [\mu]\alpha^- && \text{by Rule (Var}_\epsilon^+) \\
 &= \beta^- && \text{for some } \beta^- \in B \text{ (notice that } \beta^- \in [\mu]vars) \\
 &= \mathbf{ord} [\mu]vars \mathbf{in} \beta^- && \text{by Rule (Var}_\epsilon^+), \text{ because } \beta^- \in [\mu]vars \\
 &= \mathbf{ord} [\mu]vars \mathbf{in} [\mu]\alpha^-
 \end{aligned}$$

b. $\alpha^- \notin A$ and $\alpha^- \notin vars$

Notice that $[\mu](\mathbf{ord} vars \mathbf{in} N) = [\mu](\mathbf{ord} vars \mathbf{in} \alpha^-) = \cdot$ by Rule (Var $_{\notin}^+$). On the other hand, $\mathbf{ord} [\mu]vars \mathbf{in} [\mu]\alpha^- = \mathbf{ord} [\mu]vars \mathbf{in} \alpha^- = \cdot$. The latter equality is from Rule (Var $_{\notin}^+$), because μ is collision free on $vars$ and $\mathbf{fv} N$, so $\mathbf{fv} N \ni \alpha^- \notin \mu(A \cap vars) \cup vars \supseteq [\mu]vars$.

c. $\alpha^- \in A$ but $\alpha^- \notin vars$

Then $[\mu](\mathbf{ord} vars \mathbf{in} N) = [\mu](\mathbf{ord} vars \mathbf{in} \alpha^-) = \cdot$ by Rule (Var $_{\notin}^+$). To prove that $\mathbf{ord} [\mu]vars \mathbf{in} [\mu]\alpha^- = \cdot$, we apply Rule (Var $_{\notin}^+$). Let us show that $[\mu]\alpha^- \notin [\mu]vars$. Since $[\mu]\alpha^- = \mu(\alpha^-)$ and $[\mu]vars \subseteq \mu(A \cap vars) \cup vars$, it suffices to prove $\mu(\alpha^-) \notin \mu(A \cap vars) \cup vars$.

- (i) If there is an element $x \in A \cap vars$ such that $\mu x = \mu\alpha^-$, then $x = \alpha^-$ by bijectivity of μ , which contradicts with $\alpha^- \notin vars$. This way, $\mu(\alpha^-) \notin \mu(A \cap vars)$.
- (ii) Since μ is collision free on $vars$ and $\mathbf{fv} N$, $\mu(A \cap \mathbf{fv} N) \ni \mu(\alpha^-) \notin vars$.

d. $\alpha^- \notin A$ but $\alpha^- \in vars$

$\mathbf{ord} [\mu]vars \mathbf{in} [\mu]\alpha^- = \mathbf{ord} [\mu]vars \mathbf{in} \alpha^- = \alpha^-$. The latter is by Rule (Var $_{\notin}^+$), because $\alpha^- = [\mu]\alpha^- \in [\mu]vars$ since $\alpha^- \in vars$. On the other hand, $[\mu](\mathbf{ord} vars \mathbf{in} N) = [\mu](\mathbf{ord} vars \mathbf{in} \alpha^-) = [\mu]\alpha^- = \alpha^-$.

Case 2. $N = \uparrow P$

$$\begin{aligned}
 [\mu](\mathbf{ord} vars \mathbf{in} N) &= [\mu](\mathbf{ord} vars \mathbf{in} \uparrow P) \\
 &= [\mu](\mathbf{ord} vars \mathbf{in} P) && \text{by Rule } (\uparrow) \\
 &= \mathbf{ord} [\mu]vars \mathbf{in} [\mu]P && \text{by the induction hypothesis} \\
 &= \mathbf{ord} [\mu]vars \mathbf{in} \uparrow[\mu]P && \text{by Rule } (\uparrow) \\
 &= \mathbf{ord} [\mu]vars \mathbf{in} [\mu]\uparrow P && \text{by the definition of substitution} \\
 &= \mathbf{ord} [\mu]vars \mathbf{in} [\mu]N
 \end{aligned}$$

Case 3. $N = P \rightarrow M$

$$\begin{aligned}
 [\mu](\mathbf{ord} vars \mathbf{in} N) &= [\mu](\mathbf{ord} vars \mathbf{in} P \rightarrow M) \\
 &= [\mu](\vec{\alpha}_1, (\vec{\alpha}_2 \setminus \{\vec{\alpha}_1\})) && \text{where } \mathbf{ord} vars \mathbf{in} P = \vec{\alpha}_1 \text{ and } \mathbf{ord} vars \mathbf{in} M = \vec{\alpha}_2 \\
 &= [\mu]\vec{\alpha}_1, [\mu](\vec{\alpha}_2 \setminus \{\vec{\alpha}_1\}) \\
 &= [\mu]\vec{\alpha}_1, ([\mu]\vec{\alpha}_2 \setminus [\mu]\{\vec{\alpha}_1\}) && \text{by induction on } \vec{\alpha}_2; \text{ the inductive step is similar to case 1. Notice that } \mu \text{ is} \\
 & && \text{collision free on } \{\vec{\alpha}_1\} \text{ and } \{\vec{\alpha}_2\} \text{ since } \{\vec{\alpha}_1\} \subseteq vars \text{ and } \{\vec{\alpha}_2\} \subseteq \mathbf{fv} N \\
 &= [\mu]\vec{\alpha}_1, ([\mu]\vec{\alpha}_2 \setminus \{[\mu]\vec{\alpha}_1\}) \\
 (\mathbf{ord} [\mu]vars \mathbf{in} [\mu]N) &= (\mathbf{ord} [\mu]vars \mathbf{in} [\mu]P \rightarrow [\mu]M) \\
 &= (\vec{\beta}_1, (\vec{\beta}_2 \setminus \{\vec{\beta}_1\})) && \text{where } \mathbf{ord} [\mu]vars \mathbf{in} [\mu]P = \vec{\beta}_1 \text{ and } \mathbf{ord} [\mu]vars \mathbf{in} [\mu]M = \vec{\beta}_2 \\
 & && \text{then by the induction hypothesis, } \vec{\beta}_1 = [\mu]\vec{\alpha}_1, \vec{\beta}_2 = [\mu]\vec{\alpha}_2, \\
 &= [\mu]\vec{\alpha}_1, ([\mu]\vec{\alpha}_2 \setminus \{[\mu]\vec{\alpha}_1\})
 \end{aligned}$$

Case 4. $N = \forall \alpha^+ . M$

$$\begin{aligned}
[\mu](\mathbf{ord\,vars\,in}\,N) &= [\mu]\mathbf{ord\,vars\,in}\,\forall \alpha^+ . M \\
&= [\mu]\mathbf{ord\,vars\,in}\,M \\
&= \mathbf{ord}\,[\mu]\mathbf{vars\,in}\,[\mu]M \quad \text{by the induction hypothesis} \\
(\mathbf{ord}\,[\mu]\mathbf{vars\,in}\,[\mu]N) &= \mathbf{ord}\,[\mu]\mathbf{vars\,in}\,[\mu]\forall \alpha^+ . M \\
&= \mathbf{ord}\,[\mu]\mathbf{vars\,in}\,\forall \alpha^+ . [\mu]M \\
&= \mathbf{ord}\,[\mu]\mathbf{vars\,in}\,[\mu]M
\end{aligned}$$

□

Lemma 10 (Ordering is not affected by independent substitutions). *Suppose that $\Gamma_2 \vdash \sigma : \Gamma_1$, i.e. σ maps variables from Γ_1 into types taking free variables from Γ_2 , and \mathbf{vars} is a set of variables disjoint with both Γ_1 and Γ_2 . Then*

$$\begin{aligned}
- \mathbf{ord\,vars\,in}\,[\sigma]N &= \mathbf{ord\,vars\,in}\,N \\
+ \mathbf{ord\,vars\,in}\,[\sigma]P &= \mathbf{ord\,vars\,in}\,P
\end{aligned}$$

Proof. **Ilya:** Should be easy

□

Lemma 11 (Completeness of variable ordering). *Variable ordering is invariant under equivalence. For arbitrary \mathbf{vars} ,*

$$\begin{aligned}
- \text{If } N \simeq_1^D M \text{ then } \mathbf{ord\,vars\,in}\,N &= \mathbf{ord\,vars\,in}\,M \text{ (as lists)} \\
+ \text{If } P \simeq_1^D Q \text{ then } \mathbf{ord\,vars\,in}\,P &= \mathbf{ord\,vars\,in}\,Q \text{ (as lists)}
\end{aligned}$$

Proof. Mutual induction on $N \simeq_1^D M$ and $P \simeq_1^D Q$.

□

4.7 Normalizaion

Lemma 12. *Set of free variables is invariant under equivalence.*

$$\begin{aligned}
- \text{If } N \simeq_1^D M \text{ then } \mathbf{fv}\,N &\equiv \mathbf{fv}\,M \text{ (as sets)} \\
+ \text{If } P \simeq_1^D Q \text{ then } \mathbf{fv}\,P &\equiv \mathbf{fv}\,Q \text{ (as sets)}
\end{aligned}$$

Proof. Straightforward mutual induction on $N \simeq_1^D M$ and $P \simeq_1^D Q$

□

Lemma 13. *Free variables are not changed by the normalization*

$$\begin{aligned}
- \mathbf{fv}\,N &\equiv \mathbf{fv}\,\mathbf{nf}\,(N) \\
+ \mathbf{fv}\,P &\equiv \mathbf{fv}\,\mathbf{nf}\,(P)
\end{aligned}$$

Proof. By straightforward induction on $\mathbf{nf}\,(N) = M$.

□

Lemma 14 (Soundness of quantifier normalization).

$$\begin{aligned}
- N &\simeq_1^D \mathbf{nf}\,(N) \\
+ P &\simeq_1^D \mathbf{nf}\,(P)
\end{aligned}$$

Proof. Mutual induction on $\mathbf{nf}\,(N) = M$ and $\mathbf{nf}\,(P) = Q$. Let us consider how this judgment is formed:

Case 1. (\mathbf{Var}^-) and (\mathbf{Var}^+)

By the corresponding equivalence rules.

Case 2. (\uparrow), (\downarrow), and (\rightarrow)

By the induction hypothesis and the corresponding congruent equivalence rules.

Case 3. (\forall), i.e. $\mathbf{nf}\,(\forall \alpha^+ . N) = \forall \alpha^+ . N'$

From the induction hypothesis, we know that $N \simeq_1^D N'$. In particular, by lemma 12, $\mathbf{fv}\,N \equiv \mathbf{fv}\,N'$. Then by lemma 8, $\{\alpha^+\} \equiv \{\alpha^+\} \cap \mathbf{fv}\,N' \equiv \{\alpha^+\} \cap \mathbf{fv}\,N$, and thus, $\{\alpha^+\}' \cap \mathbf{fv}\,N' \equiv \{\alpha^+\}' \cap \mathbf{fv}\,N$.

To prove $\forall \alpha^+ . N \simeq_1^D \forall \alpha^+ . N'$, it suffices to provide a bijection $\mu : \{\alpha^+\}' \cap \mathbf{fv}\,N' \leftrightarrow \{\alpha^+\}' \cap \mathbf{fv}\,N$ such that $N \simeq_1^D [\mu]N'$. Since these sets are equal, we take $\mu = id$.

Case 4. (\exists) Same as for case 3.

□

Corollary 3 (Normalization preserves ordering). *For any vars,*

- $\mathbf{ord\,vars\,in\,nf}\,(N) = \mathbf{ord\,vars\,in}\,M$
- + $\mathbf{ord\,vars\,in\,nf}\,(P) = \mathbf{ord\,vars\,in}\,Q$

Proof. Immediately from lemmas 11 and 14. □

Lemma 15 (Distributivity of normalization over substitution). *Normalization of a term distributes over substitution. Suppose that $\Gamma_2 \vdash \sigma : \Gamma_1$, i.e. σ maps variables from Γ_1 into types taking free variables from Γ_2 . Then*

- $\mathbf{nf}\,([\sigma]N) = [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(N)$
- + $\mathbf{nf}\,([\sigma]P) = [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(P)$

where $\mathbf{nf}\,(\sigma)$ means pointwise normalization: $[\mathbf{nf}\,(\sigma)]\alpha^- = \mathbf{nf}\,([\sigma]\alpha^-)$.

Proof. Mutual induction on N and P .

Case 1. $N = \alpha^-$

$$\begin{aligned} \mathbf{nf}\,([\sigma]N) &= \mathbf{nf}\,([\sigma]\alpha^-) = [\mathbf{nf}\,(\sigma)]\alpha^-. \\ [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(N) &= [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(\alpha^-) = [\mathbf{nf}\,(\sigma)]\alpha^-. \end{aligned}$$

Case 2. $P = \alpha^+$

Similar to case 1.

Case 3. If the type is formed by \rightarrow , \uparrow , or \downarrow , the required equality follows from the congruence of the normalization and substitution, and the induction hypothesis. For example, if $N = P \rightarrow M$ then

$$\begin{aligned} \mathbf{nf}\,([\sigma]N) &= \mathbf{nf}\,([\sigma](P \rightarrow M)) \\ &= \mathbf{nf}\,([\sigma]P \rightarrow [\sigma]M) && \text{By the congruence of substitution} \\ &= \mathbf{nf}\,([\sigma]P) \rightarrow \mathbf{nf}\,([\sigma]M) && \text{By the congruence of normalization, i.e. Rule } (\rightarrow) \\ &= [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(P) \rightarrow [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(M) && \text{By the induction hypothesis} \\ &= [\mathbf{nf}\,(\sigma)](\mathbf{nf}\,(P) \rightarrow \mathbf{nf}\,(M)) && \text{By the congruence of substitution} \\ &= [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(P \rightarrow M) && \text{By the congruence of normalization} \\ &= [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(N) \end{aligned}$$

Case 4. $N = \forall \alpha^{\vec{\alpha}}. M$

$$\begin{aligned} [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(N) &= [\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(\forall \alpha^{\vec{\alpha}}. M) \\ &= [\mathbf{nf}\,(\sigma)]\forall \alpha^{\vec{\alpha}'}.\mathbf{nf}\,(M) \quad \text{Where } \vec{\alpha}^{\vec{\alpha}'} = \mathbf{ord}\,\{\vec{\alpha}^{\vec{\alpha}}\} \mathbf{in}\, \mathbf{nf}\,(M) = \mathbf{ord}\,\{\vec{\alpha}^{\vec{\alpha}}\} \mathbf{in}\, M \text{ (the latter is by corollary 3)} \end{aligned}$$

$$\begin{aligned} \mathbf{nf}\,([\sigma]N) &= \mathbf{nf}\,([\sigma]\forall \alpha^{\vec{\alpha}}. M) \\ &= \mathbf{nf}\,(\forall \alpha^{\vec{\alpha}}. [\sigma]M) && \text{Assuming } \{\vec{\alpha}^{\vec{\alpha}}\} \cap \{\Gamma_1\} = \emptyset \text{ and } \{\vec{\alpha}^{\vec{\alpha}}\} \cap \{\Gamma_2\} = \emptyset \\ &= \forall \beta^{\vec{\beta}}.\mathbf{nf}\,([\sigma]M) && \text{Where } \vec{\beta}^{\vec{\beta}} = \mathbf{ord}\,\{\vec{\alpha}^{\vec{\alpha}}\} \mathbf{in}\, \mathbf{nf}\,([\sigma]M) = \mathbf{ord}\,\{\vec{\alpha}^{\vec{\alpha}}\} \mathbf{in}\, [\sigma]M \text{ (the latter is by corollary 3)} \\ &= \forall \alpha^{\vec{\alpha}'}.\mathbf{nf}\,([\sigma]M) && \text{By lemma 10, } \vec{\beta}^{\vec{\beta}} = \vec{\alpha}^{\vec{\alpha}'} \text{ since } \{\vec{\alpha}^{\vec{\alpha}}\} \text{ is disjoint with } \Gamma_1 \text{ and } \Gamma_2 \\ &= \forall \alpha^{\vec{\alpha}'}.[\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(M) && \text{By the induction hypothesis} \end{aligned}$$

To show alpha-equivalence of $[\mathbf{nf}\,(\sigma)]\forall \alpha^{\vec{\alpha}'}.\mathbf{nf}\,(M)$ and $\forall \alpha^{\vec{\alpha}'}.[\mathbf{nf}\,(\sigma)]\mathbf{nf}\,(M)$, we can assume that $\{\vec{\alpha}^{\vec{\alpha}'}\} \cap \{\Gamma_1\} = \emptyset$, and $\{\vec{\alpha}^{\vec{\alpha}'}\} \cap \{\Gamma_2\} = \emptyset$.

Case 5. $P = \exists \alpha^{\vec{\alpha}}. Q$

Same as for case 4.

□

Corollary 4 (Commutativity of normalization and renaming). *Normalization of a term commutes with renaming. Suppose that μ is a bijection between two sets of variables $\mu : A \leftrightarrow B$. Then*

- $\mathbf{nf}\,([\mu]N) = [\mu]\mathbf{nf}\,(N)$
- + $\mathbf{nf}\,([\mu]P) = [\mu]\mathbf{nf}\,(P)$

Proof. Immediately from lemma 15, after noticing that $\mathbf{nf}(\mu) = \mu$. □

Lemma 16 (Completeness of quantified normalization). *Normalization returns the same representative for equivalent types.*

- If $N \simeq_1^D M$ then $\mathbf{nf}(N) = \mathbf{nf}(M)$
- + If $P \simeq_1^D Q$ then $\mathbf{nf}(P) = \mathbf{nf}(Q)$

(Here equality means alpha-equivalence)

Proof. Mutual induction on $N \simeq_1^D M$ and $P \simeq_1^D Q$.

Case 1. ($\forall \alpha^{\vec{+}}_1$)

From the definition of the normalization,

- $\mathbf{nf}(\forall \alpha^{\vec{+}}.N) = \forall \alpha^{\vec{+}'}. \mathbf{nf}(N)$ where $\alpha^{\vec{+}'}$ is **ord** $\{\alpha^{\vec{+}}\}$ in $\mathbf{nf}(N)$
- $\mathbf{nf}(\forall \beta^{\vec{+}}.M) = \forall \beta^{\vec{+}'}. \mathbf{nf}(M)$ where $\beta^{\vec{+}'}$ is **ord** $\{\beta^{\vec{+}}\}$ in $\mathbf{nf}(M)$

Let us take $\mu : (\{\beta^{\vec{+}}\} \cap \mathbf{fv} M) \leftrightarrow (\{\alpha^{\vec{+}}\} \cap \mathbf{fv} N)$ from the inversion of the equivalence judgment. Notice that from lemmas 8 and 13, the domain and the codomain of μ can be written as $\mu : \{\beta^{\vec{+}'}\} \leftrightarrow \{\alpha^{\vec{+}'}\}$.

To show the alpha-equivalence of $\forall \alpha^{\vec{+}'}. \mathbf{nf}(N)$ and $\forall \beta^{\vec{+}'}. \mathbf{nf}(M)$, it suffices to prove that (i) $[\mu] \mathbf{nf}(M) = \mathbf{nf}(N)$ and (ii) $[\mu] \beta^{\vec{+}'} = \alpha^{\vec{+}'}$.

(i) $[\mu] \mathbf{nf}(M) = \mathbf{nf}([\mu]M) = \mathbf{nf}(N)$. The first equality holds by corollary 4, the second—by the induction hypothesis.

$$\begin{aligned}
 \text{(ii) } [\mu] \beta^{\vec{+}'} &= [\mu] \mathbf{ord} \{\beta^{\vec{+}}\} \mathbf{in} \mathbf{nf}(M) && \text{by the definition of } \beta^{\vec{+}'} \\
 &= [\mu] \mathbf{ord} (\{\beta^{\vec{+}}\} \cap \mathbf{fv} M) \mathbf{in} \mathbf{nf}(M) && \text{from lemma 13 and corollary 2} \\
 &= \mathbf{ord} [\mu](\{\beta^{\vec{+}}\} \cap \mathbf{fv} M) \mathbf{in} [\mu] \mathbf{nf}(M) && \text{by lemma 9, because } \{\alpha^{\vec{+}}\} \cap \mathbf{fv} N \cap \mathbf{fv} \mathbf{nf}(M) \subseteq \{\alpha^{\vec{+}}\} \cap \mathbf{fv} M = \emptyset \\
 &&& \text{and } \{\alpha^{\vec{+}}\} \cap \mathbf{fv} N \cap (\{\beta^{\vec{+}}\} \cap \mathbf{fv} M) \subseteq \{\alpha^{\vec{+}}\} \cap \mathbf{fv} M = \emptyset \\
 &= \mathbf{ord} [\mu](\{\beta^{\vec{+}}\} \cap \mathbf{fv} M) \mathbf{in} \mathbf{nf}(N) && \text{since } [\mu] \mathbf{nf}(M) = \mathbf{nf}(N) \text{ is proved} \\
 &= \mathbf{ord} (\{\alpha^{\vec{+}}\} \cap \mathbf{fv} N) \mathbf{in} \mathbf{nf}(N) && \text{because } \mu \text{ is a bijection between } \{\alpha^{\vec{+}}\} \cap \mathbf{fv} N \text{ and } \{\beta^{\vec{+}}\} \cap \mathbf{fv} M \\
 &= \mathbf{ord} \{\alpha^{\vec{+}}\} \mathbf{in} \mathbf{nf}(N) && \text{from lemma 13 and corollary 2} \\
 &= \alpha^{\vec{+}'} && \text{by the definition of } \alpha^{\vec{+}'}
 \end{aligned}$$

Case 2. ($\exists \alpha^{\vec{+}}_1$) Same as for case 1.

Case 3. Other rules are congruent, and thus, proved by the corresponding congruent alpha-equivalence rule, which is applicable by the induction hypothesis. □

Lemma 17 (Idempotence of normalization). *Normalization is idempotent*

- $\mathbf{nf}(\mathbf{nf}(N)) = \mathbf{nf}(N)$
- + $\mathbf{nf}(\mathbf{nf}(P)) = \mathbf{nf}(P)$

Proof. By applying lemma 16 to lemma 14. □

Lemma 18. *The result of a substitution is normalized if and only if the initial type and the substitution are normalized.*

Suppose that σ is a substitution $\Gamma_2 \vdash \sigma : \Gamma_1$, P is a positive type ($\Gamma_1 \vdash P$), N is a negative type ($\Gamma_1 \vdash N$). Then

$$\begin{aligned}
 + [\sigma]P \text{ is normal} &\iff \begin{cases} \sigma|_{\mathbf{fv}(P)} & \text{is normal} \\ P & \text{is normal} \end{cases} \\
 - [\sigma]N \text{ is normal} &\iff \begin{cases} \sigma|_{\mathbf{fv}(N)} & \text{is normal} \\ N & \text{is normal} \end{cases}
 \end{aligned}$$

Proof. Mutual induction on $\Gamma_1 \vdash P$ and $\Gamma_1 \vdash N$.

Case 1. $N = \alpha^-$

Then N is always normal, and the normality of $\sigma|_{\{\alpha^-\}}$ by the definition means $[\sigma]\alpha^-$ is normal.

Case 2. $N = P \rightarrow M$

$$\begin{aligned}
[\sigma](P \rightarrow M) \text{ is normal} &\iff [\sigma]P \rightarrow [\sigma]M \text{ is normal} && \text{by the substitution congruence} \\
&\iff \begin{cases} [\sigma]P & \text{is normal} \\ [\sigma]M & \text{is normal} \end{cases} && \text{by congruence of normality Ilya: lemma?} \\
&\iff \begin{cases} P & \text{is normal} \\ \sigma|_{\mathbf{fv}(P)} & \text{is normal} \\ M & \text{is normal} \\ \sigma|_{\mathbf{fv}(M)} & \text{is normal} \end{cases} && \text{by the induction hypothesis} \\
&\iff \begin{cases} P \rightarrow M & \text{is normal} \\ \sigma|_{\mathbf{fv}(P) \cup \mathbf{fv}(M)} & \text{is normal} \end{cases} \iff \begin{cases} P \rightarrow M & \text{is normal} \\ \sigma|_{\mathbf{fv}(P \rightarrow M)} & \text{is normal} \end{cases}
\end{aligned}$$

Case 3. $N = \uparrow P$

By congruence and the inductive hypothesis, similar to case 2

Case 4. $N = \forall \alpha^+ . M$

$$\begin{aligned}
[\sigma](\forall \alpha^+ . M) \text{ is normal} &\iff (\forall \alpha^+ . [\sigma]M) \text{ is normal} && \text{assuming } \vec{\alpha}^+ \cap \Gamma_1 = \emptyset \text{ and } \vec{\alpha}^+ \cap \Gamma_2 = \emptyset \\
&\iff \begin{cases} [\sigma]M \text{ is normal} \\ \mathbf{ord} \{ \vec{\alpha}^+ \} \text{ in } [\sigma]M = \vec{\alpha}^+ \end{cases} && \text{by the definition of normalization} \\
&\iff \begin{cases} [\sigma]M \text{ is normal} \\ \mathbf{ord} \{ \vec{\alpha}^+ \} \text{ in } M = \vec{\alpha}^+ \end{cases} && \text{by lemma 10} \\
&\iff \begin{cases} \sigma|_{\mathbf{fv}(M)} \text{ is normal} \\ M \text{ is normal} \\ \mathbf{ord} \{ \vec{\alpha}^+ \} \text{ in } M = \vec{\alpha}^+ \end{cases} && \text{by the induction hypothesis} \\
&\iff \begin{cases} \sigma|_{\mathbf{fv}(\forall \alpha^+ . M)} \text{ is normal} \\ \forall \alpha^+ . M \text{ is normal} \end{cases} && \begin{array}{l} \text{since } \mathbf{fv}(\forall \alpha^+ . M) = \mathbf{fv}(M); \\ \text{by the definition of normalization} \end{array}
\end{aligned}$$

Case 5. $P = \dots$

The positive cases are done in the same way as the negative ones.

□

4.8 Equivalence

Lemma 19 (Type well-formedness is invariant under equivalence). *Mutual subtyping implies declarative equivalence.*

- + if $P \simeq_1^D Q$ then $\Gamma \vdash P \iff \Gamma \vdash Q$,
- if $N \simeq_1^D M$ then $\Gamma \vdash N \iff \Gamma \vdash M$

Proof. Ilya: todo

□

Corollary 5 (Normalization preserves well-formedness).

- + $\Gamma \vdash P \iff \Gamma \vdash \mathbf{nf}(P)$,
- $\Gamma \vdash N \iff \Gamma \vdash \mathbf{nf}(N)$

Proof. Immediately from lemmas 14 and 19.

□

Corollary 6 (Normalization preserves well-formedness of substitution).

$$\Gamma_2 \vdash \sigma : \Gamma_1 \iff \Gamma_2 \vdash \mathbf{nf}(\sigma) : \Gamma_1$$

Lemma 20 (Soundness of equivalence). *Declarative equivalence implies mutual subtyping.*

- + if $\Gamma \vdash P$, $\Gamma \vdash Q$, and $P \simeq_1^D Q$ then $\Gamma \vdash P \simeq_1^{\leq} Q$,

- if $\Gamma \vdash N$, $\Gamma \vdash M$, and $N \simeq_1^D M$ then $\Gamma \vdash N \simeq_1^\leq M$.

Proof. Ilya: todo □

Lemma 21 (Completeness of equivalence). *Mutual subtyping implies declarative equivalence.*

- + if $\Gamma \vdash P \simeq_1^\leq Q$ then $P \simeq_1^D Q$,
- if $\Gamma \vdash N \simeq_1^\leq M$ then $N \simeq_1^D M$.

Proof. Ilya: todo □

4.9 Upper Bounds

Lemma 22 (Decomposition of the quantifier rule). *Ilya: move somewhere* Whenever the quantifier rule (Rule $(\exists^{\geq 1})$ or Rule $(\forall^{\leq 1})$) is applied, one can assume that the rule adding quantifiers on the right-hand side was applied the last.

- If $\Gamma \vdash N \leq_1 \forall \vec{\beta}^+. M$ then $\Gamma, \vec{\beta}^+ \vdash N \leq_1 M$.
- + If $\Gamma \vdash P \geq_1 \exists \vec{\beta}^+. Q$ then $\Gamma, \vec{\beta}^+ \vdash P \geq_1 Q$.

Lemma 23 (Characterization of the Supertypes). *Let us define the set of upper bounds of a positive type $\text{UB}(P)$ in the following way:*

$\Gamma \vdash P$	$\text{UB}(\Gamma \vdash P)$
$\Gamma \vdash \beta^+$	$\{\exists \vec{\alpha}^+. \beta^+ \mid \text{for } \vec{\alpha}^+\}$
$\Gamma \vdash \exists \vec{\beta}^+. Q$	$\text{UB}(\Gamma, \vec{\beta}^+ \vdash Q) \text{ not using } \vec{\beta}^+$
$\Gamma \vdash \downarrow M$	$\left\{ \exists \vec{\alpha}^+. \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^+, M', \text{ and } \vec{N} \text{ s.t.} \\ \Gamma \vdash N_i, \Gamma, \vec{\alpha}^+ \vdash M', \text{ and } [\vec{N}/\vec{\alpha}^+] \downarrow M' \simeq_1^D \downarrow M \end{array} \right\}$
Then $\text{UB}(\Gamma \vdash P) \equiv \{Q \mid \Gamma \vdash Q \geq_1 P\}$.	

Proof. By induction on $\Gamma \vdash P$.

Case 1. $P = \beta^+$

Then the last rule that is applied to infer $\Gamma \vdash Q \geq_1 \beta^+$ must be either Rule $(\text{Var}^{+ \geq 1})$ or Rule $(\exists^{\geq 1})$. The former case means that $Q = \beta^+$. In the latter case, $Q = \exists \vec{\alpha}^+. Q'$, where Q' has no outer existential quantifiers. Then by inversion of Rule $(\exists^{\geq 1})$, $\Gamma \vdash [\vec{N}/\vec{\alpha}^+] Q' \geq_1 \beta^+$ for some \vec{N} . This time, to infer this judgment, only Rule $(\text{Var}^{+ \geq 1})$ is applicable, which means that $Q' = \beta^+$, and then $Q = \exists \vec{\alpha}^+. \beta^+$.

Case 2. $P = \exists \vec{\beta}^+. P'$

Then if $\Gamma \vdash Q \geq_1 \exists \vec{\beta}^+. P'$, then by lemma 22, $\Gamma, \vec{\beta}^+ \vdash Q \geq_1 P'$, and $\text{fv } Q \cap \{\vec{\beta}^+\} = \emptyset$ by the the Barendregt's convention. The other direction holds by Rule $(\exists^{\geq 1})$. This way, $\{Q \mid \Gamma \vdash Q \geq_1 \exists \vec{\beta}^+. P'\} = \{Q \mid \Gamma, \vec{\beta}^+ \vdash Q \geq_1 P' \text{ s.t. } \text{fv}(Q) \cap \{\vec{\beta}^+\} = \emptyset\}$. From the induction hypothesis, the latter is equal to $\text{UB}(\Gamma, \vec{\beta}^+ \vdash P')$ not using $\vec{\beta}^+$, i.e. $\text{UB}(\Gamma \vdash \exists \vec{\beta}^+. P')$.

Case 3. $P = \downarrow M$

Then let us consider two subcases upper bounds without outer quantifiers (we denote the corresponding set restriction as $|\#$) and upper bounds with outer quantifiers ($|\exists$). We prove that for both of these groups, the restricted sets are equal.

a. $Q \neq \exists \vec{\beta}^+. Q'$

Then the last applied rule to infer $\Gamma \vdash Q \geq_1 \downarrow M$ must be Rule $(\downarrow^{\geq 1})$, which means $Q = \downarrow M'$, and by inversion, $\Gamma \vdash M' \simeq_1^\leq M$, then by lemma 21 and Rule $(\downarrow^{\leq 1})$, $\downarrow M' \simeq_1^D \downarrow M$. This way, $Q = \downarrow M' \in \{\downarrow M' \mid \downarrow M' \simeq_1^D \downarrow M\} = \text{UB}(\Gamma \vdash \downarrow M)|\#$. In the other direction, $\downarrow M' \simeq_1^D \downarrow M \Rightarrow \Gamma \vdash \downarrow M' \simeq_1^\leq \downarrow M$ by lemma 20, since $\Gamma \vdash \downarrow M'$ by lemma 19 $\Rightarrow \Gamma \vdash \downarrow M' \geq_1 \downarrow M$ by inversion

b. $Q = \exists \vec{\beta}^+. Q'$ (for non-empty $\vec{\beta}^+$)

Then the last rule applied to infer $\Gamma \vdash \exists \vec{\beta}^+. Q' \geq_1 \downarrow M$ must be Rule $(\exists^{\geq 1})$. Inversion of this rule gives us $\Gamma \vdash [\vec{N}/\vec{\beta}^+] Q' \geq_1 \downarrow M$ for some $\Gamma \vdash N_i$. Notice that $[\vec{N}/\vec{\beta}^+] Q'$ has no outer quantifiers. Thus from case 3.a, $[\vec{N}/\vec{\beta}^+] Q' \simeq_1^D \downarrow M$, which is only possible if $Q' = \downarrow M'$. This way, $Q = \exists \vec{\beta}^+. \downarrow M' \in \text{UB}(\Gamma \vdash \downarrow M)|\exists$ (notice that $\vec{\beta}^+$ is not empty).

In the other direction, $[\vec{N}/\vec{\beta}^-]\downarrow M' \simeq_1^D \downarrow M \Rightarrow \Gamma \vdash [\vec{N}/\vec{\beta}^-]\downarrow M' \simeq_1^{\leq} \downarrow M$ by lemma 20, since $\Gamma \vdash [\vec{N}/\vec{\beta}^-]\downarrow M'$ by lemma 19
 $\Rightarrow \Gamma \vdash [\vec{N}/\vec{\beta}^-]\downarrow M' \geq_1 \downarrow M$ by inversion
 $\Rightarrow \Gamma \vdash \exists \vec{\beta}^-. \downarrow M' \geq_1 \downarrow M$ by Rule (\exists^{\geq_1})

□

Lemma 24 (Characterization of the Normalized Supertypes). *For a normalized positive type $P = \mathbf{nf}(P)$, let us define the set of normalized upper bounds in the following way:*

$$\frac{\Gamma \vdash P}{\text{NFUB}(\Gamma \vdash P)}$$

$$\Gamma \vdash \beta^+ \quad \{\beta^+\}$$

$$\Gamma \vdash \exists \vec{\beta}^-. P \quad \text{NFUB}(\Gamma, \vec{\beta}^- \vdash P) \text{ not using } \vec{\beta}^-$$

$$\Gamma \vdash \downarrow M \quad \left\{ \exists \vec{\alpha}^-. \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \text{ and } \vec{N} \text{ s.t. } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^-, \\ \Gamma \vdash N_i, \Gamma, \vec{\alpha}^- \vdash M', \text{ and } [\vec{N}/\vec{\alpha}^-]\downarrow M' = \downarrow M \end{array} \right\}$$

Then $\text{NFUB}(\Gamma \vdash P) \equiv \{\mathbf{nf}(Q) \mid \Gamma \vdash Q \geq_1 P\}$.

Proof. By induction on $\Gamma \vdash P$.

Case 1. $P = \beta^+$

Then from lemma 23, $\{\mathbf{nf}(Q) \mid \Gamma \vdash Q \geq_1 \beta^+\} = \{\mathbf{nf}(\exists \vec{\alpha}^-. \beta^+) \mid \text{for some } \vec{\alpha}^- = \{\beta^+\}\}$

Case 2. $P = \exists \vec{\beta}^-. P'$

$$\begin{aligned} \text{NFUB}(\Gamma \vdash \exists \vec{\beta}^-. P') &= \text{NFUB}(\Gamma, \vec{\beta}^- \vdash P') \text{ not using } \vec{\beta}^- \\ &= \{\mathbf{nf}(Q) \mid \Gamma, \vec{\beta}^- \vdash Q \geq_1 P'\} \text{ not using } \vec{\beta}^- && \text{by the induction hypothesis} \\ &= \{\mathbf{nf}(Q) \mid \Gamma, \vec{\beta}^- \vdash Q \geq_1 P' \text{ s.t. } \mathbf{fv} Q \cap \vec{\beta}^- = \emptyset\} && \text{because } \mathbf{fv} \mathbf{nf}(Q) = \mathbf{fv} Q \text{ by lemma 13} \\ &= \{\mathbf{nf}(Q) \mid Q \in \text{UB}(\Gamma, \vec{\beta}^- \vdash P') \text{ s.t. } \mathbf{fv} Q \cap \vec{\beta}^- = \emptyset\} && \text{by lemma 23} \\ &= \{\mathbf{nf}(Q) \mid Q \in \text{UB}(\Gamma \vdash \exists \vec{\beta}^-. P')\} && \text{by the definition of UB} \\ &= \{\mathbf{nf}(Q) \mid \Gamma \vdash Q \geq_1 \exists \vec{\beta}^-. P'\} && \text{by lemma 23} \end{aligned}$$

Case 3. $P = \downarrow M$

In the following reasoning, we will use the following principle of variable replacement.

Observation 1. *Suppose that $\nu : A \rightarrow A$ is an idempotent function, P is a predicate on A , $F : A \rightarrow B$ is a function. Then*

$$\begin{aligned} &\{F(\nu x) \mid x \in A \text{ s.t. } P(\nu x)\} = \\ &= \{F(x) \mid x \in A \text{ s.t. } \nu x = x \text{ and } P(x)\}. \end{aligned}$$

In our case, the idempotent ν will be normalization, variable ordering, or domain restriction.

Another observation we will use is the following.

Observation 2. *For functions F and ν , and predicates P and Q ,*

$$\begin{aligned} &\{F(\nu x) \mid x \in A \text{ s.t. } Q(\nu x) \text{ and } P(x)\} = \\ &= \{F(\nu x) \mid x \in A \text{ s.t. } Q(\nu x) \text{ and } (\exists x' \in A \text{ s.t. } P(x') \text{ and } \nu x' = \nu x)\}. \end{aligned}$$

$$\begin{aligned}
& \{\mathbf{nf}(Q) \mid \Gamma \vdash Q \succcurlyeq_1 \downarrow M\} = \\
& = \{\mathbf{nf}(Q) \mid Q \in \mathbf{UB}(\Gamma \vdash \downarrow M)\} \\
& = \left\{ \mathbf{nf}(\exists \vec{\alpha}^-. \downarrow M') \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \text{ and } \vec{N} \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash N_i, \text{ and } [\vec{N}/\vec{\alpha}^-] \downarrow M' \simeq_1^D \downarrow M \end{array} \right\} \\
& = \left\{ \mathbf{nf}(\exists \vec{\alpha}^-. \downarrow M') \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \text{ and } \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^-, \text{ and } [\sigma] \downarrow M' \simeq_1^D \downarrow M \end{array} \right\} \\
& = \left\{ \mathbf{nf}(\exists \vec{\alpha}^-. \downarrow M') \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \text{ and } \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^-, \text{ and } [\sigma|_{\mathbf{fv} M'}] \downarrow M' \simeq_1^D \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \mathbf{nf}(\downarrow M') \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^-, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\sigma|_{\mathbf{fv} M'}] \downarrow M' \simeq_1^D \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \mathbf{nf}(\downarrow M') \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^-, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } \mathbf{nf}([\sigma|_{\mathbf{fv} M'}] \downarrow M') = \mathbf{nf}(\downarrow M) \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \mathbf{nf}(\downarrow M') \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^-, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\mathbf{nf}(\sigma|_{\mathbf{fv} M'})] \downarrow \mathbf{nf}(M') = \downarrow \mathbf{nf}(M) \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^-, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\sigma|_{\mathbf{fv} M'}] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ (\exists \sigma' \text{ s.t. } \Gamma \vdash \sigma' : \vec{\alpha}^- \text{ and } \sigma|_{\mathbf{fv}(\downarrow M')} = \sigma'|_{\mathbf{fv}(\downarrow M')}) \\ \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \text{ and } [\sigma|_{\mathbf{fv} M'}] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma|_{\mathbf{fv} M'} : \vec{\alpha}^{-'}, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\sigma|_{\mathbf{fv} M'}] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^{-'}, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\sigma] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, \vec{\alpha}^-, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^- \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^{-'}, \text{ and } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\sigma] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, M', \sigma \text{ s.t. } \Gamma, \vec{\alpha}^{-'} \vdash M', \\ \Gamma \vdash \sigma : \vec{\alpha}^{-'}, \text{ and } \mathbf{ord}\{\vec{\alpha}^{-'}\} \text{ in } M' = \vec{\alpha}^{-'} \\ \text{and } [\sigma] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \left\{ \exists \vec{\alpha}^{-'} . \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^{-'}, M', \text{ and } \vec{N} \text{ s.t. } \mathbf{ord}\{\vec{\alpha}^{-'}\} \text{ in } M' = \vec{\alpha}^{-'}, \\ \Gamma \vdash N_i, \Gamma, \vec{\alpha}^{-'} \vdash M', \text{ and } [\vec{N}/\vec{\alpha}^{-'}] \downarrow M' = \downarrow M \end{array} \right\} \\
& = \mathbf{NFUB}(\downarrow M)
\end{aligned}$$

by lemma 23

by the definition of UB

we reassigned the substitution $\vec{N}/\vec{\alpha}^-$ as σ

by lemma 2

by the definition of normalization

from lemmas 14 and 16, equivalence of types can be replaced with the equality of their normal forms

by congruence of normalization and lemma 15

by lemma 18, $\downarrow M'$ and $\sigma|_{\mathbf{fv} M'}$ are already normal, since the result of the substitution is normal; M is normal by assumption

We apply observation 2 (with $\nu\sigma = \sigma|_{\mathbf{fv} M'}$, and $P(\sigma) = \Gamma \vdash \sigma : \vec{\alpha}^-$)

Notice that
“ $\exists \sigma' \text{ s.t. } (\Gamma \vdash \sigma' : \vec{\alpha}^- \text{ and } \sigma|_{\mathbf{fv}(\downarrow M')} = \sigma'|_{\mathbf{fv}(\downarrow M')})$ ”
is equivalent to $\Gamma \vdash \sigma|_{\mathbf{fv}(\downarrow M')} : \vec{\alpha}^{-'}$

We apply observation 1 to the restriction of σ , and remove $\sigma|_{\mathbf{fv} M'} = \sigma$ as it follows from $\Gamma \vdash \sigma : \vec{\alpha}^{-'}$

by lemma 4, since $\{\Gamma, \vec{\alpha}^-\} \cap \mathbf{fv} M' = \{\Gamma, \vec{\alpha}^{-'}\} \cap \mathbf{fv} M'$

We apply observation 1 to the ordering of $\vec{\alpha}^-$

By reassigning σ explicitly as $\vec{N}/\vec{\alpha}^{-'}$

by definition

□

Observation 3. Upper bounds of a type do not depend on the context as soon as the type are well-formed in it.

If $\Gamma_1 \vdash M$ and $\Gamma_2 \vdash M$ then $\mathbf{UB}(\Gamma_1 \vdash M) = \mathbf{UB}(\Gamma_2 \vdash M)$ and $\mathbf{NFUB}(\Gamma_1 \vdash M) = \mathbf{NFUB}(\Gamma_2 \vdash M)$

Proof. We prove both inclusions by induction on $\Gamma_1 \vdash M$. Notice that if $[\sigma]M' \simeq_1^D M$ and $\Gamma_2 \vdash M$ then the types from the range of $\sigma|_{\mathbf{fv} M'}$ are well-formed in 2 **Ilya: lemma**. □

Lemma 25 (Soundness of the Least Upper Bound). For types $\Gamma \vdash P_1$, and $\Gamma \vdash P_2$, if $\Gamma \models P_1 \vee P_2 = Q$ then

(i) $\Gamma \vdash Q$

(ii) $\Gamma \vdash Q \geq_1 P_1$ and $\Gamma \vdash Q \geq_1 P_2$

Proof. Induction on $\Gamma \models P_1 \vee P_2 = Q$.

Case 1. $\Gamma \models \alpha^+ \vee \alpha^+ = \alpha^+$

Then $\Gamma \vdash \alpha^+$ by assumption, and $\Gamma \vdash \alpha^+ \geq_1 \alpha^+$ by Rule (Var⁺₁).

Case 2. $\Gamma \models \exists \vec{\alpha}^-. P_1 \vee \exists \vec{\beta}^-. P_2 = Q$

Then by inversion of $\Gamma \vdash \exists \vec{\alpha}^-. P_i$ and weakening, $\Gamma, \vec{\alpha}^-, \vec{\beta}^- \vdash P_i$, hence, the induction hypothesis applied to $\Gamma, \vec{\alpha}^-, \vec{\beta}^- \models P_1 \vee P_2 = Q$. Then

- (i) $\Gamma, \vec{\alpha}^-, \vec{\beta}^- \vdash Q$,
- (ii) $\Gamma, \vec{\alpha}^-, \vec{\beta}^- \vdash Q \geq_1 P_1$,
- (iii) $\Gamma, \vec{\alpha}^-, \vec{\beta}^- \vdash Q \geq_1 P_2$.

To prove $\Gamma \vdash Q$, it suffices to show that $\mathbf{fv}(Q) \cap \{\Gamma, \vec{\alpha}^-, \vec{\beta}^-\} = \mathbf{fv}(Q) \cap \{\Gamma\}$ (and then apply lemma 4). The inclusion right-to-left is self-evident. To show $\mathbf{fv}(Q) \cap \{\Gamma, \vec{\alpha}^-, \vec{\beta}^-\} \subseteq \mathbf{fv}(Q) \cap \{\Gamma\}$, we prove that $\mathbf{fv}(Q) \subseteq \{\Gamma\}$

$$\mathbf{fv}(Q) \subseteq \mathbf{fv} P_1 \cap \mathbf{fv} P_2$$

by lemma 1

$$\begin{aligned} &\subseteq (\{\Gamma, \vec{\alpha}^-\} \setminus \{\vec{\beta}^-\}) \cap (\{\Gamma, \vec{\beta}^-\} \setminus \{\vec{\alpha}^-\}) && \text{since } \Gamma \vdash \exists \vec{\alpha}^-. P_1, \mathbf{fv}(P_1) \subseteq \{\Gamma, \vec{\alpha}^-\} = \{\Gamma, \vec{\alpha}^-\} \setminus \{\vec{\beta}^-\} \\ & && \text{(the latter is because by the Barendregt's convention, } \{\Gamma, \vec{\alpha}^-\} \cap \{\vec{\beta}^-\} = \emptyset\text{); similarly, } \mathbf{fv}(P_2) \subseteq \{\Gamma, \vec{\beta}^-\} \setminus \{\vec{\alpha}^-\} \\ &\subseteq \{\Gamma\} \end{aligned}$$

To show $\Gamma \vdash Q \geq_1 \exists \vec{\alpha}^-. P_1$, we apply Rule ($\exists \geq_1$). Then $\Gamma, \vec{\alpha}^- \vdash Q \geq_1 P_1$ holds since $\Gamma, \vec{\alpha}^-, \vec{\beta}^- \vdash Q \geq_1 P_1$ (by the induction hypothesis), $\Gamma, \vec{\alpha}^- \vdash Q$ (by weakening), and $\Gamma, \vec{\alpha}^- \vdash P_1$.

Judgment $\Gamma \vdash Q \geq_1 \exists \vec{\beta}^-. P_2$ is proved symmetrically.

Case 3. $\Gamma \models \downarrow N \vee \downarrow M = \exists \vec{\alpha}^-. [\vec{\alpha}^- / \Xi] P$ By the inversion, $\Gamma, \cdot \models \downarrow N \stackrel{a}{\simeq} \downarrow M \Rightarrow (\Xi, P, \hat{\tau}_1, \hat{\tau}_2)$. Then by lemma 5,

(i) $\Gamma; \Xi \vdash P$, then by ??,

$$\Gamma, \vec{\alpha}^- \vdash [\vec{\alpha}^- / \Xi] P \quad (1)$$

(ii) $\Gamma; \cdot \vdash \hat{\tau}_1 : \Xi$ and $\Gamma; \cdot \vdash \hat{\tau}_2 : \Xi$. Assuming that $\Xi = \hat{\beta}_1^-, \dots, \hat{\beta}_n^-$, the antiunification solutions $\hat{\tau}_1$ and $\hat{\tau}_2$ can be put explicitly as $\hat{\tau}_1 = (\hat{\beta}_1^- : \approx N_1, \dots, \hat{\beta}_n^- : \approx N_n)$, and $\hat{\tau}_2 = (\hat{\beta}_1^- : \approx M_1, \dots, \hat{\beta}_n^- : \approx M_n)$. Then

$$\hat{\tau}_1 = (\vec{N} / \vec{\alpha}^-) \circ (\vec{\alpha}^- / \Xi) \text{ (as substitutions)} \quad (2)$$

$$\hat{\tau}_2 = (\vec{M} / \vec{\alpha}^-) \circ (\vec{\alpha}^- / \Xi) \text{ (as substitutions)} \quad (3)$$

(iii) $[\hat{\tau}_1]Q = P_1$ and $[\hat{\tau}_2]Q = P_1$, which, by 2 and 3, means

$$[\vec{N} / \vec{\alpha}^-][\vec{\alpha}^- / \Xi] P = \downarrow N \quad (4)$$

$$[\vec{M} / \vec{\alpha}^-][\vec{\alpha}^- / \Xi] P = \downarrow M \quad (5)$$

Then $\Gamma \vdash \exists \vec{\alpha}^-. [\vec{\alpha}^- / \Xi] P$ follows directly from 1.

To show $\Gamma \vdash \exists \vec{\alpha}^-. [\vec{\alpha}^- / \Xi] P \geq_1 \downarrow N$, we apply Rule ($\exists \geq_1$), instantiating $\vec{\alpha}^-$ with \vec{N} . Then $\Gamma \vdash [\vec{N} / \vec{\alpha}^-][\vec{\alpha}^- / \Xi] P \geq_1 \downarrow N$ follows from 4 and reflexivity of subtyping (??).

Analogously, instantiating $\vec{\alpha}^-$ with \vec{M} , gives us $\Gamma \vdash [\vec{M} / \vec{\alpha}^-][\vec{\alpha}^- / \Xi] P \geq_1 \downarrow M$ (from 5), and hence, $\Gamma \vdash \exists \vec{\alpha}^-. [\vec{\alpha}^- / \Xi] P \geq_1 \downarrow M$. \square

Lemma 26 (Completeness of the Least Upper Bound). *For normalized types $\Gamma \vdash P_1$, $\Gamma \vdash P_2$, and $\Gamma \vdash Q$ such that $\Gamma \vdash Q \geq_1 P_1$ and $\Gamma \vdash Q \geq_1 P_2$, there exists Q' s.t. $\Gamma \models P_1 \vee P_2 = Q'$.*

Proof. Induction on the pair (P_1, P_2) . From lemma 24, $Q \in \text{UB}(\Gamma \vdash P_1) \cap \text{UB}(\Gamma \vdash P_2)$. Let us consider the cases what P_1 and P_2 are (i.e. the last rules to infer $\Gamma \vdash P_1$).

Case 1. $P_1 = \exists \vec{\beta}^-_1. Q_1$, $P_2 = \exists \vec{\beta}^-_2. Q_2$ where $\vec{\beta}^-_1$ or $\vec{\beta}^-_2$ is not empty

Then $Q \in \text{UB}(\Gamma \vdash \exists \vec{\beta}^-_1. Q_1) \cap \text{UB}(\Gamma \vdash \exists \vec{\beta}^-_2. Q_2)$

$$\subseteq \text{UB}(\Gamma, \vec{\beta}^-_1 \vdash Q_1) \cap \text{UB}(\Gamma, \vec{\beta}^-_2 \vdash Q_2)$$

from the definition of UB

$$= \text{UB}(\Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \vdash Q_1) \cap \text{UB}(\Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \vdash Q_2)$$

by observation 3, weakening and exchange

$$= \{Q' \mid \Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \vdash Q \geq_1 Q_1\} \cap \{Q' \mid \Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \vdash Q \geq_1 Q_2\} \quad \text{by lemma 23,}$$

meaning that $\Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \vdash Q \geq_1 Q_1$ and $\Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \vdash Q \geq_1 Q_2$. Then after one step, the algorithm terminates by the induction hypothesis. In other words, $\exists Q'$ s.t. $\Gamma, \vec{\beta}^-_1, \vec{\beta}^-_2 \models Q_1 \vee Q_2 = Q'$, and thus, Rule (\exists^\vee) is applicable.

Case 2. $P_1 = \alpha^+$ and $P_2 = \downarrow N$

Then the set of common upper bounds of $\downarrow N$ and α^+ is empty, and thus, $Q \in \text{UB}(\Gamma \vdash P_1) \cap \text{UB}(\Gamma \vdash P_2)$ gives a contradiction:
 $Q \in \text{UB}(\Gamma \vdash \alpha^+) \cap \text{UB}(\Gamma \vdash \downarrow N)$

$$= \{\exists \vec{\alpha}^+. \alpha^+ \mid \dots\} \cap \{\exists \vec{\beta}^-. \downarrow M' \mid \dots\} \quad \text{by the definition of UB}$$

$$= \emptyset \quad \text{since } \alpha^+ \neq \downarrow M' \text{ for any } M'$$

Case 3. $P_1 = \downarrow N$ and $P_2 = \alpha^+$

Symmetric to case 2

Case 4. $P_1 = \alpha^+$ and $P_2 = \beta^+$ (where $\beta^+ \neq \alpha^+$)

Similarly to case 2, the set of common upper bounds is empty, which leads to the contradiction:

$$Q \in \text{UB}(\Gamma \vdash \alpha^+) \cap \text{UB}(\Gamma \vdash \beta^+)$$

$$= \{\exists \vec{\alpha}^+. \alpha^+ \mid \dots\} \cap \{\exists \vec{\beta}^+. \beta^+ \mid \dots\} \quad \text{by the definition of UB}$$

$$= \emptyset \quad \text{since } \alpha^+ \neq \beta^+$$

Case 5. $P_1 = \alpha^+$ and $P_2 = \alpha^+$

Then the algorithm terminates in one step (Rule (Var^\vee)): $\Gamma \models \alpha^+ \vee \alpha^+ = \alpha^+$.

Case 6. $P_1 = \downarrow M_1$ and $P_2 = \downarrow M_2$

Then on the next step, the algorithm tries to anti-unify $\downarrow M_1$ and $\downarrow M_2$. By lemma 6, to show that the anti-unification algorithm terminates, it suffices to demonstrate that a sound anti-unification solution exists.

Notice that

$$\text{nf}(Q) \in \text{NFUB}(\Gamma \vdash \exists \vec{\beta}^-_1. Q_1) \cap \text{NFUB}(\Gamma \vdash \exists \vec{\beta}^-_2. Q_2)$$

$$\begin{aligned} & \left\{ \exists \vec{\alpha}^-. \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \text{ and } \vec{N} \text{ s.t. } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^-, \\ \Gamma \vdash N_i, \Gamma, \vec{\alpha}^- \vdash M', \text{ and } [\vec{N}/\vec{\alpha}^-] \downarrow M' = \downarrow M_1 \end{array} \right\} \\ &= \bigcap \left\{ \exists \vec{\alpha}^-. \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \text{ and } \vec{N} \text{ s.t. } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^-, \\ \Gamma \vdash \vec{N}_1, \Gamma \vdash \vec{N}_2, \Gamma, \vec{\alpha}^- \vdash M', \text{ and } [\vec{N}/\vec{\alpha}^-] \downarrow M' = \downarrow M_2 \end{array} \right\} \\ &= \left\{ \exists \vec{\alpha}^-. \downarrow M' \mid \begin{array}{l} \text{for } \vec{\alpha}^-, M', \vec{N}_1 \text{ and } \vec{N}_2 \text{ s.t. } \mathbf{ord}\{\vec{\alpha}^-\} \text{ in } M' = \vec{\alpha}^-, \\ \Gamma \vdash \vec{N}_1, \Gamma \vdash \vec{N}_2, \Gamma, \vec{\alpha}^- \vdash M', [\vec{N}_1/\vec{\alpha}^-] \downarrow M' = \downarrow M_1, \text{ and } [\vec{N}_2/\vec{\alpha}^-] \downarrow M' = \downarrow M_2 \end{array} \right\} \end{aligned}$$

The fact that the latter set is non-empty means that there exist $\vec{\alpha}^-, M', \vec{N}_1$ and \vec{N}_2 such that

- (i) $\Gamma, \vec{\alpha}^- \vdash M'$,
- (ii) $\Gamma \vdash \vec{N}_1$ and $\Gamma \vdash \vec{N}_2$,
- (iii) $[\vec{N}_1/\vec{\alpha}^-] \downarrow M' = \downarrow M_1$ and $[\vec{N}_2/\vec{\alpha}^-] \downarrow M' = \downarrow M_2$

For each negative variable α^- from $\vec{\alpha}^-$, let us choose a fresh negative antiunification variable $\hat{\alpha}^-$, and denote the list of these variables as $\vec{\hat{\alpha}}^-$. Let us show that $(\vec{\hat{\alpha}}^-, [\vec{\hat{\alpha}}^-/\vec{\alpha}^-] \downarrow M', \vec{N}_1/\vec{\hat{\alpha}}^-, \vec{N}_2/\vec{\hat{\alpha}}^-)$ is a sound anti-unifier of $\downarrow M_1$ and $\downarrow M_2$ in context Γ :

- $\vec{\hat{\alpha}}^-$ is negative by construction,
- $\Gamma; \vec{\hat{\alpha}}^- \vdash [\vec{\hat{\alpha}}^-/\vec{\alpha}^-] \downarrow M'$ because $\Gamma, \vec{\alpha}^- \vdash \downarrow M'$ **Ilya: lemma!**,
- $\Gamma; \cdot \vdash (\vec{N}_1/\vec{\hat{\alpha}}^-) : \vec{\hat{\alpha}}^-$ because $\Gamma \vdash \vec{N}_1$ and $\Gamma; \cdot \vdash (\vec{N}_2/\vec{\hat{\alpha}}^-) : \vec{\hat{\alpha}}^-$ because $\Gamma \vdash \vec{N}_2$,

- $[\vec{N}_1/\vec{\alpha}^-][\vec{\alpha}^-/\vec{\alpha}^-]\downarrow M' = [\vec{N}_1/\vec{\alpha}^-]\downarrow M' = \downarrow M_1$; analogously, $[\vec{N}_2/\vec{\alpha}^-][\vec{\alpha}^-/\vec{\alpha}^-]\downarrow M' = i[\vec{N}_2/\vec{\alpha}^-]\downarrow M' = \downarrow M_2$.

Then by the completeness of the anti-unification (lemma 6), the anti-unification algorithm terminates, so is the Least Upper Bound algorithm invoking it. \square

Lemma 27 (Initiality of the Least Upper Bound). *For normalized types $\Gamma \vdash P_1$, $\Gamma \vdash P_2$, and $\Gamma \vdash Q$ such that $\Gamma \vdash Q \geq_1 P_1$ and $\Gamma \vdash Q \geq_1 P_2$, If $\Gamma \models P_1 \vee P_2 = Q'$ then $\Gamma \vdash Q \geq_1 Q'$.*

Proof. By induction on a pair (P_1, P_2) , similarly to the proof of lemma 26.

Let us consider the cases what P_1 and P_2 are (i.e. the last rules to infer $\Gamma \vdash P_1$).

Case 1. $P_1 = \exists \vec{\beta}^{-}_1. Q_1$, $P_2 = \exists \vec{\beta}^{-}_2. Q_2$ where $\vec{\beta}^{-}_1$ or $\vec{\beta}^{-}_2$ is not empty

Then by the same reasoning as in case 1 of the proof of lemma 26, $\Gamma, \vec{\beta}^{-}_1, \vec{\beta}^{-}_2 \vdash Q \geq_1 Q_1$ and $\Gamma, \vec{\beta}^{-}_1, \vec{\beta}^{-}_2 \vdash Q \geq_1 Q_2$.

On the other hand, the inversion of $\Gamma \models \exists \vec{\beta}^{-}_1. Q_1 \vee \exists \vec{\beta}^{-}_2. Q_2 = Q'$ gives us $\Gamma, \vec{\beta}^{-}_1, \vec{\beta}^{-}_2 \models Q_1 \vee Q_2 = Q'$. Hence, by the induction hypothesis, $\Gamma, \vec{\beta}^{-}_1, \vec{\beta}^{-}_2 \vdash Q \geq_1 Q'$.

Since both Q and Q' are sound, $\Gamma \vdash Q$ and $\Gamma \vdash Q'$, and therefore, $\Gamma, \vec{\beta}^{-}_1, \vec{\beta}^{-}_2 \vdash Q \geq_1 Q'$ can be strengthened to $\Gamma \vdash Q \geq_1 Q'$.

Ilya: lemma!

Case 2. ($P_1 = \alpha^+$ and $P_2 = \downarrow N$) or ($P_1 = \downarrow N$ and $P_2 = \alpha^+$) or ($P_1 = \alpha^+$ and $P_2 = \beta^+$)

By the same argument as in case 2 of the proof of lemma 26, the set of common supertypes of P_1 and P_2 is empty, hence contradiction.

Case 3. $P_1 = \alpha^+$ and $P_2 = \alpha^+$

Since $Q \in \text{UB}(\Gamma \vdash \alpha^+)$, $Q = \exists \vec{\alpha}^-. \alpha^+$. Then $\Gamma \vdash \exists \vec{\alpha}^-. \alpha^+ \geq_1 \alpha^+$ by Rule (\exists^{\geq_1}): $\vec{\alpha}^-$ can be instantiated with arbitrary negative types (for example $\forall \beta^+. \uparrow \beta^+$), since the substitution for unused variables does not change the term $[\vec{N}/\vec{\alpha}^-]\alpha^+ = \alpha^+$, and then $\Gamma \vdash \alpha^+ \geq_1 \alpha^+$ by Rule (Var^{\geq_1}).

Case 4. $P_1 = \downarrow M_1$ and $P_2 = \downarrow M_2$

By the same reasoning as in case 6 of the proof of lemma 26, $\mathbf{nf}(Q) = \exists \vec{\alpha}^-. \downarrow M'$ for some $\vec{\alpha}^-$ and $\downarrow M'$ such that there exist \vec{N}_1 and \vec{N}_2 such that:

- (i) $\Gamma, \vec{\alpha}^- \vdash M'$,
- (ii) $\Gamma \vdash \vec{N}_1$ and $\Gamma \vdash \vec{N}_2$,
- (iii) $[\vec{N}_1/\vec{\alpha}^-]\downarrow M' = \downarrow M_1$ and $[\vec{N}_2/\vec{\alpha}^-]\downarrow M' = \downarrow M_2$

For each negative variable α^- from $\vec{\alpha}^-$, let us choose a fresh negative antiunification variable $\hat{\alpha}^-$, and denote the list of these variables as $\vec{\hat{\alpha}}^-$. As shown in case 6 of the proof of lemma 26, $(\vec{\hat{\alpha}}^-, [\vec{\hat{\alpha}}^-/\vec{\alpha}^-]\downarrow M', \vec{N}_1/\vec{\hat{\alpha}}^-, \vec{N}_2/\vec{\hat{\alpha}}^-)$ is a sound anti-unifier of $\downarrow M_1$ and $\downarrow M_2$ in context Γ .

By the inversion of $\Gamma \models \downarrow M_1 \vee \downarrow M_2 = Q'$, we conclude that $Q' = \exists \vec{\beta}^-. [\vec{\beta}^-/\Xi]P$, where $(\Xi, P, \hat{\tau}_1, \hat{\tau}_2)$ is the result of the antiunification of $\downarrow M_1$ and $\downarrow M_2$ in context Γ .

Then by the initiality of the anti-unification (lemma 7), there exists $\hat{\tau}$ such that $\Gamma; \Xi \vdash \hat{\tau} : \vec{\hat{\alpha}}^-$ and $[\hat{\tau}][\vec{\hat{\alpha}}^-/\vec{\alpha}^-]\downarrow M' = P$.

Let σ be a sequential Kleisli composition of the following substitutions: (i) $\vec{\hat{\alpha}}^-/\vec{\alpha}^-$, (ii) $\hat{\tau}$, and (iii) $\vec{\beta}^-/\Xi$. Notice that $\Gamma, \vec{\beta}^- \vdash \sigma : \vec{\alpha}^-$ and $[\sigma]\downarrow M' = [\vec{\beta}^-/\Xi][\hat{\tau}][\vec{\hat{\alpha}}^-/\vec{\alpha}^-]\downarrow M' = [\vec{\beta}^-/\Xi]P$. In particular, from the reflexivity of subtyping: $\Gamma, \vec{\beta}^- \vdash [\sigma]\downarrow M' \geq_1 [\vec{\beta}^-/\Xi]P$.

It allows us to show $\Gamma \vdash \mathbf{nf}(Q) \geq_1 Q'$, i.e. $\Gamma \vdash \exists \vec{\alpha}^-. \downarrow M' \geq_1 \exists \vec{\beta}^-. [\vec{\beta}^-/\Xi]P$, by applying Rule (\exists^{\geq_1}), instantiating $\vec{\alpha}^-$ with respect to σ . Finally, $\Gamma \vdash Q \geq_1 Q'$ since $\Gamma \vdash \mathbf{nf}(Q) \simeq_1^s Q$, and equivalence implies subtyping by **Ilya: lemma**.

\square

Lemma 28 (Soundness of Upgrade). *For $\Delta \subseteq \Gamma$, suppose that $\mathbf{upgrade} \Gamma \vdash P$ to $\Delta = Q$.*

- (i) $\Delta \vdash Q$
- (ii) $\Gamma \vdash Q \geq_1 P$

Lemma 29 (Completeness of Upgrade). *For $\Delta \subseteq \Gamma$, $\Gamma \vdash P$ and $\Delta \vdash Q'$, such that $\Gamma \vdash Q' \geq_1 P$, there exists Q s.t. $\mathbf{upgrade} \Gamma \vdash P$ to $\Delta = Q$, and $\Delta \vdash Q' \geq_1 Q$.*