

Introduction


Once each month from July to October of 2017 we surveyed 32 Michigan lakes. At the start of each month, water conditions were recorded, samples were taken, and Solid Phase Adsorption Toxin Tracking (SPATT) bags were deployed and collected. Analysis of the water samples included analysis of nutrients, qPCR for cyanobacteria genes, and microcystins and other toxins. SPATTs are a new way of tracking toxins that provide a time weighted relative measure.

Microcystin and Aquatic Toxin Summary



Enzyme Linked Immunosorbent Assay (ELISA) is a most commonly used analytical method for microcystin (MC) detection. Further data will demonstrate the validity of the test across multiple aquatic environments. We will be ultimately comparing ELISA with Mass Spectrometry (MS) results because it is the most precise and accurate method of quantification. Different MC congeners were measured and added as a sum (Sum of MC Congeners) to be compared with ELISA.

The information collected reflects the conditions at the time of sampling, but toxins recovered from the SPATT bags are indicative of toxin levels during the weeks between sampling. The SPATT bags were deployed for the month-long period between sampling events and then collected. SPATT data are represented by the total toxins collected during the period. The reported concentrations are from the analytical procedures used on the SPATT bags and are not directly indicative of average toxin levels. There is no USEPA guidance for this procedure yet and we will use the 2017 and 2018 data to provide interim guidance.

Grab Sample



SPATTS



qPCR


QPCR is a DNA test to rapidly measure the amount of total cyanobacteria (16s rRNA) and toxin genes (mcyE) present. Phytoxigene™ CyanoDtec test was performed with Applied Biosystem StepOnePlus PCR. Total Cyanobacteria 16s rRNA and toxin gene assay were analyzed in parallel for each month of grab samples. Data for each month is listed in table below. CyrA and SxtA were not detected for this year. The calculated values are expressed as “GeneCopies/mL”.

16s rRNA is a ubiquitous gene which is found in mostly all cyanobacteria. The 16s rRNA gene copies is used to measure relatively how much cyanobacteria is found in your lake. McyE gene is one of the few genes responsible of producing microcystin. Presence of these genes do not indicate that the toxins are present. Detection of McyE gene would indicate that the lake has the potential to produce microcystin and is advised to continue monitoring the lake. We do not become concerned until total cyanobacteria (16s rRNA) are above 200,000 and toxin genes (McyE) are above 10,000.

Month	16s rRNA (copies/mL)	mcyE (copies/mL)
July	14706	NA
August	3040181	5
September	67700	38
October	85809	12

Nutrient Summary

The chart and table show the nutrient levels below. The key nutrients we measure are the forms of nitrogen and phosphorous. Phosphorus has long been known to increase the algal and cyanobacteria growth.



Month	Orthophosphate (mg P/L)	Nitrate + Nitrite (mg N/L)	Ammonia (mg N/L)	Total Phosphorus (mg P/L)
July	0.0080	0	0.3710	0.025
August	0.0052	0	0.2719	0.042
September	0.0073	0	0.0275	0.045
October	0.0096	0	0.0000	0.085

Water Parameters

Month	Temperature (°C)	Turbidity (NTU)	pH	Dissolve Oxygen (mg/L)	Conductivity (S)	Chlorophyll-a (R)
July	23.52	2.41	8.52	8.57	165	
August	27.23	29.60	8.40	12.35	317	
September	20.01	4.97	8.11	10.74	330	
October	16.32	1.56	8.82	8.41	163	