

## Lecture 14: Compatible Properties in Histories

Fri Sep 20 2019

We don't want to make the mistake of discussing incompatible properties. To do this with histories, we start with some family of histories  $\{Y^\alpha\}$  and we require  $Y^\alpha Y^\beta = Y^\beta Y^\alpha$ ,  $\forall \alpha \beta$ . This is a complete family of histories, so  $\sum_{\vec{\alpha}} Y^{\vec{\alpha}} = \tilde{I}$ . Therefore, **logical negation** is  $\neg Y^{\vec{\alpha}} = \tilde{I} - Y^{\vec{\alpha}}$ .

**Example.** Coin toss:  $\{(H, H), (H, T), (T, H), (T, T)\}$ . The negation of  $(H, H)$  is  $\neg(H, H) = \{(H, T), (T, H), (T, T)\}$   $\diamond$

**Example.**

$$\neg[z+] \odot [x+] = [z-] \odot [x+] + [z+] \odot [x-] + [z-] \odot [x-] \quad (1)$$

$\diamond$

We can also have **conjunction**,  $Y \wedge Y' = YY'$ .

**Example.**

$$Y = [z+]_0 \odot I_1 \quad (2)$$

$$Y' = I_0 \odot [x+]_1 \quad (3)$$

$$YY' = [z+] \odot [x+] \quad (4)$$

$\diamond$

Additionally, **disjunction** is defined by  $Y \vee Y' = Y + Y' - YY'$ .

### 0.1 Chainket

If we start in a pure state, we can define a product history

$$Y^{\vec{\alpha}} = [\psi_0] \odot P_1^{\alpha_1} \odot \dots \odot P_f^{\alpha_f} \in \tilde{\mathcal{H}} \quad (5)$$

**Definition 1.** A **chainket** is defined by

$$|\vec{\alpha}\rangle = P_f^{\alpha_f} T_{f,f-1} \dots T_{21} P_1^{\alpha_1} T_{10} |\psi_0\rangle \quad (6)$$

**Theorem 1.** Generalized Born Rule:

$$Pr(\vec{\alpha}) = \langle \vec{\alpha} | \vec{\alpha} \rangle \quad (7)$$

Is it correct? Let's check some cases.

#### 0.1.1 Two-Time History

$$Y^k = [\psi_0] \odot [\phi_1^k] \quad (8)$$

All of our states start at  $|\psi_0\rangle$ , so we need to define a complete set of histories by adding all the things that don't start there:

$$Z = (I - [\psi_0]) \odot I_1 \quad (9)$$

The chainket for this state is

$$|k\rangle = [\phi_1^k] T_{10} |\psi_0\rangle \quad (10)$$

$$\langle k|k\rangle = \langle \psi_0 | T_{01} [\phi_1^k] \langle \phi_1^k | T_{10} |\psi_0\rangle = |\langle \phi_1^k | \psi_1\rangle|^2 = Pr([\phi_1^k] | \psi_0) \quad (11)$$

N.B.

$$|\langle \phi_1^k | \psi_1\rangle|^2 = \langle \phi_1^k | \psi_1\rangle \langle \psi_1 | \phi_1^k\rangle = \langle \psi_1 | (\phi_1^k)^2 | \psi_1\rangle = \langle \psi_1 | [\phi_1^k] | \psi_1\rangle \quad (12)$$

### 0.1.2 Unitary History

$$|\psi_0\rangle \rightarrow |\psi_1\rangle = T_{10} |\psi_0\rangle \rightarrow |\psi_2\rangle = T_{21} |\psi_1\rangle = T_{20} |\psi_0\rangle \quad (13)$$

$$U = [\psi_0] \odot [\psi_1] \odot [\psi_2] \quad (14)$$

$$|U\rangle = |\psi_2\rangle \langle \psi_2 | T_{21} |\psi_1\rangle \langle \psi_1 | T_{10} |\psi_0\rangle \quad (15)$$

There's another way of thinking about this:

$$|U\rangle = |\psi_2\rangle \langle \psi_2 | T_{21} |\psi_1\rangle \langle \psi_1 | T_{10} |\psi_0\rangle \quad (16)$$

so

$$Pr(U) = \langle U|U\rangle = \langle \psi_2 | \psi_2\rangle = 1 \quad (17)$$