Manifold-regression to predict from MEG/EEG brain signals without source modeling

D. Sabbagh, P. Ablin, G. Varoquaux, A. Gramfort, D.Engemann

September 2019

JDSE2019 - 4th edition
Junior Conference on Data Science and Engineering

Non-invasive measure of brain activity

Non-invasive measure of brain activity





First EEG recordings in 1929 by H. Berger



Hôpital La Timone
Marseille, France

Objective: predict a variable from M/EEG brain signals

Objective: predict a variable from M/EEG brain signals

- We want to predict a continuous variable
- From M/EEG brain signals

Objective: predict a variable from M/EEG brain signals

- We want to predict a continuous variable
- From M/EEG brain signals



Model: generative model of M/EEG data

Model: generative model of M/EEG data

We measure M/EEG signal of subject $i = 1 \dots N$ on P channels:

$$\boxed{\boldsymbol{x}_i(t) = \boldsymbol{A}\ \boldsymbol{s}_i(t) + \boldsymbol{n}_i(t)} \in \mathbb{R}^P \quad t = 1 \dots T$$

- *mixing matrix* $\boldsymbol{A} = [\boldsymbol{a}_1, \dots, \boldsymbol{a}_Q] \in \mathbb{R}^{P \times Q}$ fixed accross subjects
- *source patterns* $\boldsymbol{a}_j \in \mathbb{R}^P, j = 1 \dots Q$
- *source vector* $\boldsymbol{s}_i(t) \in \mathbb{R}^Q$
- *noise* $\boldsymbol{n}_i(t) \in \mathbb{R}^P$

Model: generative model of M/EEG data

We measure M/EEG signal of subject $i = 1 \dots N$ on P channels:

$$\boxed{\boldsymbol{x}_i(t) = \boldsymbol{A}\ \boldsymbol{s}_i(t) + \boldsymbol{n}_i(t)} \in \mathbb{R}^P \quad t = 1 \dots T$$

- *mixing matrix* $\boldsymbol{A} = [\boldsymbol{a}_1, \dots, \boldsymbol{a}_Q] \in \mathbb{R}^{P \times Q}$ fixed accross subjects
- *source patterns* $\boldsymbol{a}_j \in \mathbb{R}^P, j = 1 \dots Q$
- *source vector* $\boldsymbol{s}_i(t) \in \mathbb{R}^Q$
- *noise* $\boldsymbol{n}_i(t) \in \mathbb{R}^P$

Typically, $N \simeq 10^2$, $T \simeq 10^5$, $P \simeq 10^2$, $Q \simeq 10^4$

Model: generative model of M/EEG data

We measure M/EEG signal of subject $i = 1 \dots N$ on P channels:

$$\boxed{\boldsymbol{x}_i(t) = \boldsymbol{A}\ \boldsymbol{s}_i(t) + \boldsymbol{n}_i(t)} \in \mathbb{R}^P \quad t = 1 \dots T$$

- *mixing matrix* $\boldsymbol{A} = [\boldsymbol{a}_1, \dots, \boldsymbol{a}_Q] \in \mathbb{R}^{P \times Q}$ fixed accross subjects
- *source patterns* $\boldsymbol{a}_j \in \mathbb{R}^P, j = 1 \dots Q$
- *source vector* $\boldsymbol{s}_i(t) \in \mathbb{R}^Q$
- *noise* $\boldsymbol{n}_i(t) \in \mathbb{R}^P$

Typically, $N \simeq 10^2$, $T \simeq 10^5$, $P \simeq 10^2$, $Q \simeq 10^4$

Under stationnarity and gaussianity assumptions, we can represent band-pass filtered signal by its second-order statistics

$$\boldsymbol{C}_i = \mathbb{E}[\boldsymbol{x}_i(t)\boldsymbol{x}_i(t)^\top] \simeq \frac{\boldsymbol{X}_i \boldsymbol{X}_i^\top}{T} \in \mathbb{R}^{P \times P} \quad \text{with} \quad \boldsymbol{X}_i \in \mathbb{R}^{P \times T}$$

Model: generative model of target variable

We want to predict a continuous variable: $\boxed{y_i = \sum_{j=1}^{Q} \alpha_j f(p_{i,j})} \in \mathbb{R}$

with $p_{i,j} = \mathbb{E}_t[s_{i,j}^2(t)]$ band-power of sources

Model: generative model of target variable

We want to predict a continuous variable: $\boxed{y_i = \sum_{j=1}^{Q} \alpha_j f(p_{i,j})} \in \mathbb{R}$

with $p_{i,j} = \mathbb{E}_t[s_{i,j}^2(t)]$ band-power of sources

- linear $y_i = \sum_{j=1}^{Q} \alpha_j p_{i,j}$

Model: generative model of target variable

We want to predict a continuous variable: $\boxed{y_i = \sum_{j=1}^{Q} \alpha_j f(p_{i,j})} \in \mathbb{R}$

with $p_{i,j} = \mathbb{E}_t[s_{i,j}^2(t)]$ band-power of sources

- linear $y_i = \sum_{j=1}^{Q} \alpha_j p_{i,j}$

Euclidean vectorization leads to consistent model

$y_i = \sum_{k \leq l} \Theta_{k,l} \boldsymbol{C}_i(k,l)$ *i.e.* y_i is linear in coeff. of Upper($\boldsymbol{C}_i$)

Model: generative model of target variable

We want to predict a continuous variable: $\boxed{y_i = \sum_{j=1}^{Q} \alpha_j f(p_{i,j})} \in \mathbb{R}$

with $p_{i,j} = \mathbb{E}_t[s_{i,j}^2(t)]$ band-power of sources

- linear $y_i = \sum_{j=1}^{Q} \alpha_j p_{i,j}$

Euclidean vectorization leads to consistent model

$y_i = \sum_{k \leq l} \Theta_{k,l} \boldsymbol{C}_i(k,l)$ *i.e.* y_i is linear in coeff. of Upper($\boldsymbol{C}_i$)

- log linear $y_i = \sum_{j=1}^{Q} \alpha_j \log(p_{i,j})$

Model: generative model of target variable

We want to predict a continuous variable: $\boxed{y_i = \sum_{j=1}^{Q} \alpha_j f(p_{i,j})} \in \mathbb{R}$

with $p_{i,j} = \mathbb{E}_t[s_{i,j}^2(t)]$ band-power of sources

- linear $y_i = \sum_{j=1}^{Q} \alpha_j p_{i,j}$

Euclidean vectorization leads to consistent model

$y_i = \sum_{k \leq l} \Theta_{k,l} \boldsymbol{C}_i(k,l)$ *i.e.* y_i is linear in coeff. of Upper($\boldsymbol{C}_i$)

- log linear $y_i = \sum_{j=1}^{Q} \alpha_j \log(p_{i,j})$

  $\boldsymbol{C}_i$ live on a Riemannian manifold so can't be naively vectorized

Riemannian matrix manifolds (in a nutshell)



Riemannian matrix manifolds (in a nutshell)



[Absil & al. Optimization algorithms on matrix manifolds. 2009]

Riemannian matrix manifolds (in a nutshell)



Vectorization operator:

$$\mathcal{P}_{\boldsymbol{M}}(\boldsymbol{M'}) = \phi_{\boldsymbol{M}}(\mathrm{Log}_{\boldsymbol{M}}(\boldsymbol{M'})) \simeq \mathrm{Upper}(\mathrm{Log}_{\boldsymbol{M}}(\boldsymbol{M'}))$$

$$d(\boldsymbol{M}, \boldsymbol{M'}) = \|\mathcal{P}_{\boldsymbol{M}}(\boldsymbol{M'})\|_2 + o(\|\mathcal{P}_{\boldsymbol{M}}(\boldsymbol{M'})\|_2)$$

$$d(\boldsymbol{M}_i, \boldsymbol{M}_j) \simeq \|\mathcal{P}_{\overline{\boldsymbol{M}}}(\boldsymbol{M}_i) - \mathcal{P}_{\overline{\boldsymbol{M}}}(\boldsymbol{M}_j)\|_2$$

Vectorization operator key for ML

[Absil & al. Optimization algorithms on matrix manifolds. 2009]

Regression on matrix manifolds

Given a training set of samples $\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N \in \mathcal{M}$ and target continuous variables $y_1, \ldots, y_N \in \mathbb{R}$:

Regression on matrix manifolds

Given a training set of samples $\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N \in \mathcal{M}$ and target continuous variables $y_1, \ldots, y_N \in \mathbb{R}$:

- compute the mean of the samples $\overline{\boldsymbol{M}} = \mathrm{Mean}_d(\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N)$

Regression on matrix manifolds

Given a training set of samples $\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N \in \mathcal{M}$ and target continuous variables $y_1, \ldots, y_N \in \mathbb{R}$:

- compute the mean of the samples $\overline{\boldsymbol{M}} = \mathrm{Mean}_d(\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N)$
- compute the vectorization of the samples *w.r.t.* this mean: $\boldsymbol{v}_1, \ldots, \boldsymbol{v}_N \in \mathbb{R}^K$ as $\boldsymbol{v}_i = \mathcal{P}_{\overline{\boldsymbol{M}}}(\boldsymbol{M}_i)$

Regression on matrix manifolds

Given a training set of samples $\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N \in \mathcal{M}$ and target continuous variables $y_1, \ldots, y_N \in \mathbb{R}$:

- compute the mean of the samples $\overline{\boldsymbol{M}} = \mathrm{Mean}_d(\boldsymbol{M}_1, \ldots, \boldsymbol{M}_N)$
- compute the vectorization of the samples *w.r.t.* this mean: $\boldsymbol{v}_1, \ldots, \boldsymbol{v}_N \in \mathbb{R}^K$ as $\boldsymbol{v}_i = \mathcal{P}_{\overline{\boldsymbol{M}}}(\boldsymbol{M}_i)$
- use those vectors as features in regularized linear regression algorithm (e.g. ridge regression) with parameters $\boldsymbol{\beta} \in \mathbb{R}^K$ assuming that $y_i \simeq \boldsymbol{v}_i^\top \boldsymbol{\beta}$

Distance and invariance on positive matrix manifolds

[Bhatia. Positive Definite Matrices. Princeton University Press, 2007]

Distance and invariance on positive matrix manifolds

- **Manifold of positive definite matrices**: $\boldsymbol{M}_i = \boldsymbol{C}_i \in \mathcal{S}_P^{++}$

[Bhatia. Positive Definite Matrices. Princeton University Press, 2007]

Distance and invariance on positive matrix manifolds

- **Manifold of positive definite matrices**: $\boldsymbol{M}_i = \boldsymbol{C}_i \in \mathcal{S}_P^{++}$
- **Geometric distance**:

  $$d_G(\boldsymbol{S}, \boldsymbol{S'}) = \| \log(\boldsymbol{S}^{-1}\boldsymbol{S'}) \|_F = \left[\sum_{i=1}^{P} \log^2 \lambda_k \right]^{\frac{1}{2}}$$

  where $\lambda_k, k = 1 \ldots P$ are the real eigenvalues of $\boldsymbol{S}^{-1}\boldsymbol{S'}$.
- **Tangent Space Projection**:
  $\mathcal{P}_{\boldsymbol{S}}(\boldsymbol{S'}) = \text{Upper}(\log(\boldsymbol{S}^{-\frac{1}{2}}\boldsymbol{S'}\boldsymbol{S}^{-\frac{1}{2}}))$

[Bhatia. Positive Definite Matrices. Princeton University Press, 2007]

Distance and invariance on positive matrix manifolds

- **Manifold of positive definite matrices**: $\boldsymbol{M}_i = \boldsymbol{C}_i \in \mathcal{S}_P^{++}$
- **Geometric distance**:

  $$d_G(\boldsymbol{S}, \boldsymbol{S'}) = \| \log(\boldsymbol{S}^{-1}\boldsymbol{S'}) \|_F = \left[\sum_{i=1}^{P} \log^2 \lambda_k \right]^{\frac{1}{2}}$$

  where $\lambda_k, k = 1 \ldots P$ are the real eigenvalues of $\boldsymbol{S}^{-1}\boldsymbol{S'}$.
- **Tangent Space Projection**:
  $\mathcal{P}_{\boldsymbol{S}}(\boldsymbol{S'}) = \text{Upper}(\log(\boldsymbol{S}^{-\frac{1}{2}}\boldsymbol{S'}\boldsymbol{S}^{-\frac{1}{2}}))$
- **Affine invariance property**:
  For $\boldsymbol{W}$ invertible, $d_G(\boldsymbol{W}^\top \boldsymbol{S} \boldsymbol{W}, \boldsymbol{W}^\top \boldsymbol{S'} \boldsymbol{W}) = d_G(\boldsymbol{S}, \boldsymbol{S'})$

[Bhatia. Positive Definite Matrices. Princeton University Press, 2007]

Distance and invariance on positive matrix manifolds

- **Manifold of positive definite matrices**: $\boldsymbol{M}_i = \boldsymbol{C}_i \in \mathcal{S}_P^{++}$
- **Geometric distance**:
  $$d_G(\boldsymbol{S}, \boldsymbol{S'}) = \| \log(\boldsymbol{S}^{-1}\boldsymbol{S'}) \|_F = \left[\sum_{i=1}^{P} \log^2 \lambda_k \right]^{\frac{1}{2}}$$
  where $\lambda_k, k = 1 \dots P$ are the real eigenvalues of $\boldsymbol{S}^{-1}\boldsymbol{S'}$.
- **Tangent Space Projection**:
  $\mathcal{P}_{\boldsymbol{S}}(\boldsymbol{S'}) = \text{Upper}(\log(\boldsymbol{S}^{-\frac{1}{2}}\boldsymbol{S'}\boldsymbol{S}^{-\frac{1}{2}}))$
- **Affine invariance property**:
  For $\boldsymbol{W}$ invertible, $d_G(\boldsymbol{W}^\top \boldsymbol{S} \boldsymbol{W}, \boldsymbol{W}^\top \boldsymbol{S'} \boldsymbol{W}) = d_G(\boldsymbol{S}, \boldsymbol{S'})$

Affine invariance is key: working with $\boldsymbol{C}_i$ is then equivalent to working with covariance matrices of sources $\boldsymbol{s}_i$

[Bhatia. Positive Definite Matrices. Princeton University Press, 2007]

Consistency of linear regression in tangent space of $\mathcal{S}_P^{++}$

Consistency of linear regression in tangent space of $\mathcal{S}_P^{++}$

Geometric vectorization

Assume $y_i = \sum_{j=1}^{Q} \alpha_j \log(p_{i,j})$. Denote $\overline{\boldsymbol{C}} = \text{Mean}_G(\boldsymbol{C}_1, \ldots, \boldsymbol{C}_N)$ and $\boldsymbol{v}_i = \mathcal{P}_{\overline{\boldsymbol{C}}}(\boldsymbol{C}_i)$. Then, the relationship between y_i and $\boldsymbol{v}_i$ is linear.

Consistency of linear regression in tangent space of $\mathcal{S}_P^{++}$

Geometric vectorization

Assume $y_i = \sum_{j=1}^{Q} \alpha_j \log(p_{i,j})$. Denote $\overline{\boldsymbol{C}} = \text{Mean}_G(\boldsymbol{C}_1, \ldots, \boldsymbol{C}_N)$ and $\boldsymbol{v}_i = \mathcal{P}_{\overline{\boldsymbol{C}}}(\boldsymbol{C}_i)$. Then, the relationship between y_i and $\boldsymbol{v}_i$ is linear.

We generate i.i.d. samples following the log linear generative model. $\boldsymbol{A} = \exp(\mu \boldsymbol{B})$ with $\boldsymbol{B} \in \mathbb{R}^{P \times P}$ random



And in the real world ?

We investigated 3 violations from previous idealized model:

And in the real world ?

We investigated 3 violations from previous idealized model:

- <u>Noise in target variable</u>

$$y_i = \sum_j \alpha_j \log(p_{ij}) + \varepsilon_i \ , \quad \text{with} \quad \varepsilon_i \sim \mathcal{N}(0, \sigma^2)$$

And in the real world ?

We investigated 3 violations from previous idealized model:

- Noise in target variable

$$y_i = \sum_j \alpha_j \log(p_{ij}) + \varepsilon_i \ , \quad \text{with} \quad \varepsilon_i \sim \mathcal{N}(0, \sigma^2)$$

- Subject-dependent mixing matrix

$$\boldsymbol{A}_i = \boldsymbol{A} + \boldsymbol{E}_i \ , \quad \text{with entries of} \quad \boldsymbol{E}_i \sim \mathcal{N}(0, \sigma^2)$$

And in the real world ?

We investigated 3 violations from previous idealized model:

- Noise in target variable

$$y_i = \sum_j \alpha_j \log(p_{ij}) + \varepsilon_i \ , \quad \text{with} \quad \varepsilon_i \sim \mathcal{N}(0, \sigma^2)$$

- Subject-dependent mixing matrix

$$\boldsymbol{A}_i = \boldsymbol{A} + \boldsymbol{E}_i \ , \quad \text{with entries of} \quad \boldsymbol{E}_i \sim \mathcal{N}(0, \sigma^2)$$





And in the real world ?

- Rank-deficient signals (*e.g.* cleaning process): $\boldsymbol{C}_i \in \mathcal{S}^+_{P,R}$

[Massart, Absil. Quotient geometry with simple geodesics for the manifold of fixed-rank positive-semidefinite matrices. Technical report, UCLouvain, 2018]

And in the real world ?

- Rank-deficient signals (*e.g.* cleaning process): $\boldsymbol{C}_i \in \mathcal{S}^+_{P,R}$

We can manipulate them in their native manifolds $\mathcal{S}^+_{P,R}$:

- **Wasserstein distance**:
  $d_W(\boldsymbol{S}, \boldsymbol{S}') = \left[\mathrm{Tr}(\boldsymbol{S}) + \mathrm{Tr}(\boldsymbol{S}') - 2\mathrm{Tr}((\boldsymbol{S}^{\frac{1}{2}}\boldsymbol{S}'\boldsymbol{S}^{\frac{1}{2}})^{\frac{1}{2}})\right]^{\frac{1}{2}}$
- **Tangent Space Projection**:
  $\mathcal{P}_{\boldsymbol{Y}\boldsymbol{Y}^\top}(\boldsymbol{Y}'\boldsymbol{Y}'^\top) = \mathrm{vect}(\boldsymbol{Y}'\boldsymbol{Q}^* - \boldsymbol{Y}) \in \mathbb{R}^{PR}$
  where $\boldsymbol{U}\boldsymbol{\Sigma}\boldsymbol{V}^\top = \boldsymbol{Y}^\top\boldsymbol{Y}'$, $\boldsymbol{Q}^* = \boldsymbol{V}\boldsymbol{U}^\top$
- **Orthogonal invariance property**:
  For $\boldsymbol{W}$ orthogonal, $d_W(\boldsymbol{W}^\top\boldsymbol{S}\boldsymbol{W}, \boldsymbol{W}^\top\boldsymbol{S}'\boldsymbol{W}) = d_W(\boldsymbol{S}, \boldsymbol{S}')$

[Massart, Absil. Quotient geometry with simple geodesics for the manifold of fixed-rank positive-semidefinite matrices. Technical report, UCLouvain, 2018]

And in the real world ?

- Rank-deficient signals (*e.g.* cleaning process): $\boldsymbol{C}_i \in \mathcal{S}^+_{P,R}$

We can manipulate them in their native manifolds $\mathcal{S}^+_{P,R}$:

- **Wasserstein distance**:
  $d_W(\boldsymbol{S}, \boldsymbol{S}') = \left[\mathrm{Tr}(\boldsymbol{S}) + \mathrm{Tr}(\boldsymbol{S}') - 2\mathrm{Tr}((\boldsymbol{S}^{\frac{1}{2}}\boldsymbol{S}'\boldsymbol{S}^{\frac{1}{2}})^{\frac{1}{2}})\right]^{\frac{1}{2}}$
- **Tangent Space Projection**:
  $\mathcal{P}_{\boldsymbol{Y}\boldsymbol{Y}^\top}(\boldsymbol{Y}'\boldsymbol{Y}'^\top) = \mathrm{vect}(\boldsymbol{Y}'\boldsymbol{Q}^* - \boldsymbol{Y}) \in \mathbb{R}^{PR}$
  where $\boldsymbol{U}\boldsymbol{\Sigma}\boldsymbol{V}^\top = \boldsymbol{Y}^\top\boldsymbol{Y}'$, $\boldsymbol{Q}^* = \boldsymbol{V}\boldsymbol{U}^\top$
- **Orthogonal invariance property**:
  For $\boldsymbol{W}$ orthogonal, $d_W(\boldsymbol{W}^\top\boldsymbol{S}\boldsymbol{W}, \boldsymbol{W}^\top\boldsymbol{S}'\boldsymbol{W}) = d_W(\boldsymbol{S}, \boldsymbol{S}')$

Wasserstein vectorization

Assume $y_i = \sum_{j=1}^{Q} \alpha_j \sqrt{p_{i,j}}$ and $\boldsymbol{A}$ orthogonal. Denote $\overline{\boldsymbol{C}} = \mathrm{Mean}_W(\boldsymbol{C}_1, \ldots, \boldsymbol{C}_N)$ and $\boldsymbol{v}_i = \mathcal{P}_{\overline{\boldsymbol{C}}}(\boldsymbol{C}_i)$.
Then the relationship between y_i and $\boldsymbol{v}_i$ is linear.

[Massart, Absil. Quotient geometry with simple geodesics for the manifold of fixed-rank positive-semidefinite matrices. Technical report, UCLouvain, 2018]

Experiment: predict age from MEG data

Experiment: predict age from MEG data

- Task-free MEG recordings from Cam-CAN dataset

Experiment: predict age from MEG data

- Task-free MEG recordings from Cam-CAN dataset
- Age is a dominant driver of cross-person variance in neuroscience data

Experiment: predict age from MEG data

- Task-free MEG recordings from Cam-CAN dataset
- Age is a dominant driver of cross-person variance in neuroscience data
- Dimension: $N = 595, P = 102, T \simeq 520,000, 65 \leq R_i \leq 73$

Experiment: predict age from MEG data

- Task-free MEG recordings from Cam-CAN dataset
- Age is a dominant driver of cross-person variance in neuroscience data
- Dimension: $N = 595, P = 102, T \simeq 520,000, 65 \leq R_i \leq 73$
- Signals is filtered into 9 frequency bands

Experiment: predict age from MEG data

- Task-free MEG recordings from Cam-CAN dataset
- Age is a dominant driver of cross-person variance in neuroscience data
- Dimension: $N = 595, P = 102, T \simeq 520,000, 65 \leq R_i \leq 73$
- Signals is filtered into 9 frequency bands

Conclusion

- Proposed Riemannian method for regression from M/EEG data

Conclusion

- Proposed Riemannian method for regression from M/EEG data
- With theoritical guarantees and empirical robustness

Conclusion

- Proposed Riemannian method for regression from M/EEG data
- With theoritical guarantees and empirical robustness
- Working to translate proposed method into the clinic

Conclusion

- Proposed Riemannian method for regression from M/EEG data
- With theoritical guarantees and empirical robustness
- Working to translate proposed method into the clinic

[Sabbagh, Ablin, Varoquaux, Gramfort, Engemann (2019), *Manifold-regression to predict from MEG/EEG brain signals without source modeling*, Proc. NeurIPS 2019]

`https://arxiv.org/abs/1906.02687`