

First Principle 2017-Fall Homework 3 Solution

Kai-Hsin Wu (吳愷訢)*

*Department of Physics and Center of Theoretical Sciences,
National Taiwan University, Taipei 10607, Taiwan*

1. The following shows the result of Al and Na:

- Al

(1) a_0 using volume optimization:

$$a_0 = 4.05000$$

(2) Variation with different a_0 :

$a_0(\text{\AA})$	$E(\text{eV})$
3.90	-14.541085
3.95	-14.665726
4.00	-14.735369
4.05	-14.757538
4.10	-14.738699
4.15	-14.684395
4.20	-14.599834

(3) the following figure shows the energy (E) v.s. V :




FIG. 1. Al-fcc E-V

* r05222003@ntu.edu.tw

By using third order polyfit, and with the following formula, we can get the bulk-modulus B and the minimum a_0 :

$$B = V \frac{\partial^2}{\partial V^2} E$$

$$V = a_0^3$$

$$a_0 = 4.050723 \text{ \AA}$$

$$B = 74.608739 \text{ GPa}$$

(4) Consider the free electron HF energy density and seek for the minimum of r_s :

$$e^{HF} = \frac{A}{r_s^2} - \frac{\beta}{r_s}$$

$$A = 2.21$$

$$B = 0.916$$

Where the unit of energy is Ry and r_s is in unit of bohr radius (*bohr*). The minimum is at :

$$r_s^* = \frac{2A}{\beta}(\text{bohr}) = 2.553467 \text{ \AA}$$

using following relation by which we consider 4 free electrons per unit-cell, we can estimate the lattice constant a_0 :

$$\frac{4\pi}{3} r_s^3 = n^{-1}$$

$$\frac{N_{free}}{a_0^3} = n$$

$$a_0 = \left(\frac{16\pi}{3} \right)^{1/3} r_s$$

$$\approx 6.1737 \text{ \AA}$$

For Bulk modulus, we have following relation. by which we express Ω in terms of r_s :

$$\frac{N}{\Omega} = n = \frac{3}{4\pi r_s^3}$$

$$B = \Omega \frac{\partial^2 E}{\partial \Omega^2}$$

$$= \frac{1}{12\pi r_s} \left[\frac{-2}{r_s} e^{HF'} + e^{HF''} \right]$$

$$= \frac{1}{6\pi r_s^4} \left[\frac{5A}{r_s} - 2\beta \right]$$

we have the bulk modulus at r_s^*

$$B = \frac{\beta^4}{6\pi(2A)^4} \frac{\beta}{2} (Ry/bohr^3)$$

$$\approx 18.468 GPa$$

- Na

(1) a_0 using volume optimization:

$$a_0 = 4.04669$$

(2) Variation with different a_0 :

$a_0(\text{\AA})$	$E(eV)$
3.9241059	-1.437731
3.9645606	-1.443730
4.0050153	-1.447706
4.0454700	-1.449866
4.0859247	-1.449972
4.1263794	-1.448133
4.1668341	-1.445039
4.2072888	-1.440498

(3) the following figure shows the energy (E) v.s. V :




FIG. 2. Na-fcc E-V

By using third order polyfit, and with the following formula, we can get the bulk-modulus B and the minimum a_0 :

$$B = V \frac{\partial^2}{\partial V^2} E$$

$$V = a_0^3$$

$$a_0 = 4.06802437 \text{ \AA}$$

$$B = 4.83347453 \text{ GPa}$$

(4) Consider the free electron HF energy density and seek for the minimum of r_s :

$$e^{HF} = \frac{A}{r_s^2} - \frac{\beta}{r_s}$$

$$A = 2.21$$

$$B = 0.916$$

Where the unit of energy is *Ry* and r_s is in unit of bohr radius (*bohr*). The minimum is at :

$$r_s^* = \frac{2A}{\beta}(\text{bohr}) = 2.553467 \text{ \AA}$$

using following relation by which we consider 2 free electrons per unit-cell, we can estimate the lattice constant a_0 :

$$\frac{4\pi}{3}r_s^3 = n^{-1}$$

$$\frac{N_{free}}{a_0^3} = n$$

$$a_0 = \left(\frac{8\pi}{3}\right)^{1/3} r_s$$

$$\approx 5.186047 \text{ \AA}$$

For Bulk modulus, we have following relation. by which we express Ω in terms of r_s :

$$\frac{N}{\Omega} = n = \frac{3}{4\pi r_s^3}$$

$$B = \Omega \frac{\partial^2 E}{\partial \Omega^2}$$

$$= \frac{1}{12\pi r_s} \left[\frac{-2}{r_s} e^{HF'} + e^{HF''} \right]$$

$$= \frac{1}{6\pi r_s^4} \left[\frac{5A}{r_s} - 2\beta \right]$$

we have the bulk modulus at r_s^*

$$B = \frac{\beta^4}{6\pi(2A)^4} \frac{\beta}{2} (\text{Ry}/\text{bohr}^3)$$

$$\approx 18.468 \text{ GPa}$$

2. For bcc Fe , doing LDA and GGA method with FM and NM :

- LDA

(1) a_0 using volumn optimization :

$$a_0^{FM} = 2.744294 \text{ \AA}$$

$$a_0^{NM} = 2.694591 \text{ \AA}$$

(2) minimization energy with variated a_0 and are showed in figure below:

FM		NM	
$a_0(\text{\AA})$	$E(\text{eV})$	$a_0(\text{\AA})$	$E(\text{eV})$
2.60	-8.967318	2.55	-8.672935
2.65	-9.121350	2.60	-8.863644
2.70	-9.206677	2.65	-8.964152
2.75	-9.228247	2.70	-8.988743
2.80	-9.202380	2.75	-8.952579
2.85	-9.135795	2.80	-8.867398
2.90	-9.048763	2.85	-8.742378
2.95	-8.935725	2.90	-8.585827




FIG. 3. Fe-bcc LDA E-V

The minimized a_0 and B of bcc Fe with FM and NM are :

$$a_0^{FM} = 2.746787 \text{ \AA}$$

$$B^{FM} = 134.99746 \text{ GPa}$$

$$a_0^{NM} = 2.694740 \text{ \AA}$$

$$B^{NM} = 175.45431 \text{ GPa}$$

We see that the FM case has minimal energy, which shows the ground state is Ferromagnetic.

- GGA

(1) a_0 using volumn optimization :

$$a_0^{FM} = 2.82214 \text{ \AA}$$

$$a_0^{NM} = 2.75412 \text{ \AA}$$

(2) minimization energy with variated a_0 and are showed in figure below:

FM		NM	
$a_0(\text{\AA})$	$E(\text{eV})$	$a_0(\text{\AA})$	$E(\text{eV})$
2.70	-8.009989	2.70	-7.724527
2.75	-8.108553	2.75	-7.757336
2.80	-8.151154	2.80	-7.737622
2.85	-8.158445	2.85	-7.674045
2.90	-8.135495	2.90	-7.575350
2.95	-8.077445	2.95	-7.448175
3.00	-8.000995	3.00	-7.298845




FIG. 4. Fe-bcc GGA E-V

The minimized a_0 and B of bcc Fe with FM and NM are :

$$a_0^{FM} = 2.834269 \text{ \AA}$$

$$B^{FM} = 93.907626 \text{ GPa}$$

$$a_0^{NM} = 2.755061 \text{ \AA}$$

$$B^{NM} = 124.57160 \text{ GPa}$$

We see that the FM case has minimal energy, which consist with the GGA results that the ground state is Ferromagnetic.

3. GGA band gaps of typical semiconductors, the following experiment value comes from [1]

- Si

(1) a_0 using volumn optimization :

$$a_0^{Si} = 5.46078 \text{ \AA}$$

(2) The band structure and density of state are shown below :




FIG. 5. Si band-structure

The band gap E_g^{Si} is :

$$E_g^{Si} = 0.6245 \text{ eV}$$

Experiment values :

$$E_g^{exp} = 1.17 \text{ eV}$$

$$a_0^{exp} = 5.431 \text{ \AA}$$

- Ge

(1) a_0 using volumn optimization :

$$a_0^{Ge} = 5.76822 \text{ \AA}$$

(2) The band structure and density of state are shown below :




FIG. 6. Ge band-structure

The band gap E_g^{Ge} is :

$$E_g^{Ge} = 0.0 \text{ eV}$$

Experiment values :

$$E_g^{exp} = 0.74 \text{ eV}$$

$$a_0^{exp} = 5.658 \text{ \AA}$$

- GaAs

(1) a_0 using volumn optimization :

$$a_0^{GaAs} = 5.751291 \text{ \AA}$$

(2) The band structure and density of state are shown below :




FIG. 7. GaAs band-structure

The band gap E_g^{GaAs} is :

$$E_g^{GaAs} = 0.1755 \text{ eV}$$

Experiment values :

$$E_g^{exp} = 1.52 \text{ eV}$$

$$a_0^{exp} = 5.65325 \text{ \AA}$$

4. GGA, GGA+U of NiO in AF-II

- GGA

- (1) the bandstructure




FIG. 8. NiO-AFII band-structure

Energy gap E_g and total magnetic moment m :

$$E_g = 0.8936 eV$$

$$m = 0.0000$$

- GGA+U

- (1) the bandstructure

Energy gap E_g and total magnetic moment m :

$$E_g = 1.0973 eV$$

$$m = 0.0000$$

[1] K. C., *Introduction to Solid State Physics, 6th Ed. (1986), p. 185. (Wiley).*




FIG. 9. NiO-AFII band-structure