

Generalized Group Lasso for Patient Subgroup Selection

Wenxuan Deng

Takeda Pharmaceuticals U.S.A., Inc.

Wenxuan.Deng@takeda.com

July 17, 2018

1 Introduction

- Prognostic and Predictive Biomarkers
- Why not regression trees?

2 Methods



3 Algorithm

- Algorithm Framework
- Computation of Proximal Operator



4 Criteria

5 Simulation

Prognostic Biomarkers



Predictive Biomarkers



Regression trees GUIDE[Loh, 2018]:

- piecewise-linear Model
- examine residual patterns for each treatment level

Cannot repeat even the most naive simulation in GUIDE paper.

Reason: limited sample size. Even two splits will results in small sample size in each branch. The results would be highly unstable. Tree-based method is not appropriate to clinical trial dataset and identify prognostic and predictive biomarkers.

Ordinary Linear Model

$$Y = X\beta + W\tau + G\alpha + W \otimes G\gamma + \epsilon$$

- X : Baseline variables
- W : Treatment variables
- G : Main effects of genes, i.e. expression levels, SNP or mutation
- $W \otimes G$: Interaction effects of genes and treatment
- ϵ : Random errors

We choose group lasso for its ability to

- handle high dimensional data
- allow hierarchical structure

However, the current group lasso based methods

- penalize on all parameters
- have no efficient adaptive penalty weights
- do not specifically target on patients treatment subgroup identification

Loss Function

We assume the hierarchical relationship between prognostic biomarkers and predictive biomarkers, that is a predictive biomarker should be a prognostic biomarker.

The loss function is

$$\min_{\theta} f(\theta|Y, X, W, G) + g(\theta)$$

where

$$g(\theta) = \lambda \sum_i \phi_i^I |\gamma_i| + \lambda \sum_i \phi_i^M \sqrt{\alpha_i^2 + \gamma_i^2}$$

where $f(\theta|Y, X, W, G)$ is L-2 loss function, i.e. sum of squared errors for ordinary linear model.

$\theta = (\beta, \tau, \alpha, \gamma)$ is parameter vector.

Loss function for ordinary linear model

$$\min_{\theta} \| Y - (X\beta + W\tau + G\alpha + W \otimes G\gamma) \|^2 + \lambda \sum_i \phi_i^l |\gamma_i| + \lambda \sum_i \phi_i^M \sqrt{\alpha_i^2 + \gamma_i^2}$$

Denote $X^{(i)} = [G_i, W \otimes G_i]$ is the i th group of the main and interaction effects of gene i . Then we let

$$\phi_i^l = \| X^{(i)} \|_2$$

$$\phi_i^M = \sqrt{\| G_i \|_2^2 + 2(1 - \sqrt{\frac{2}{\pi}}) \| W \otimes G_i \|_2^2}$$

- Fast iterative shrinkage-thresholding algorithm with backtracking[Beck and Teboulle, 2009]
- Adaptive restart for rippling behavior [O'Donoghue and Candes, 2009]
- Adaptive stepsize of cyclic Barzilai-Borwein spectral approach[Wright, 2009]
- Warm start in cross validation

Proximal Operator

Definition

Let

$$Q_{\tau_i, g}(t, u) = g(t) + \frac{1}{2\tau} \|t - u\|^2$$

then the proximal operator is defined as

$$\tilde{t} = \arg \min Q_{\tau_i, g}(t, u)$$

For convenience, we denote $P_{\tau_i, g}(u) = \tilde{t}$

Remark: Proximal operator is a point that compromises between minimizing g and being near to u .

Algorithm

initialization $\theta_0 = 0$ or warm start from previous run, $\tau_0 = 0.1$, stepsize $\eta = 0.5$;

while $i \leq k$ **do**

$u_i = \theta_{i-1} - \tau_i \nabla f(\theta_{i-1})$ Find the smallest nonnegative integers s_i such that with $\tau_i = \eta^{s_i-1} \tau_{i-1}$,

$$(f + g)(P_{\tau_i, g}(u_i)) \leq Q_{\tau_i, g}(P_{\tau_i, g}(u_i), u_i);$$

Then, we compute $t_i = P_{\tau_i, g}(u_i)$ And accelerate the computation by setting **if** $f(\theta_i + g(\theta_i)) > f(\theta_{i-1}) + g(\theta_{i-1})$ **then**

$$\quad \rho_i = 1$$

else

$$\quad \rho_i = \frac{1 + \sqrt{1 + 4\rho_{i-1}^2}}{2}$$

end

$\theta_i = t_i + (\frac{\rho_{i-1}-1}{\rho_i})(t_i - t_{i-1})$ and find τ_{i+1} that $\tau_{i+1} /$ can mimic the Hessian $\nabla^2 f(\theta_i)$

end

Algorithm 1: Patient Subgroup Identification Group Lasso Algorithm

Table

Treatments	Response 1	Response 2
Treatment 1	0.0003262	0.562
Treatment 2	0.0015681	0.910
Treatment 3	0.0009271	0.296

Table: Table caption

References



Loh, Wei-Yin, Michael Man, and Shuaicheng Wang.

"Subgroups from regression trees with adjustment for prognostic effects and postselection inference."

Statistics in medicine (2018).



Beck, Amir, and Marc Teboulle.

"A fast iterative shrinkage-thresholding algorithm for linear inverse problems."

SIAM journal on imaging sciences 2.1 (2009): 183-202. (2009).



Wright, Stephen J., Robert D. Nowak, and Mário AT Figueiredo.

"Sparse reconstruction by separable approximation."

IEEE Transactions on Signal Processing 57.7 : 2479-2493.(2009)



O'Donoghue, B., and E. Candes.

"Adaptive restart for accelerated gradient schemes."

Foundations of computational mathematics 15.3: 715-732. (2015)

The End