

References:

1. Mulholland, M. R., Morse, R., Egerton, T., Bernhardt, P. W., & Filippino, K. C. (2018). Blooms of dinoflagellate mixotrophs in a lower Chesapeake Bay tributary: Carbon and nitrogen uptake over diurnal, seasonal, and interannual timescales. *Estuaries and Coasts*, 41(6), 1744–1765. <https://doi.org/10.1007/s12237-018-0388-5>
2. Harding, L. W., Mallonee, M. E., Perry, E. S., David Miller, W., Adolf, J. E., Gallegos, C. L., & Paerl, H. W. (2020). Seasonal to inter-annual variability of primary production in Chesapeake Bay: Prospects to reverse eutrophication and change trophic classification. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-58702-3>
3. Rattner, B. A., Wazniak, C. E., Lankton, J. S., McGowan, P. C., Drovetski, S. V., & Egerton, T. A. (2022). Review of harmful algal bloom effects on birds with implications for avian wildlife in the Chesapeake Bay Region. *Harmful Algae*, 120, 102319. <https://doi.org/10.1016/j.hal.2022.102319>
4. Owens, O. V., & Esaias, W. E. (1976). Physiological responses of phytoplankton to major environmental factors. *Annual Review of Plant Physiology*, 27(1), 461–483. <https://doi.org/10.1146/annurev.pp.27.060176.002333>
5. Cotner, J. B., & Wetzel, R. G. (1992). Uptake of dissolved inorganic and organic BPHOSPHORUS compounds by phytoplankton and bacterioplankton. *Limnology and Oceanography*, 37(2), 232–243. <https://doi.org/10.4319/lo.1992.37.2.0232>
6. Marshall, Harold G., and Todd A. Egerton. "Phytoplankton blooms: Their occurrence and composition within Virginia's tidal tributaries." *Virginia Journal of Science* 60.3 (2009): 3.
7. Guildford, S. J., & Hecky, R. E. (2000). Total nitrogen, total phosphorus, and nutrient limitation in lakes and oceans: Is there a common relationship? *Limnology and Oceanography*, 45(6), 1213–1223. <https://doi.org/10.4319/lo.2000.45.6.1213>