



[Course](#) > [Unit 3: Logistic Regression](#) > [Assignment 3](#) > Predicting Parole Violators

Audit Access Expires Aug. 12, 2019

You lose all access to this course, including your progress, on Aug. 12, 2019.

Predicting Parole Violators

Predicting parole violators

In many criminal justice systems around the world, inmates deemed not to be a threat to society are released from prison under the parole system prior to completing their sentence. They are still considered to be serving their sentence while on parole, and they can be returned to prison if they violate the terms of their parole.

Parole boards are charged with identifying which inmates are good candidates for release on parole. They seek to release inmates who will not commit additional crimes after release. In this problem, we will build and validate a model that predicts if an inmate will violate the terms of his or her parole. Such a model could be useful to a parole board when deciding to approve or deny an application for parole.

For this prediction task, we will use data from the [United States 2004 National Corrections Reporting Program](#), a nationwide census of parole releases that occurred during 2004. We limited our focus to parolees who served no more than 6 months in prison and whose maximum sentence for all charges did not exceed 18 months. The dataset contains all such parolees who either successfully completed their term of parole during 2004 or those who violated the terms of their parole during that year. The dataset contains the following variables:

- **male:** 1 if the parolee is male, 0 if female
- **race:** 1 if the parolee is white, 2 otherwise
- **age:** the parolee's age (in years) when he or she was released from prison

- **state:** a code for the parolee's state. 2 is Kentucky, 3 is Louisiana, 4 is Virginia, and 1 is any other state. The three states were selected due to having a high representation in the dataset.
- **time.served:** the number of months the parolee served in prison (limited by the inclusion criteria to not exceed 6 months).
- **max.sentence:** the maximum sentence length for all charges, in months (limited by the inclusion criteria to not exceed 18 months).
- **multiple.offenses:** 1 if the parolee was incarcerated for multiple offenses, 0 otherwise.
- **crime:** a code for the parolee's main crime leading to incarceration. 2 is larceny, 3 is drug-related crime, 4 is driving-related crime, and 1 is any other crime.
- **violator:** 1 if the parolee violated the parole, and 0 if the parolee completed the parole without violation.

Problem 1.1 - Loading the Dataset

1 point possible (graded)

Load the dataset `parole.csv` into a data frame called `parole`, and investigate it using the `str()` and `summary()` functions.

How many parolees are contained in the dataset?

Answer: 675

Explanation

You can load the dataset into R with the following command:

```
parole = read.csv("parole.csv")
```

Then you can count the number of parolees in the dataset with `str(parole)` or with `nrow(parole)`.

Submit

You have used 0 of 3 attempts

i Answers are displayed within the problem

Problem 1.2 - Loading the Dataset

1 point possible (graded)

How many of the parolees in the dataset violated the terms of their parole?

Answer: 78

Explanation

This can be observed by running `table(parole$violator)`

Submit

You have used 0 of 3 attempts

i Answers are displayed within the problem

Problem 2.1 - Preparing the Dataset

1 point possible (graded)

You should be familiar with unordered factors (if not, review the Week 2 homework problem "Reading Test Scores"). Which variables in this dataset are unordered factors with at least three levels? Select all that apply.

☐ male

☐ race

☐ age

☒ state ✓

☐ time.served

☐ max.sentence

☐ multiple.offenses

☒ crime ✓

☐ violator

Explanation

While the variables male, race, state, crime, and violator are all unordered factors, only state and crime have at least 3 levels in this dataset.

Submit

You have used 0 of 2 attempts

i Answers are displayed within the problem

Problem 2.2 - Preparing the Dataset

1 point possible (graded)

In the last subproblem, we identified variables that are unordered factors with at least 3 levels, so we need to convert them to factors for our prediction problem (we introduced this idea in the "Reading Test Scores" problem last week). Using the `as.factor()` function, convert these variables to factors. Keep in mind that we are not changing the values, just the way R understands them (the values are still numbers).

How does the output of `summary()` change for a factor variable as compared to a numerical variable?

- ☒ The output becomes similar to that of the `table()` function applied to that variable ✓
- ☐ The output becomes similar to that of the `str()` function applied to that variable
- ☐ There is no change

Explanation

To convert to factors, the following commands should be run:

```
parole$state = as.factor(parole$state)
```

```
parole$crime = as.factor(parole$crime)
```

The output of `summary(parole$state)` or `summary(parole$crime)` now shows a breakdown of the number of parolees with each level of the factor, which is most similar to the output of the `table()` function.

Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem

Problem 3.1 - Splitting into a Training and Testing Set

1 point possible (graded)

To ensure consistent training/testing set splits, run the following 5 lines of code (do not include the line numbers at the beginning):

1) `set.seed(144)`

2) `library(caTools)`

3) `split = sample.split(parole$violator, SplitRatio = 0.7)`

4) `train = subset(parole, split == TRUE)`

5) `test = subset(parole, split == FALSE)`

Roughly what proportion of parolees have been allocated to the training and testing sets?

☒ 70% to the training set, 30% to the testing set ✓

☐ 50% to the training set, 50% to the testing set

☐ 30% to the training set, 70% to the testing set

Explanation

`SplitRatio=0.7` causes `split` to take the value `TRUE` roughly 70% of the time, so `train` should contain roughly 70% of the values in the dataset. You can verify this by running `nrow(train)` and `nrow(test)`.

Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem

Problem 3.2 - Splitting into a Training and Testing Set

3 points possible (graded)

Now, suppose you re-ran lines [1]-[5] of Problem 3.1. What would you expect?

- ☒ The exact same training/testing set split as the first execution of [1]-[5] ✓
- ☐ A different training/testing set split from the first execution of [1]-[5]

If you instead ONLY re-ran lines [3]-[5], what would you expect?

- ☐ The exact same training/testing set split as the first execution of [1]-[5]
- ☒ A different training/testing set split from the first execution of [1]-[5] ✓

If you instead called `set.seed()` with a different number and then re-ran lines [3]-[5] of Problem 3.1, what would you expect?

- ☐ The exact same training/testing set split as the first execution of [1]-[5]
- ☒ A different training/testing set split from the first execution of [1]-[5] ✓

Explanation

If you set a random seed, split, set the seed again to the same value, and then split again, you will get the same split. However, if you set the seed and then split twice, you will get different splits. If you set the seed to different values, you will get different splits.

You can also verify this by running the specified code in R. If you have training sets `train1` and `train2`, the function `sum(train1 != train2)` will count the number of values in those two data frames that are different.

Submit

You have used 0 of 1 attempt

Problem 4.1 - Building a Logistic Regression Model

1 point possible (graded)

If you tested other training/testing set splits in the previous section, please re-run the original 5 lines of code to obtain the original split.

Using `glm` (and remembering the parameter `family="binomial"`), train a logistic regression model on the training set. Your dependent variable is "violator", and you should use all of the other variables as independent variables.

What variables are significant in this model? Significant variables should have a least one star, or should have a probability less than 0.05 (the column `Pr(> | z |)` in the summary output). Select all that apply.

☐ male☒ race ✓☐ age☐ state2☐ state3☒ state4 ✓☐ time.served☐ max.sentence☒ multiple.offenses ✓☐ crime2☐ crime3☐ crime4

Explanation

The following lines of code show the summary of the logistic regression model:
`mod = glm(violator~., data=train, family="binomial")`
`summary(mod)`

Submit

You have used 0 of 3 attempts

i Answers are displayed within the problem

Problem 4.2 - Building a Logistic Regression Model

1 point possible (graded)

What can we say based on the coefficient of the multiple.offenses variable?

The following two properties might be useful to you when answering this question:

1) If we have a coefficient c for a variable, then that means the log odds (or Logit) are increased by c for a unit increase in the variable.

2) If we have a coefficient c for a variable, then that means the odds are multiplied by e^c for a unit increase in the variable.

- ☐ Our model predicts that parolees who committed multiple offenses have 1.61 times higher odds of being a violator than the average parolee.
- ☐ Our model predicts that a parolee who committed multiple offenses has 1.61 times higher odds of being a violator than a parolee who did not commit multiple offenses but is otherwise identical.
- ☐ Our model predicts that parolees who committed multiple offenses have 5.01 times higher odds of being a violator than the average parolee.
- ☒ Our model predicts that a parolee who committed multiple offenses has 5.01 times higher odds of being a violator than a parolee who did not commit multiple offenses but is otherwise identical. ✓

Explanation

For parolees A and B who are identical other than A having committed multiple offenses, the predicted log odds of A is 1.61 more than the predicted log odds of B. Then we have:

$$\ln(\text{odds of A}) = \ln(\text{odds of B}) + 1.61$$

$$\exp(\ln(\text{odds of A})) = \exp(\ln(\text{odds of B}) + 1.61)$$

$$\exp(\ln(\text{odds of A})) = \exp(\ln(\text{odds of B})) * \exp(1.61)$$

$$\text{odds of A} = \exp(1.61) * \text{odds of B}$$

$$\text{odds of A} = 5.01 * \text{odds of B}$$

In the second step we raised e to the power of both sides. In the third step we used the exponentiation rule that $e^{(a+b)} = e^a * e^b$. In the fourth step we used the rule that $e^{(\ln(x))} = x$.

Submit

You have used 0 of 2 attempts

i Answers are displayed within the problem

Problem 4.3 - Building a Logistic Regression Model

0.0/4.0 points (graded)

Consider a parolee who is male, of white race, aged 50 years at prison release, from the state of Maryland, served 3 months, had a maximum sentence of 12 months, did not commit multiple offenses, and committed a larceny. Answer the following questions based on the model's predictions for this individual. (HINT: You should use the coefficients of your model, the Logistic Response Function, and the Odds equation to solve this problem.)

According to the model, what are the odds this individual is a violator?

Answer: 0.1825687

According to the model, what is the probability this individual is a violator?

Answer: 0.1543831

Explanation

From the logistic regression equation, we have $\log(\text{odds}) = -4.2411574 + 0.3869904 \cdot \text{male} + 0.8867192 \cdot \text{race} - 0.0001756 \cdot \text{age} + 0.4433007 \cdot \text{state2} + 0.8349797 \cdot \text{state3} - 3.3967878 \cdot \text{state4} - 0.1238867 \cdot \text{time.served} + 0.0802954 \cdot \text{max.sentence} + 1.6119919 \cdot \text{multiple.offenses} + 0.6837143 \cdot \text{crime2} - 0.2781054 \cdot \text{crime3} - 0.0117627 \cdot \text{crime4}$. This parolee has $\text{male}=1$, $\text{race}=1$, $\text{age}=50$, $\text{state2}=0$, $\text{state3}=0$, $\text{state4}=0$, $\text{time.served}=3$, $\text{max.sentence}=12$, $\text{multiple.offenses}=0$, $\text{crime2}=1$, $\text{crime3}=0$, $\text{crime4}=0$. We conclude that $\log(\text{odds}) = -1.700629$.

Therefore, the odds ratio is $\exp(-1.700629) = 0.183$, and the predicted probability of violation is $1/(1+\exp(1.700629)) = 0.154$.

Submit

You have used 0 of 7 attempts

i Answers are displayed within the problem

Problem 5.1 - Evaluating the Model on the Testing Set

1 point possible (graded)

Use the `predict()` function to obtain the model's predicted probabilities for parolees in the testing set, remembering to pass `type="response"`.

What is the maximum predicted probability of a violation?

Answer: 0.907

Explanation

The following commands make the predictions and display a summary of the values:

```
predictions = predict(mod, newdata=test, type="response")
summary(predictions)
```

Submit

You have used 0 of 5 attempts

i Answers are displayed within the problem

Problem 5.2 - Evaluating the Model on the Testing Set

3 points possible (graded)

In the following questions, evaluate the model's predictions on the test set using a threshold of 0.5.

What is the model's sensitivity?

Answer: 0.522

What is the model's specificity?

Answer: 0.933

What is the model's accuracy?

Answer: 0.886

Explanation

To obtain the confusion matrix, use the following command:

```
table(test$violation, as.numeric(predictions >= 0.5))
```

There are 202 observations in the test set. The accuracy (percentage of values on the diagonal) is $(167+12)/202 = 0.886$. The sensitivity (proportion of the actual

violators we got correct) is $12/(11+12) = 0.522$, and the specificity (proportion of the actual non-violators we got correct) is $167/(167+12) = 0.933$.

Submit

You have used 0 of 5 attempts

i Answers are displayed within the problem

Problem 5.3 - Evaluating the Model on the Testing Set

1 point possible (graded)

What is the accuracy of a simple model that predicts that every parolee is a non-violator?

Answer: 0.886

Explanation

If you table the outcome variable using the following command:

```
table(test$violator)
```

you can see that there are 179 negative examples, which are the ones that the baseline model would get correct. Thus the baseline model would have an accuracy of $179/202 = 0.886$.

Submit

You have used 0 of 5 attempts

i Answers are displayed within the problem

Problem 5.4 - Evaluating the Model on the Testing Set

1 point possible (graded)

Consider a parole board using the model to predict whether parolees will be violators or not. The job of a parole board is to make sure that a prisoner is ready to be released into free society, and therefore parole boards tend to be particularly concerned about releasing prisoners who will violate their parole. Which of the following most likely describes their preferences and best course of action?

- ☐ The board assigns more cost to a false negative than a false positive, and should therefore use a logistic regression cutoff higher than 0.5.
- ☒ The board assigns more cost to a false negative than a false positive, and should therefore use a logistic regression cutoff less than 0.5. ✓
- ☐ The board assigns equal cost to a false positive and a false negative, and should therefore use a logistic regression cutoff equal to 0.5.
- ☐ The board assigns more cost to a false positive than a false negative, and should therefore use a logistic regression cutoff higher than 0.5.
- ☐ The board assigns more cost to a false positive than a false negative, and should therefore use a logistic regression cutoff less than 0.5.

Explanation

If the board used the model for parole decisions, a negative prediction would lead to a prisoner being granted parole, while a positive prediction would lead to a prisoner being denied parole. The parole board would experience more regret for releasing a prisoner who then violates parole (a negative prediction that is actually positive, or false negative) than it would experience for denying parole to a prisoner who would not have violated parole (a positive prediction that is actually negative, or false positive).

Decreasing the cutoff leads to more positive predictions, which increases false positives and decreases false negatives. Meanwhile, increasing the cutoff leads to more negative predictions, which increases false negatives and decreases false positives. The parole board assigns high cost to false negatives, and therefore should decrease the cutoff.

Submit

You have used 0 of 2 attempts

i Answers are displayed within the problem

Problem 5.5 - Evaluating the Model on the Testing Set

1 point possible (graded)

Which of the following is the most accurate assessment of the value of the logistic

regression model with a cutoff 0.5 to a parole board, based on the model's accuracy as compared to the simple baseline model?

- ☐ The model is of limited value to the board because it cannot outperform a simple baseline, and using a different logistic regression cutoff is unlikely to improve the model's value.
- ☐ The model is of limited value to the board because it cannot outperform a simple baseline, and using a different logistic regression cutoff is likely to improve the model's value.
- ☐ The model is likely of value to the board, and using a different logistic regression cutoff is unlikely to improve the model's value.
- ☒ The model is likely of value to the board, and using a different logistic regression cutoff is likely to improve the model's value. ✓

Explanation

The model at cutoff 0.5 has 12 false positives and 11 false negatives, while the baseline model has 0 false positives and 23 false negatives. Because a parole board is likely to assign more cost to a false negative, the model at cutoff 0.5 is likely of value to the board.

From the previous question, the parole board would likely benefit from decreasing the logistic regression cutoffs, which decreases the false negative rate while increasing the false positive rate.

Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem

Problem 5.6 - Evaluating the Model on the Testing Set

0.0/2.0 points (graded)

Using the ROCR package, what is the AUC value for the model?

Answer: 0.8945834

Explanation

This can be obtained with the following code:

```
library(ROCR)
pred = prediction(predictions, test$violator)
as.numeric(performance(pred, "auc")@y.values)
```

Submit

You have used 0 of 5 attempts

i Answers are displayed within the problem

Problem 5.7 - Evaluating the Model on the Testing Set

1 point possible (graded)

Describe the meaning of AUC in this context.

- ☒ The probability the model can correctly differentiate between a randomly selected parole violator and a randomly selected parole non-violator. ✓
- ☐ The model's accuracy at logistic regression cutoff 0.5.
- ☐ The model's accuracy at the logistic regression cutoff at which it is most accurate.

Explanation

The AUC deals with differentiating between a randomly selected positive and negative example. It is independent of the regression cutoff selected.

Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem

Problem 6.1 - Identifying Bias in Observational Data

1 point possible (graded)

Our goal has been to predict the outcome of a parole decision, and we used a publicly available dataset of parole releases for predictions. In this final problem,

we'll evaluate a potential source of bias associated with our analysis. It is always important to evaluate a dataset for possible sources of bias.

The dataset contains all individuals released from parole in 2004, either due to completing their parole term or violating the terms of their parole. However, it does not contain parolees who neither violated their parole nor completed their term in 2004, causing non-violators to be underrepresented. This is called "selection bias" or "selecting on the dependent variable," because only a subset of all relevant parolees were included in our analysis, based on our dependent variable in this analysis (parole violation). How could we improve our dataset to best address selection bias?

- ☐ There is no way to address this form of biasing.
- ☐ We should use the current dataset, expanded to include the missing parolees. Each added parolee should be labeled with violator=0, because they have not yet had a violation.
- ☐ We should use the current dataset, expanded to include the missing parolees. Each added parolee should be labeled with violator=NA, because the true outcome has not been observed for these individuals.
- ☒ We should use a dataset tracking a group of parolees from the start of their parole until either they violated parole or they completed their term.



Explanation

While expanding the dataset to include the missing parolees and labeling each as violator=0 would improve the representation of non-violators, it does not capture the true outcome, since the parolee might become a violator after 2004. Though labeling these new examples with violator=NA correctly identifies that we don't know their true outcome, we cannot train or test a prediction model with a missing dependent variable.

As a result, a prospective dataset that tracks a cohort of parolees and observes the true outcome of each is more desirable. Unfortunately, such datasets are often more challenging to obtain (for instance, if a parolee had a 10-year term, it might require tracking that individual for 10 years before building the model). Such a prospective analysis would not be possible using the 2004 National Corrections Reporting Program dataset.

Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem

Please remember not to ask for or post complete answers to homework questions in this discussion forum.

Discussion

Show Discussion

Topic: Unit 3 / Unit 3, Homework: Predicting Parole Violators

© All Rights Reserved