Week 3 Cheat Sheet


Statistics and Data Analysis with Excel, Part 1

Charlie Nuttelman


Here, I provide the mathematical equations and some of the important Excel functions required to perform various calculations in Week 3 of the course. The headings represent the screencasts in which you will find those calculations and concepts. Not all screencasts are referenced below – just the ones that have complex mathematical formulas or Excel formulas that are tricky to use.

Sample Spaces and Events


Union of two events:



Intersection of two events:



Complement of an event:



Mutually exclusive events: Two events are mutually exclusive if their intersection is the null set (zero) $(E_1 \cap E_2 = \emptyset)$.

Permutations and Combinations

Permutation: The number of ways we can put r items into n spots if order matters.

Denoted by: ${}_{n}P_{r}$

Calculation: ${}_{n}P_{r} = \frac{n!}{(n-r)!}$ (! denotes the factorial)

Easy to calculate in Excel: PERMUT(n,r)

Combination: The number of ways we can put r items into n spots if order does not matter.

Denoted by: ${}_{n}C_{r}$

Calculation: ${}_{n}C_{r} = \frac{n!}{(n-r)!r!}$ (! denotes the factorial)

Alternative notation: $\binom{n}{r}$ (This is especially important in some of the functions in Week 4)

In Excel: COMBIN(n,r)

Permutations of like objects: When members of subgroups are indistinguishable, the number of ways that we can arrange them if order matters is:

permutations of like objects =
$$\frac{n!}{n_1! \cdot n_1! \cdot \dots \cdot n_k!}$$


Here, the items are separated into k distinct groups, and n_1 , n_2 , ..., n_k represent the number of items in each of those distinct groups.

Probability Rules


Probability can be thought of as the ratio of the number of ways of satisfying a particular constraint to the total number of possibilities:

$$P = \frac{Total\ ways\ of\ satisfying\ a\ constraint}{Total\ possibilities}$$

Probability of a union:



Mutually exclusive events (no intersection):




Probability of a complement:


$$P[A']=1-P[A]$$
 where A' is complement of A
$$S \longrightarrow A'$$

$$P[S]=1$$

Conditional Probability


The conditional probability of B given A is defined by:





Multiplication Rule

Starting with rearranged versions of conditional probabilities (see above), we can set the left-hand side of the following two equations equal to each other:



$$P[A \cap B] = P[B|A]P[A]$$


$$P[A \cap B] = P[A|B]P[B]$$

to arrive at the Multiplication Rule:

$$P[B|A] \cdot P[A] = P[A|B] \cdot P[B]$$

Total Probability Rule

Consider two events A and A' that are exhaustive (meaning that they take up the entire sample set) and each has an intersection with event B:




The total probability of B is given by

$$P[B] = P[B \cap A] + P[B \cap A']$$

Using the rules of conditional probability (see above), we arrive at the Total Probability Rule:

$$P[B] = P[B|A] \cdot P[A] + P[B|A'] \cdot P[A']$$


This can be extended to cases with multiple exhaustive events (E_1 through E_5 , in the following example), where each of those events has an intersection with event B:



$$P[B] = P[B|E_1] \cdot P[E_1] + P[B|E_2] \cdot P[E_2] + P[B|E_3] \cdot P[E_3] + P[B|E_4] \cdot P[E_4] + P[B|E_5] \cdot P[E_5]$$

Bayes' Theorem


Again consider two events A and A' that are exhaustive (meaning that they take up the entire sample set) and each has an intersection with event B:



We can rearrange the Multiplication Rule and substitute in the Total Probability Rule to arrive at Bayes' Theorem:

$$P[A|B] = \frac{P[B|A]P[A]}{P[B|A]P[A] + P[B|A']P[A']}$$

Like the Total Probability Rule, this can be extended to cases with multiple exhaustive events (E_1 through E_5 , in the following example), where each of those events has an intersection with event B:



$$P[E_{j}|B] = \frac{P[B|E_{j}] \cdot P[E_{j}]}{\sum_{i=1}^{k} P[B|E_{i}] P[E_{i}]}$$