## **Thermodynamics**



Summary of material in FIRST YEAR lectures (skeleton notes)

You must put 'flesh' on these 'bare bones'

(...not everything is covered)

CHEM1011: Thermodynamics © Steve Colbran Slide 83

# **Copyright Notice**

Some images used in these lectures are taken, with permission, from "Chemistry", Blackman et al. (Wiley, Brisbane, 2008 & 2012), and other sources as indicated, in accordance with the Australian copyright regulations.

#### COMMONWEALTH OF AUSTRALIA Copyright Regulation Warning

This material has been reproduced and communicated to you by or on behalf of the University of New South Wales pursuant to Part VA of the Copyright Act 1968 (the Act)

The material in this communication may be subject to copyright under the Act.

Any further reproduction or communication of this material by you may be subject to copyright protection under the act.

Do not remove this notice.

All of these lecture notes are made available to UNSW students through Moodle.

These lecture notes may not be posted on any other website.

## — Thermochemistry: Bare Bones —

### Heat (energy transferred due to $\Delta T$ )

$$q = n \times C_p \times \Delta T$$

where n = amount,  $C_p$  = heat capacity,  $\Delta T$  = temperature change

and for a calorimeter

$$q = q_{contents} + q_{calorimeter}$$

$$= (n \times C_p \times \Delta T) + (C_{calorimeter} \times \Delta T)$$

Heat out: q < 0; Heat in: q > 0 (by definition)

#### **Work (at constant pressure)**

$$w = -p\Delta V$$
$$= -\Delta n_{gas} RT$$

(negative for expansion)
(assuming Ideal Gas Equation)

#### **First Law of Thermodynamics**


$$\Delta U = q + w$$

(state function → path independent)

#### **Enthalpy (Heat of Reaction at constant P)**



Hess's Law: enthalpy is a state function, so path independent



Be able to use tabulated *standard\** enthalpies to calculate  $\Delta_r H^\circ$ :

- Enthalpy (heat) of formation,  $\Delta_f H^\circ$
- Enthalpy (heat) of combustion,  $\Delta_c H^\circ$
- Bond Enthalpy (energy), Δ<sub>X-Y</sub>H°
- Enthalpy of melting or vaporisation, Heat capacity etc.

\*Standard states: P = 1.0 bar, pure (unmixed) substances or 1.0 mol L-1 solutions at the specified temperature

#### **Second Law of Thermodynamics**

 $\Delta S$  (universe) > 0 for a spontaneous process

where **S = entropy, increases as disorder increases** 

#### **Third Law of Thermodynamics**

S (pure crystal)  $\rightarrow$  constant value as  $T \rightarrow 0$  K Define S° (pure crystal) = 0 at T = 0 K

so can calc. So at any T

 $S^{\circ} = Sum \ m_i \ \Delta S^{\circ}_{products \ i} - Sum \ m_j \ \Delta S^{\circ}_{reactants \ j}$ 

#### **Gibbs Free Energy**

$$\Delta_r G = \Delta_r H - T \Delta_r S$$

$$= Sum \ m_i \ \Delta G_{products \ i} - Sum \ m_j \ \Delta G_{reactants \ j}$$

 $\Delta G = -T \Delta S$  (universe)


*so...*  $\Delta G < 0$  implies process (reaction) is **spontaneous**  $\Delta G > 0$  implies process (reaction) is **non-spontaneous** 

#### At equilibrium:

$$\Delta_r G = 0$$
  
and  $\Delta_r G^\circ = \Delta_r H^\circ - T \Delta_r S^\circ = -RT \ln K \ (\neq 0)$ 

... and a little extra (not examinable, but it may help)

( to clear up confusion between  $\Delta G$  and  $\Delta G^{\circ}$  -- they are different!! )



# How should I study for this topic?

