



# Pymble Ladies' College

## Physics

2001

Trial Examination

### General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Board-approved calculators may be used
- Write using black or blue pen
- Draw diagrams using pencil

### Section I

Total marks (75)

This section has two parts, Part A and Part B

#### Part A Multiple choice Total marks (15)

- Attempt Questions 1–15
- Allow about 30 minutes for this part

#### Part B Extended Answers Total marks (60)

- Attempt Questions 16–30
- Allow about 1 hour and 45 minutes for this part

### Section II

Total marks (25)

- Attempt ONE question - Question 31
- Allow about 45 minutes for this section

Physics  
2001  
Trial Examination

Multiple Choice Answer Sheet

Select the alternative A, B, C or D that best answers the question.  
Fill in the response space completely. If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.  
If you change your mind and have crossed out what you consider to be the correct answer, then indicate this by writing the word **correct** and drawing an arrow.

Question	A	B	C	D
1	A	B	C	D
2	A	B	C	D
3	A	B	C	D
4	A	B	C	D
5	A	B	C	D
6	A	B	C	D
7	A	B	C	D
8	A	B	C	D
9	A	B	C	D
10	A	B	C	D
11	A	B	C	D
12	A	B	C	D
13	A	B	C	D
14	A	B	C	D
15	A	B	C	D

## Section 1

Total marks (75)

This section has two parts, Part A and Part B

### Part A

Multiple choice Total marks (15)

- Attempt Questions 1–15
- Allow about 30 minutes for this part

#### Question 1

Jill has a weight of 550 N on the earth. What is her weight on a planet with half the mass of earth and half the radius of earth?

- A 69 N
- B 275 N
- C 550 N
- D 1100 N

#### Question 2

Which of the following factors does not affect the escape velocity of an object from earth?

- A the mass of the object
- B the mass of the earth
- C the radius of the earth
- D the gravitational constant  $G$

#### Question 3

A satellite in orbit at a distance  $R$  from the centre of the earth has a period of 12 hours. What is the period of a satellite orbiting at a distance  $3R$ ?

- A 4 hours
- B 21 hours
- C 36 hours
- D 62 hours


#### Question 4

Which of the following is an inertial frame of reference?

- A a rocket just after takeoff
- B a deep space probe without fuel
- C a satellite in geostationary orbit around the earth
- D a sub-orbital rocket at the point of maximum height in its trajectory

#### Question 5

Trainee astronauts could have the experience of 'weightlessness' by flying in a plane that is travelling in vertical, circular path, as shown in the diagram below.



What is the radius  $R$  of the vertical circle if the plane is flying at a constant speed of  $20 \text{ m.s}^{-1}$  and the astronauts feel 'weightless' at the top of the circle?

- A 20 m
- B 40 m
- C 80 m
- D 160 m


#### Question 6

Who was the scientist who discovered that an electric current could be induced by moving a magnet near a coil of wire?

- A Ampere
- B Lenz
- C Faraday
- D Tesla

### Question 7

The diagram below shows a current carrying wire in a magnetic field.




In which direction will the wire tend to move?

- A up
- B down
- C into the page
- D out of the page

### Question 8

Two straight current-carrying conductors are placed parallel to each other, 4 cm apart. One has a current of 2 A travelling through it and the other has a current of 5 A travelling through it. Both currents travel in the same direction.




What is the force on 1 m of the 5 A wire due to the 2 A wire?

- A  $5 \times 10^{-5}$  N towards the 2 A wire.
- B  $5 \times 10^{-5}$  N away from the 2 A wire.
- C  $5 \times 10^{-7}$  N towards the 2 A wire.
- D  $5 \times 10^{-7}$  N away from the 2 A wire.

### Question 9

The square loop shown in the diagram below has sides 50 mm x 50 mm and is supported on a central axle, parallel to the sides AB and CD. It carries a current of 5 A and is in a uniform magnetic field of  $2.0 \times 10^{-2}$  T.



What is the torque experienced by the loop when the plane of the loop is lying parallel to the magnetic field as shown?

- A 0 Nm
- B  $2.5 \times 10^{-4}$  Nm
- C  $5.0 \times 10^{-3}$  Nm
- D 2.5 Nm


### Question 10

Which of the following methods is used to reduce energy losses in electrical transmission wires?

- A using good insulation
- B keeping voltage as low as possible
- C keeping current as low as possible
- D keeping resistance as high as possible

### Question 11


What was the equipment below used for?



- A To demonstrate the photoelectric effect
- B Hertz' experiment with electromagnetic waves
- C The first radio
- D To demonstrate thermionic conduction

### Question 12

The diagram below shows two charged, parallel plates.




An electron is fired into the space between the two plates in the direction shown. The electron will travel through without being deflected if a magnetic field is also present between the plates. What would the direction of the magnetic field have to be?

- A into the page
- B out of the page
- C towards the positive plate
- D towards the negative plate


### Question 13

The resistance ( $R$ ) of a superconductor is plotted as a function of temperature ( $T$ ). Which graph would most closely represent the results obtained?



### Question 14

Two charged particles, A and B, are fired into a uniform magnetic field as shown below.



The initial velocity of particle A is twice that of particle B. Particle A has a charge of  $-0.5Q$  coulombs. Particle B has a charge of  $+Q$  coulombs.  $F_A$  is the force acting on particle A due to the magnetic field.  $F_B$  is the force acting on particle B due to the magnetic field.


Which of the following statements is true?

- A  $F_A$  is the same size as  $F_B$ .
- B  $F_A$  is twice the size of  $F_B$ .
- C  $F_A$  is half the size of  $F_B$ .
- D  $F_A$  is a quarter the size of  $F_B$ .

### Question 15

The diagram below shows one of the cathode ray tubes that can be used to demonstrate the properties of cathode rays. Which of the following can be deduced from the effect observed from this particular cathode ray tube?

rotating wheel



- A Cathode rays are negatively charged.
- B Cathode rays are fast moving electrons.
- C Cathode rays have energy and momentum.
- D Cathode rays are electromagnetic.

### Part B

Extended Answers

Total marks (60)

- Attempt Questions 16–30
- Allow about 1 hour and 45 minutes for this part

### Question 16: (3 marks)

Describe difficulties associated with effective and reliable communications between satellites and earth.

Marks

3

.....

.....

.....

.....

.....

.....

.....

.....

.....

### Question 17: (4 marks)

Marks

A rocket is fired from its launch pad with an initial speed of  $80 \text{ m.s}^{-1}$  at an angle of  $35^\circ$  to the horizontal.

Calculate:

- (a) its total time of flight.

3

.....

.....

.....

.....

.....

.....

.....

.....

.....

Continued on next page ...

(b) its range.

.....

.....

.....

.....

.....

Question 19: (5 marks)

Describe a first-hand investigation to determine a value for the acceleration due to gravity using pendulum motion.

The relevant equation is  $g = 4\pi^2 \ell / T^2$

where  $g$  is the acceleration due to gravity

$\ell$  is the length of the pendulum

$T$  is the period of oscillation of the pendulum

Question 18: (4 marks)

A rocket is travelling to the star closest to earth, Proxima Centauri, which is a distance of 4.3 light years away. The rocket travels at a speed of  $0.7c$  and the time taken to accelerate and decelerate is negligible.

(a) Calculate the number of years that will pass, as measured by the crew of the rocket, as they travel to Proxima Centauri.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Calculate the distance to Proxima Centauri, as measured by the crew, in light years.

.....

.....

.....

.....

.....

Question 20: (4 marks)

Marks


Explain how space probes may use planets to provide a slingshot effect.

4

Question 21: (5 marks)

Marks

The diagram below shows a generator.



(a) Explain how the generator works.

4

.....

.....

.....

.....

.....

.....

.....

(b) Describe how this generator could be transformed into a DC generator.

1

.....

.....


.....




**Question 22:** (3 marks)

**Marks**

Below is a diagram of a square coil of wire attached to a split-ring commutator and a power source that provided a current of 2 A. The coil had 250 turn and sides of 4 cm x 4 cm.



A student placed some permanent magnets at A and B and the motor started spinning. Attaching a torque meter to the axle, the student was able to determine the torque at various angles  $\theta$  (theta). The student then plotted a graph of torque (Nm) against cosine  $\theta$ , as shown below.



Use the graph and the information given to calculate the strength of the magnetic field provided by the magnets. Show all working.

**3**

**Question 23:** (3 marks)

**Marks**


Explain the advantages of induction motors compared with conventional A.C. motors.

**3**

Continued on next page ...

**Question 24: (4 marks)**

Two magnets are brought near to a spinning aluminium disc, as shown in the diagram below.



- (a) Explain what happens when the magnets are brought near.

- (b) Explain how this effect could be reduced.

**Question 25: (5 marks)**

A transformer has 300 turns in the primary coil and 10 turns in the secondary coil. The primary voltage is 240 V AC and the primary current is 2 A.

- (a) Calculate the secondary voltage in the transformer.

- (b) Explain why an experimentally observed value might be different to your answer to part (a)?

- (c) Explain why some electrical appliances in the home that are connected to the mains domestic power supply use a transformer.

- (b) Explain how this effect could be reduced.

**Marks**

- (a) Explain what happens when the magnets are brought near.

- (b) Explain how this effect could be reduced.

## 4

4

4

## 4

५

५

b) Explain the photoelectric effect using Einstein's model for light.

5

5



## Section II

Total marks (25)

Allow about 45 minutes for this section.

Answer Question 31 on the writing paper provided.

Extra writing paper is available.

### Question 31 – From Quanta to Quarks (25 marks)

Marks

a) Carbon-13 is one isotope of the element carbon. With reference to Carbon explain the term "isotope". 1

b) i) By considering the various forces within the nucleus explain why there must be a strong nuclear force. 2


ii) State one property of the strong nuclear force. 1

c) i) Compare and contrast a controlled and uncontrolled nuclear chain reaction 3

ii) Explain how a controlled nuclear chain reaction is maintained in a nuclear reactor. 3

d) Write an equation for the nuclear reaction that occurs when Plutonium-241 undergoes  $\alpha$  decay. 2

e) A typical fission reaction is




Calculate the amount of energy released in this reaction.

Data:	${}^1_0\text{n}$	1.008665 u	${}^{141}_{56}\text{Ba}$	140.9141 u
	${}^{235}_{92}\text{U}$	235.043925 u	${}^{92}_{36}\text{Kr}$	91.9250 u

### Question 31 continued.


f) The graph below shows the relative number of beta particles emitted by a radioactive source as a function of the beta particle's kinetic energy.



i) Explain the difficulty in understanding this pattern of energy distribution when it was first observed. 3

ii) Describe how this difficulty was overcome. 1

g) An experiment was done in which an aluminium barrier was placed between a radioactive source and a detector. The radioactive source emitted  $\alpha$  particles and the number of counts during a 10 second time interval was recorded. The diagram below shows the experimental arrangement. When the radioactive source was removed, the detector registered 4 counts in the 10 second interval.




In the experiment a number of different thicknesses of aluminium were used.

The graph of the experimental results is shown below.

Question 31 continued on next page

# Penetration of Beta Particles as a Function of Barrier Thickness



Analyse the experimental results.

6

## Numerical values of several constants

Charge on the electron, $q_e$	$-1.602 \times 10^{-19} \text{ C}$
Mass of electron, $m_e$	$9.109 \times 10^{-31} \text{ kg}$
Mass of neutron, $m_n$	$1.675 \times 10^{-27} \text{ kg}$
Mass of proton, $m_p$	$1.673 \times 10^{-27} \text{ kg}$
Speed of sound in air	$340 \text{ m s}^{-1}$
Earth's gravitational acceleration, $g$	$9.8 \text{ m s}^{-2}$
Speed of light (in vacuo), $c$	$3.00 \times 10^8 \text{ m s}^{-1}$
Magnetic force constant, $\left(k \equiv \frac{\mu_0}{2\pi}\right)$	$2.0 \times 10^{-7} \text{ N A}^{-2}$
Universal gravitational constant, $G$	$6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
Mass of Earth	$6.0 \times 10^{24} \text{ kg}$
Planck's constant, $h$	$6.626 \times 10^{-34} \text{ J s}$
Rydberg's constant, $R_H$	$1.097 \times 10^7 \text{ m}^{-1}$
Atomic mass unit, $u$	$1.661 \times 10^{-27} \text{ kg}$ $931.5 \text{ MeV}/c^2$
1 eV	$1.602 \times 10^{-19} \text{ J}$
Density of water, $\rho$	$1.00 \times 10^3 \text{ kg m}^{-3}$
Specific heat capacity of water	$4.18 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

$$c = f\lambda$$

$$\text{Intensity} \propto \frac{1}{d^2}$$

$$\frac{v_1}{v_2} = \frac{\sin i}{\sin r}$$

$$E = \frac{F}{q}$$

$$R = \frac{V}{I}$$

$$P = VI$$

$$\text{Energy} = VIt$$

$$v_{av} = \frac{\Delta x}{\Delta t}$$

$$a_{av} = \frac{\Delta v}{\Delta t} = \frac{v-u}{t}$$

$$\Sigma F = ma$$

$$E_k = \frac{1}{2}mv^2$$

$$p = mv$$

$$\Delta p = Ft$$

$$F = \frac{Gm_1m_2}{r^2}$$

$$\frac{r^3}{T^2} = \frac{GM}{4\pi^2}$$

$$m_1 + m_2 = \frac{4\pi^2 r^3}{GT^2}$$

$$M = m - 5 \log \left( \frac{d}{10} \right)$$

$$\frac{I_A}{I_B} = 100(m_B - m_A)/5$$

$$d = \frac{1}{p}$$

$$F = BIl \sin \theta$$

$$E = k \frac{1}{l} \frac{I_1 I_2}{d}$$

$$\tau = Fd$$

$$\tau = nBI A \cos \theta$$

$$\frac{V_p}{V_s} = \frac{n_p}{n_s}$$

$$E_p = \frac{Gm_1m_2}{r}$$

$$v = u + at$$

$$v_x^2 = u_x^2$$

$$v_y^2 = u_y^2 + 2a_y \Delta y$$

$$\Delta x = u_x t$$

$$\Delta y = u_y t + \frac{1}{2} a_y t^2$$

$$\frac{s}{t} = \frac{u+v}{2}$$

$$l_v = l_o \sqrt{1 - \frac{v^2}{c^2}}$$

$$l_v = \frac{l_o}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$F = qvB \sin \theta$$

$$E = \frac{V}{d}$$

$$E = hf$$

$$Z = \rho v$$

$$\frac{l_r}{l_o} = \frac{[Z_2 - Z_1]^2}{[Z_2 + Z_1]^2}$$

$$\frac{1}{\lambda} = R_H \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

$$\lambda = \frac{h}{mv}$$

$$\text{Amplifier gain} = \frac{V_{out}}{V_{in}}$$

$$A_o = \frac{V_o}{V_+ - V_-}$$

PERIODIC TABLE OF THE ELEMENTS

1 H 1.008	2 He 4.003	3 Li 6.941	4 Be 9.012	5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18	11 Na 22.99	12 Mg 24.31	13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.61	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc [98.91]	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3	55 Cs 132.9	56 Ba 137.3	57-71 Lanthanides	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po [210.0]	85 At [222.0]	86 Rn [222.0]	87 Fr [223.0]	88 Ra [226.0]	89-103 Actinides	104 Rf [261.1]	105 Db [262.1]	106 Sg [263.1]	107 Bh [264.1]	108 Hs [265.1]	109 Mt [268]	110 Un [270]	111 Uub [271]	112 Uut [272]	113 Uuq [273]	114 Uuq [274]	115 Uub [275]	116 Uuh [276]	117 Uus [277]	118 Uuo [278]	119 Uu [279]	120 Uu [280]
-----------------	------------------	------------------	------------------	-----------------	-----------------	-----------------	-----------------	-----------------	-------------------	-------------------	-------------------	-------------------	-------------------	------------------	------------------	-------------------	-------------------	------------------	-------------------	-------------------	-------------------	------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	------------------	-------------------	-------------------	-------------------	---------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	------------------	-------------------	-------------------	-------------------	----------------------	-------------------	-------------------	------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	---------------------	---------------------	---------------------	---------------------	---------------------	---------------------	----------------------	----------------------	----------------------	----------------------	----------------------	--------------------	--------------------	---------------------	---------------------	---------------------	---------------------	---------------------	---------------------	---------------------	---------------------	--------------------	--------------------

KEY  
Atomic Number  
Atomic Weight  
Symbol of element  
Name of element  
Cold

57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm [146.9]	62 Sm 150.4	63 Eu 152.0	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0	Actinides	89 Ac [227.0]	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np [237.0]	94 Pu [239.1]	95 Am [241.1]	96 Cm [244.1]	97 Bk [249.1]	98 Cf [252.1]	99 Es [252.1]	100 Fm [257.1]	101 Md [258.1]	102 No [259.1]	103 Lr [262.1]	Lanthanides
-------------------	-------------------	-------------------	-------------------	---------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-----------	---------------------	-------------------	-------------------	------------------	---------------------	---------------------	---------------------	---------------------	---------------------	---------------------	---------------------	----------------------	----------------------	----------------------	----------------------	-------------

Where the atomic weight is not known, the relative atomic mass of the most common radioactive isotope is shown in brackets.  
The atomic weights of Np and Tc are given for the isotopes <sup>237</sup>Np and <sup>99</sup>Tc.

This sheet should be REMOVED for your convenience.