



**CATHOLIC SECONDARY SCHOOLS ASSOCIATION OF NSW
2013 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION
MATHEMATICS EXTENSION 1**

**Section I
10 marks**

Questions 1-10 (1 mark each)

Question 1 (1 mark)

Outcomes Assessed: HE6

Targeted Performance Bands: E2

Solution	Answer	Mark
$\int 2 \sin^2 x \, dx = \int (1 - \cos 2x) \, dx$ $= x - \frac{1}{2} \sin 2x + C$	B	1

Question 2 (1 mark)

Outcomes Assessed: PE3

Targeted Performance Bands: E2

Solution	Answer	Mark
$(x, y) = \left(\frac{3(9) + 2(-1)}{3+2}, \frac{3(-6) + 2(4)}{3+2} \right)$ $= (5, -2)$	D	1

Question 3 (1 mark)

Outcomes Assessed: PE3

Targeted Performance Bands: E2

Solution	Answer	Mark
$P\left(\frac{-1}{2}\right) = 0 \Rightarrow 8\left(\frac{-1}{2}\right)^3 + a\left(\frac{-1}{2}\right)^2 - 4\left(\frac{-1}{2}\right) + 1 = 0$ $\therefore a = -8$	A	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 4 (1 mark)**Outcomes Assessed: PE3****Targeted Performance Bands: E3**

Solution	Answer	Mark
<p>Total number of arrangements of TWITTER = $\frac{7!}{3!}$</p> <p>Number of arrangements with TTT grouped is the same as the number of arrangements of TWIER = $5!$</p> <p>\therefore probability Ts are grouped together = $\frac{5!}{\left(\frac{7!}{3!}\right)} = \frac{1}{7}$</p>	C	1

Question 5 (1 mark)**Outcomes Assessed: PE3****Targeted Performance Bands: E2**

Solution	Answer	Mark
$\alpha + \beta + \gamma = -\frac{-5}{1} = 5$ $\alpha\beta + \alpha\gamma + \beta\gamma = \frac{1}{1} = 1$ $\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma)$ $= 5^2 - 2 \times 1$ $= 23$	B	1

Question 6 (1 mark)**Outcomes Assessed: PE2, PE3****Targeted Performance Bands: E3**

Solution	Answer	Mark
<p>The coefficient of x^3 in $(1-3x)^7$ is</p> $\binom{7}{3} 1^4 (-3)^3 = -945$	A	1

Question 7 (1 mark)**Outcomes Assessed: PE2****Targeted Performance Bands: E2**

Solution	Answer	Mark
$\sin x - \sqrt{3} \cos x \equiv 2 \sin x \cos \alpha + 2 \cos x \sin \alpha$ $\therefore 2 \cos \alpha = 1, 2 \sin \alpha = -\sqrt{3} \Rightarrow \alpha = -\frac{\pi}{3}$	B	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 8 (1 mark)**Outcomes Assessed: HE4****Targeted Performance Bands: E3**

Solution	Answer	Mark
$\frac{d}{dx} \sin^{-1}\left(\frac{3x}{4}\right) = \frac{1}{\sqrt{1-\left(\frac{3x}{4}\right)^2}} \times \frac{3}{4}$ $= \frac{3}{\sqrt{16-9x^2}}$	D	1

Question 9 (1 mark)**Outcomes Assessed: HE3****Targeted Performance Bands: E2**

Solution	Answer	Mark
<p>The vertical motion is the same for all three stones.</p> <p>\therefore All three stones reach the ground at the same time.</p>	D	1

Question 10 (1 mark)**Outcomes Assessed: HE4****Targeted Performance Bands: E4**

Solution	Answer	Mark
<p>The graph of $y = f^{-1}(x)$ is the reflection of the graph of $y = f(x)$ about the line $y = x$.</p> $\int_{-2}^6 f^{-1}(x) dx = \text{area } A + \text{area of } 4 \times 6 \text{ rectangle} - \text{area } B$ $= 3 + 24 - 7$ $= 20 \text{ square units}$	C	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Section II

60 marks

Question 11 (15 marks)

(a) (2 marks)

Outcomes assessed: HE6

Targeted Performance Bands: E2

Criteria	Mark
• Correct answer in terms of π	2
• Correct integration of $\frac{1}{\sqrt{4-x^2}}$	1

Sample answer:

$$\begin{aligned}\int_0^1 \frac{dx}{\sqrt{4-x^2}} &= \left[\sin^{-1} \frac{x}{2} \right]_0^1 \\ &= \sin^{-1} \frac{1}{2} - \sin^{-1} 0 \\ &= \frac{\pi}{6}\end{aligned}$$

(b) (3 marks)

Outcomes Assessed: PE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct range of solutions given	3
• Significant progress towards solution (e.g. inequality if using solution presented below or points of intersection if using graphical techniques)	2
• Some progress towards answer (considering technique used e.g. graphical or algebraic)	1

Sample answer:

$$\begin{aligned}\frac{t-2}{t+3} &> -2 \\ (t+3)(t-2) &> -2(t+3)^2 \\ 3t^2 + 13t + 12 &> 0 \\ (3t+4)(t+3) &> 0 \\ \therefore t &< -3, t > \frac{-4}{3}\end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (3 marks)

Outcomes Assessed: HE6

Targeted Performance Bands: E3

Criteria	Mark
• Correct answer	3
• Significant progress towards solution	2
• Correct integral after applying given substitution	1

Sample answer:

$$\text{Let } u = \sqrt{x}, du = \frac{dx}{2\sqrt{x}}$$

$$\begin{aligned}\int \frac{e^{3\sqrt{x}}}{\sqrt{x}} dx &= 2 \int \frac{e^{3\sqrt{x}}}{2\sqrt{x}} dx \\ &= 2 \int e^{3u} du \\ &= \frac{2}{3} e^{3u} + C \\ &= \frac{2}{3} e^{3\sqrt{x}} + C\end{aligned}$$

(d)(i) (2 marks)

Outcomes Assessed: HE5

Targeted Performance Bands: E2

Criteria	Mark
• Correct derivative	2
• Significant progress applying the product rule	1

Sample answer:

$$\frac{d}{dx}(x \sin 2x) = 2x \cos 2x + \sin 2x$$

(d)(ii) (2 marks)

Outcomes Assessed: PE5

Targeted Performance Bands: E3

Criteria	Mark
• Determines correct integral	2
• Significant progress towards the answer	1

Sample answer:

$$\begin{aligned}\int 2x \cos 2x dx + \int \sin 2x dx &= x \sin 2x \\ \int x \cos 2x dx &= \frac{1}{2} x \sin 2x - \frac{1}{2} \int \sin 2x dx \\ &= \frac{1}{2} x \sin 2x + \frac{1}{4} \cos 2x + C\end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(e) (3 marks)

Outcomes Assessed: PE3

Targeted Performance Bands: E3

Criteria	Mark
• Correctly finds the height of the tower	3
• Makes significant progress towards finding the height of the tower	2
• Finds the correct expression for AC or BC in terms of h	1

Sample answer:

$$\tan 30^\circ = \frac{h}{BC} \Rightarrow BC = \sqrt{3}h$$

$$\tan 45^\circ = \frac{h}{AC} \Rightarrow AC = h$$

$$(\sqrt{3}h)^2 - h^2 = 400^2$$

$$2h^2 = 400^2$$

$$h = 200\sqrt{2}$$

\therefore The height of the tower is $200\sqrt{2}$ metres.

Question 12 (15 marks)

(a)(i) (1 mark)

Outcomes assessed: HE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct verification that T is a solution of the differential equation	1

Sample answer:

$$T = 20 + Ae^{-kt}$$

$$\frac{dT}{dt} = -kAe^{-kt}$$

$$= -k(T - 20) \text{ since } Ae^{-kt} = T - 20$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(a)(ii) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct values for A and k	2
• Significant progress towards solution	1

Sample answer:

$$t = 0, T = 100 \Rightarrow A = 80$$

$$t = 5, T = 70 \Rightarrow 70 = 20 + 80e^{-5k}$$

$$e^{-5k} = \frac{5}{8}$$

$$\therefore k = \frac{-1}{5} \ln\left(\frac{5}{8}\right) \approx 0.094 \text{ (3 d.p.)}$$

(a)(iii) (1 mark)

Outcomes assessed: HE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct answer	1

Sample answer:

$$T = 20 + 80e^{\frac{1}{5} \ln\left(\frac{5}{8}\right) \times 15}$$

$$= 39.531...$$

$$\approx 40 \text{ (nearest degree)}$$

\therefore The temperature of the soup after 15 minutes is 40°C (to the nearest degree)

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.


(b) (2 marks)

Outcomes assessed: HE4

Targeted Performance Bands: E3

Criteria	Mark
• Correct sketch	2
• Correct shape with either correct domain or range	1

Sample answer:



(c) (2 marks)

Outcomes assessed: PE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct answer	2
• Some progress towards the correct answer	1

Sample answer:

If Jack is seated first, there are 4 seats (not next to Jack) in which Jill can sit. There are then 5! ways to seat the remaining students.

\therefore The total number of ways is $1 \times 4 \times 5! = 480$.

(d)(i) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct solution, including mentioning $f(x)$ is continuous	2
• Evaluates $f(0.7)$ and $f(0.8)$	1

Sample answer:

$$f(0.7) = \ln 0.7 - \sin 0.7 + 1 = -0.00089... < 0$$

$$f(0.8) = \ln 0.8 - \sin 0.8 + 1 = 0.05950... > 0$$

Since $f(x)$ is continuous for $x > 0$ and changes sign from $x = 0.7$ to $x = 0.8$, $f(x)$ has a zero between 0.7 and 0.8.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(d)(ii) (2 marks)

Outcomes assessed: PE5

Targeted Performance Bands: E3

Criteria	Mark
• Correct solution	2
• Correct application using halving-the-interval method	1

Sample answer:

$$f(0.75) = \ln 0.75 - \sin 0.75 + 1 = 0.030679... > 0$$

$\therefore f(x)$ has a zero between 0.7 and 0.75, hence the zero is 0.7 correct to one decimal place.


(e) (3 marks)

Outcomes assessed: PE3, HE7

Targeted Performance Bands: E3

Criteria	Mark
• Correct proof	3
• Significant progress towards required result	2
• Establishes one correct pair of equal or complementary angles leading to result	1

Sample answer:



Let $\angle TPB = \theta$

Hence, $\angle PAB = \theta$ (The angle between a tangent and a chord equals the angle in the alternate segment).

$\angle APB = 90^\circ$ (angle subtended at the circumference by a diameter is 90°)

$\angle PBA = 90^\circ - \theta$ (angle sum of $\triangle APB$ is 180°)

$\angle BPN = \theta$ (angle sum of $\triangle NPB$ is 180°)

$\therefore \angle TPB = \angle BPN$, hence BP bisects $\angle NPT$.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 13 (15 marks)

(a)(i) (1 mark)

Outcomes assessed: HE3, HE5**Targeted Performance Bands: E3**

Criteria	Mark
• Correct expression for acceleration in terms of x	1

Sample answer:

$$\begin{aligned}\ddot{x} &= \frac{d}{dx} \left(\frac{1}{2} v^2 \right) \\ &= \frac{d}{dx} \left(-\frac{1}{2} x^2 - 2x + 6 \right) \\ &= -x - 2\end{aligned}$$

(a)(ii) (2 marks)

Outcomes assessed: HE3, HE5**Targeted Performance Bands: E3**

Criteria	Mark
• Correct centre and period of motion	2
• Correct centre or period of motion	1

Sample answer:

Since $\ddot{x} = -1^2(x+2)$,

Centre of motion is $x = -2$.

Period of motion is $\frac{2\pi}{n} = \frac{2\pi}{1} = 2\pi$ seconds.

(a)(iii) (1 mark)

Outcomes assessed: HE3, HE5**Targeted Performance Bands: E3**

Criteria	Mark
• Correct maximum speed	1

Sample answer:

Maximum speed occurs at the centre of motion, $x = -2$.

$$v^2 = -(-2)^2 - 4(-2) + 12 = 16$$

$$v = \pm 4$$

\therefore The maximum speed of the particle is 4 ms^{-1} .

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (3 marks)

Outcomes assessed: PE3

Targeted Performance Bands: E4

Criteria	Mark
• Correct co-ordinates for P	3
• Correct solutions for θ	2
• Significant progress towards solving $2 \cos \theta - \cos 2\theta = 1$	1

Sample answer:

Solve $x = 1$ and $x = 2 \cos \theta - \cos 2\theta$ simultaneously for points of intersection:

$$2 \cos \theta - \cos 2\theta = 1$$

$$2 \cos \theta - 2 \cos^2 \theta + 1 = 1$$

$$2 \cos \theta (1 - \cos \theta) = 0$$

$$\cos \theta = 0, \cos \theta = 1$$

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2}, 0 \text{ for } 0 \leq \theta < 2\pi$$

$$\theta = \frac{\pi}{2} \Rightarrow y = 2 \sin \frac{\pi}{2} - \sin \pi = 2 \text{ which is in the first quadrant,}$$

Hence P has coordinates $(1, 2)$.

$$\text{Note: } \theta = \frac{3\pi}{2} \Rightarrow y = 2 \sin \frac{3\pi}{2} - \sin 3\pi = -2 \text{ which is the point of intersection in the 4}^{\text{th}} \text{ quadrant}$$

$$\text{and } \theta = 0 \Rightarrow y = 2 \sin 0 - \sin 0 = 0 \text{ which is the point of intersection on the x-axis.}$$

(c)(i) (2 marks)

Outcomes assessed: PE3

Targeted Performance Bands: E2

Criteria	Mark
• Correct proof	2
• Some progress towards the proof	1

Sample answer:

$$m_{OP} = \frac{ap^2}{2ap} = \frac{p}{2}$$

$$m_{OQ} = \frac{aq^2}{2aq} = \frac{q}{2}$$

$$OP \perp OQ \Rightarrow \frac{p}{2} \times \frac{q}{2} = -1.$$

$$\therefore pq = -4.$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c)(ii) (2 marks)

Outcomes assessed: PE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct equation of the parabola	2
• Some attempt to eliminate p and q	1

Sample answer:

$$x = 2a(p + q) \Rightarrow p + q = \frac{x}{2a}$$

$$y = a(p^2 + q^2) \Rightarrow p^2 + q^2 = \frac{y}{a}$$

$$pq = -4 \text{ (from part i)}$$

$$(p + q)^2 = p^2 + q^2 + 2pq$$

$$\Rightarrow \left(\frac{x}{2a}\right)^2 = \frac{y}{a} + 2(-4)$$

$$\Rightarrow \frac{x^2}{4a^2} = \frac{y}{a} - 8$$

$$\Rightarrow x^2 = 4a(y - 8a)$$

\therefore The equation of the locus of R is a parabola.

(c)(iii) (2 marks)

Outcomes assessed: HE5

Targeted Performance Bands: E3

Criteria	Mark
• Correct solution	2
• Progress towards correct solution	1

Sample answer:

$$pq = -4 \Rightarrow q = \frac{-4}{p} \Rightarrow \frac{dq}{dp} = \frac{4}{p^2}$$

$$\frac{dq}{dt} = \frac{dq}{dp} \times \frac{dp}{dt}$$

$$= \frac{4}{p^2} \times \frac{dp}{dt}$$

$$= \frac{-q}{p} \times \frac{dp}{dt} \text{ (since } pq = -4, \frac{4}{p} = -q \text{)}.$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c)(iv) (2 marks)

Outcomes assessed: HE5

Targeted Performance Bands: E4

Criteria	Mark
• Correct solution	2
• Progress towards finding an expression for $\frac{dy_Q}{dt}$ by applying the chain rule	1

Sample answer:

Given $\frac{dy_P}{dt} = 1$, we are required to prove $\frac{dy_Q}{dt} = \frac{-q^2}{p^2}$

$$y_P = ap^2 \Rightarrow \frac{dy_P}{dy_p} = 2ap$$

$$y_Q = aq^2 \Rightarrow \frac{dy_Q}{dy_q} = 2aq$$

$$\frac{dy_P}{dt} = \frac{dy_P}{dp} \times \frac{dp}{dt}$$

$$1 = 2ap \times \frac{dp}{dt}$$

$$\frac{dp}{dt} = \frac{1}{2ap}$$

$$\frac{dy_Q}{dt} = \frac{dy_Q}{dq} \times \frac{dq}{dt}$$

$$= 2aq \times \frac{-q}{p} \times \frac{dp}{dt} \text{ (using part (i))}$$

$$= 2aq \times \frac{-q}{p} \times \frac{1}{2ap} \text{ (from above)}$$

$$= \frac{-q^2}{p^2}$$

\therefore The y -coordinate of Q is decreasing at the rate $\frac{q^2}{p^2}$ units per second.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 14 (15 marks)

(a) (4 marks)

Outcomes assessed: HE2**Targeted Performance Bands:** E4

Criteria	Mark
• Complete proof	4
• Significant progress towards the correct proof	3
• Makes use of the assumption	2
• Proof for $P(1)$	1

Sample answer:

Let $P(n)$ be the given proposition. $P(1)$ is true since $1^3 + 5(1) = 6$ which is divisible by 6.

Assume $P(k)$ is true for some positive integer k .

i.e. $k^3 + 5k = 6M$ for some integer M .

Prove $P(k+1)$ is true:

$$\begin{aligned}
 (k+1)^3 + 5(k+1) &= k^3 + 3k^2 + 8k + 6 \\
 &= 6M - 5k + 3k^2 + 8k + 6 \text{ (using assumption)} \\
 &= 6M + 6 + 3k^2 + 3k \\
 &= 6(M+1) + 3k(k+1).
 \end{aligned}$$

Since either k or $k+1$ is even, $k(k+1)$ is divisible by 2. $\therefore 3k(k+1)$ is divisible by 6.

Also, as M is an integer, $6(M+1)$ is divisible by 6.

Hence, the above expression is divisible by 6.

\therefore By the Principle of Mathematical Induction, $P(n)$ is true for integers $n \geq 1$.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b)(i) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct simplified expression	2
• Correct expansion of $(a+b)^n$ or $(a-b)^n$	1

Sample answer:

$$(a+b)^n = \binom{n}{0}a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \binom{n}{3}a^{n-3}b^3 + \binom{n}{4}a^{n-4}b^4 + \dots + \binom{n}{n}b^n$$

$$(a-b)^n = \binom{n}{0}a^n - \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 - \binom{n}{3}a^{n-3}b^3 + \binom{n}{4}a^{n-4}b^4 + \dots + \binom{n}{n}b^n$$

$$(a+b)^n + (a-b)^n = 2\binom{n}{0}a^n + 2\binom{n}{2}a^{n-2}b^2 + 2\binom{n}{4}a^{n-4}b^4 + \dots + 2\binom{n}{n}b^n$$

(b)(ii) (1 mark)

Outcomes assessed: HE3

Targeted Performance Bands: E3

Criteria	Mark
• Correct expression	1

Sample answer:

The probability of obtaining exactly two sixes is given by $\binom{n}{2}\left(\frac{1}{6}\right)^2\left(\frac{5}{6}\right)^{n-2}$.

(b)(iii) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E4

Criteria	Mark
• Complete proof	2
• Some progress towards an expression for the correct probability	1

Sample answer:

$$\begin{aligned} P(\text{even number of sixes}) &= \binom{n}{0}\left(\frac{5}{6}\right)^n + \binom{n}{2}\left(\frac{1}{6}\right)^2\left(\frac{5}{6}\right)^{n-2} + \binom{n}{4}\left(\frac{1}{6}\right)^4\left(\frac{5}{6}\right)^{n-4} + \dots + \binom{n}{n}\left(\frac{1}{6}\right)^n \\ &= \frac{1}{2}\left(\left(\frac{5}{6} + \frac{1}{6}\right)^n + \left(\frac{5}{6} - \frac{1}{6}\right)^n\right) \text{ using part i} \\ &= \frac{1}{2}\left(1 + \left(\frac{2}{3}\right)^n\right) \end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c)(i) (1 mark)

Outcomes assessed: HE7

Targeted Performance Bands: E3

Criteria	Mark
• Correct answer	1

Sample answer:

$$0 \leq t \leq 1$$

(c)(ii) (3 marks)

Outcomes assessed: HE7

Targeted Performance Bands: E4

Criteria	Mark
• Correct proof and expression for y -coordinate	3
• Significant progress towards the correct proof and expression for y -coordinate	2
• Progress towards finding the equation of AP and solving simultaneously for P	1

Sample answer:

The equation of the line through $A(-1, 0)$ with gradient t is $y = t(x + 1)$.

For the coordinates of P solve $y = t(x + 1)$ and $x^2 + y^2 = 1$ simultaneously:

$$\begin{aligned}x^2 + (t(x + 1))^2 &= 1 \\(1 + t^2)x^2 + 2xt^2 + (t^2 - 1) &= 0\end{aligned}$$

Since one of the roots of this quadratic is $x = -1$ (x -coordinate of A) and the product of roots is

$$\frac{t^2 - 1}{1 + t^2}, \text{ the other root is } \frac{1 - t^2}{1 + t^2}.$$

Hence the x coordinate of P is $\frac{1 - t^2}{1 + t^2}$.

Substituting into $y = t(x + 1)$ gives $y = t\left(\frac{1 - t^2}{1 + t^2} + 1\right) = \frac{2t}{1 + t^2}$

Hence the y coordinate of P is $\frac{2t}{1 + t^2}$.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c)(iii) (1 mark)

Outcomes assessed: HE7

Targeted Performance Bands: E3

Criteria	Mark
• Correct solution	1

Sample answer:

$$\tan \theta = m_{OP}$$

$$= \frac{2t}{1+t^2} \div \frac{1-t^2}{1+t^2}$$

$$= \frac{2t}{1-t^2}, t \neq 1$$

(c)(iv) (1 mark)

Outcomes assessed: HE7

Targeted Performance Bands: E4

Criteria	Mark
• Correct value for t	1

Sample answer:

$\theta = 2\alpha$ since the angle at the centre of a circle is twice the angle at the circumference, subtended by the same arc.

$$\alpha = 22\frac{1}{2}^\circ \Rightarrow \theta = 45^\circ$$

$$\therefore \tan \theta = 1$$

$$\text{Hence, } \frac{2t}{1-t^2} = 1$$

$$t^2 + 2t - 1 = 0$$

$$t = \frac{-2 \pm \sqrt{8}}{2}$$

$$t = -1 \pm \sqrt{2}$$

$$t = -1 + \sqrt{2}, \text{ since } 0 \leq t \leq 1$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.