

Question 1

Begin a new page


(a) Find the value of $\sum_{k=1}^4 (-1)^k k!$ 2

(b) $A(-2, -5)$ and $B(1, 4)$ are two points. Find the acute angle θ between the line AB and the line $x + 2y + 1 = 0$, giving the answer correct to the nearest minute. 3

(c) The polynomial $P(x) = x^5 + ax^3 + bx$ leaves a remainder of 5 when it is divided by $(x - 2)$, where a and b are numerical constants.

- (i) Show that $P(x)$ is odd. 1
 (ii) Hence find the remainder when $P(x)$ is divided by $(x + 2)$. 2

(d)




$ABCD$ is a cyclic quadrilateral. The tangent at A to the circle through A , B , C and D is parallel to BD .

- (i) Copy the diagram. 1
 (ii) Give a reason why $\angle ACB = \angle MAB$. 1
 (iii) Give a reason why $\angle ACD = \angle ABD$. 2
 (iv) Hence show that AC bisects $\angle BCD$.

Question 2

Begin a new page

- (a) Find $\frac{d^2}{dx^2} e^{x^2}$. 2
- (b) $A(-1, 4)$ and $B(x, y)$ are two points. The point $P(14, -6)$ divides the interval AB externally in the ratio $5:3$. Find the coordinates of B . 3
- (c) Find the number of ways in which the letters of the word EXTENSION can be arranged in a straight line so that no two vowels are next to each other. 3
- (d)



$P(2t, t^2)$ is a variable point which moves on the parabola $x^2 = 4y$. The tangent to the parabola at P cuts the x axis at T . M is the midpoint of PT .

- (i) Show that the tangent PT has equation $tx - y - t^2 = 0$. 1
- (ii) Show that M has coordinates $\left(\frac{3t}{2}, \frac{t^2}{2}\right)$. 2
- (iii) Hence find the Cartesian equation of the locus of M as P moves on the parabola. 1

Question 3


Begin a new page

- (a) (i) By expanding $\cos(2A + A)$, show that $\cos 3A = 4\cos^3 A - 3\cos A$. 2
- (ii) Hence show that if $2\cos A = x + \frac{1}{x}$ then $2\cos 3A = x^3 + \frac{1}{x^3}$. 3
- (b) The function $f(x)$ is given by $f(x) = \sqrt{x+6}$ for $x \geq -6$.
- (i) Find the inverse function $f^{-1}(x)$ and find its domain. 2
- (ii) On the same diagram, sketch the graphs of $y = f(x)$ and $y = f^{-1}(x)$, showing clearly the intercepts on the coordinate axes. Draw in the line $y = x$. 3
- (iii) Show that the x coordinates of any points of intersection of the graphs $y = f(x)$ and $y = f^{-1}(x)$ satisfy the equation $x^2 - x - 6 = 0$. Hence find any points of intersection of the two graphs. 2

Question 4

Begin a new page

- (a) Use Mathematical Induction to show that $5^n + 2(11^n)$ is a multiple of 3 for all positive integers n . 5
- (b)



Two concentric circles with centre O have radii 2 cm and 4 cm. The points P and Q lie on the larger circle and $\angle POQ = x$, where $0 < x < \frac{\pi}{2}$.

- (i) If the area $A \text{ cm}^2$ of the shaded region is $\frac{1}{16}$ the area of the larger circle, show that x satisfies the equation $8\sin x - 2x - \pi = 0$. 3
- (ii) Show that this equation has a solution $x = \alpha$, where $0.5 < \alpha < 0.6$. 2
- (iii) Taking 0.6 as a first approximation for α , use one application of Newton's Method to find a second approximation, giving the answer correct to two decimal places. 2

Question 5


Begin a new page

- (a) Evaluate $\int_1^{49} \frac{1}{4(x + \sqrt{x})} dx$ using the substitution $u^2 = x$, $u > 0$. Give the answer in simplest exact form. 4
- (b) At any point on the curve $y = f(x)$, the gradient function is given by $\frac{dy}{dx} = \sin^2 x$. 4
Find the value of $f\left(\frac{3\pi}{4}\right) - f\left(\frac{\pi}{4}\right)$.
- (c) A particle is performing Simple Harmonic Motion in a straight line. At time t seconds it has velocity v metres per second, and displacement x metres from a fixed point O on the line, where $x = 5 \cos \frac{\pi t}{2}$. 1
- (i) Find the period of the motion. 1
- (ii) Find an expression for v in terms of t , and hence show that $v^2 = \frac{\pi^2}{4} (25 - x^2)$. 3
Find the speed of the particle when it is 4 metres to the right of O .

Question 6

Begin a new page

(a)



A person on horizontal ground is looking at an aeroplane A through a telescope T . The aeroplane is approaching at a speed of 80 ms^{-1} at a constant altitude of 200 metres above the telescope. When the horizontal distance of the aeroplane from the telescope is x metres, the angle of elevation of the aeroplane is θ radians.

(i) Show that $\theta = \tan^{-1} \frac{200}{x}$. 1

(ii) Show that $\frac{d\theta}{dt} = \frac{16000}{x^2 + 40000}$. 2

(iii) Find the rate at which θ is changing when $\theta = \frac{\pi}{4}$, giving the answer in degrees per second correct to the nearest degree. 2

(b) A particle moves in a straight line. At time t seconds its displacement is x metres from a fixed point O on the line, its acceleration is $a \text{ ms}^{-2}$, and its velocity is $v \text{ ms}^{-1}$ where v is given by $v = \frac{32}{x} - \frac{x}{2}$.

(i) Find an expression for a in terms of x . 1

(ii) Show that $t = \int \frac{2x}{64 - x^2} dx$, and hence show that $x^2 = 64 - 60e^{-t}$. 3

(iii) Sketch the graph of x^2 against t and describe the limiting behaviour of the particle. 3

Question 7

Begin a new page

- (a) Four fair dice are rolled. Any die showing 6 is left alone, while the remaining dice are rolled again.
- (i) Find the probability (correct to 2 decimal places) that after the first roll of the dice, exactly one of the four dice is showing 6. 1
- (ii) Find the probability (correct to 2 decimal places) that after the second roll of the dice exactly two of the four dice are showing 6. 4
- (b) A particle is projected from a point O with speed 50 ms^{-1} at an angle of elevation θ , and moves freely under gravity, where $g = 10 \text{ ms}^{-2}$.
- (i) Write down expressions for the horizontal and vertical displacements of the particle at time t seconds referred to axes Ox and Oy . 1
- (ii) Hence show that the equation of the path of the projectile, given as a quadratic equation in $\tan \theta$, is $x^2 \tan^2 \theta - 500x \tan \theta + (x^2 + 500y) = 0$. 2
- (iii) Hence show that there are two values of θ , $0 < \theta < \frac{\pi}{2}$, for which the projectile passes through a given point (X, Y) provided that $500Y < 62500 - X^2$. 2
- (iv) If the projectile passes through the point (X, X) whose coordinates satisfy this inequality, and the two values of θ are α and β , find expressions in terms of X for $\tan \alpha + \tan \beta$ and $\tan \alpha \tan \beta$, and hence show that $\alpha + \beta = \frac{3\pi}{4}$. 2

MATHEMATICS EXTENSION 1
TRIAL EXAMINATION 2001

AMENDMENTS:

Page 4 QUESTION 2(c) Find the number of ways in which the letters of the word EXTENSION can be arranged in a straight line so that no two vowels **and no two consonants** are next to each other.

Page 7 QUESTION 6(b) A particle moves in a straight line. At time t seconds its displacement is x metres from a fixed point O on the line, its acceleration is $a \text{ ms}^{-2}$, and its velocity is $v \text{ ms}^{-1}$ where v is given by $v = \frac{32}{x} - \frac{x}{2}$. **The particle starts 2 metres to the right of O**

Question 1

(a) **Outcomes Assessed: H5, H9**

Marking Guidelines	
Criteria	Marks
<ul style="list-style-type: none"> • one mark for simplification of sum • one mark for value of sum 	2

Answer:

$$\sum_{k=1}^4 (-1)^k k! = -1! + 2! - 3! + 4! = 19$$

(b) **Outcomes Assessed: P4**

Marking Guidelines	
Criteria	Marks
<ul style="list-style-type: none"> • one mark for values of gradients • one mark for value of $\tan \theta$ • one mark for size of angle 	3

Answer:

$$\left. \begin{array}{l} AB \text{ has gradient } m_1 = 3 \\ x + 2y + 1 = 0 \text{ has gradient } m_2 = -\frac{1}{2} \end{array} \right\} \Rightarrow \tan \theta = \left| \frac{3 - (-\frac{1}{2})}{1 + 3(-\frac{1}{2})} \right| = 7 \quad \therefore \theta = 81^\circ 52'$$

(c) **Outcomes Assessed: (i) P5 (ii) PE3**

Marking Guidelines	
Criteria	Marks
(i) • one mark for showing $P(x)$ is odd (ii) • one mark for showing remainder is $-P(2)$ • one mark for value of remainder	3

Answer:

(i)

$$\begin{aligned} P(-x) &= (-x)^5 + a(-x)^3 + b(-x) \\ &= -x^5 - ax^3 - bx \\ &= -(x^5 + ax^3 + bx) \\ &= -P(x) \quad \text{for all } x \\ \therefore P(x) &\text{ is odd.} \end{aligned}$$

(ii) When $P(x)$ is divided by $(x+2)$,

$$\begin{aligned} \text{remainder is } P(-2) &= -P(2) \text{ since } P(x) \text{ is odd} \\ &= -5 \quad \text{since } P(2) = 5 \end{aligned}$$


(d) Outcomes Assessed: (i) (ii) PE3 (iii) PE3 (iv) H5, PE2, PE3

Marking Guidelines

Criteria	Marks
(i) • no marks for copying diagram (ii) • one mark for reason (iii) • one mark for reason (iv) • one mark for showing $\angle MAB = \angle ABD$ • one mark for showing $\angle ACB = \angle ACD$	4

Answer:

(i)



(iii) $\angle ACD = \angle ABD$ because the angles subtended in the same segment at B and C by the arc AD are equal.

(iv)

$\angle MAB = \angle ABD$ (equal alternate angles, $MN \parallel BD$)

$\angle ACB = \angle ACD$ ($\angle MAB = \angle ACB$, $\angle ABD = \angle ACD$)

$\therefore AC$ bisects $\angle BCD$

(ii) $\angle ACB = \angle MAB$ because the angle between the tangent MA and the chord AB through the point of contact A is equal to the angle ACB in the alternate segment.

Question 2

(a) Outcomes Assessed: P7, PE5

Marking Guidelines

Criteria	Marks
• one mark for first derivative	2
• one mark for second derivative using product rule.	

Answer:

$$\frac{d}{dx} e^{x^2} = 2x e^{x^2} \quad \frac{d^2}{dx^2} e^{x^2} = \frac{d}{dx} 2x e^{x^2} = 2(e^{x^2}) + (2x)(2x e^{x^2}) = 2(1 + 2x^2) e^{x^2}$$

(b) Outcomes Assessed: P4

Marking Guidelines

Criteria	Marks
• one mark for equation in x • one mark for equation in y • one mark for coordinates of B	3

Answer:

$$\frac{5x - 3 \times (-1)}{5 - 3} = 14 \Rightarrow 5x + 3 = 28 \quad \therefore x = 5$$

$$\therefore B(5, 0)$$

$$\frac{5y - 3 \times (4)}{5 - 3} = -6 \Rightarrow 5y - 12 = -12 \quad \therefore y = 0$$

Marking Guidelines

Criteria	Marks
<ul style="list-style-type: none"> • one mark for number of arrangements of vowels • one mark for number of arrangements of consonants • one mark for total number of arrangements 	3

Answer:

The vowels (E, E, I, O) can be arranged in positions 2, 4, 6, 8 in $\frac{4!}{2!} = 12$ ways.

The consonants (N, N, S, T, X) can be arranged in positions 1, 3, 5, 7, 9 in $\frac{5!}{2!} = 60$ ways.


Hence the total number of arrangements is $12 \times 60 = 720$.

(d) Outcomes Assessed: (i) PE3, PE4 (ii) PE3 (iii) PE3

Marking Guidelines

Criteria	Marks
(i) • one mark for equation of tangent (ii) • one mark for coordinates of T • one mark for coordinates of M (iii) • one mark for equation of locus	4

Answer:



$$(i) \quad y = \frac{1}{4}x^2 \Rightarrow \frac{dy}{dx} = \frac{1}{2}x$$

\therefore tangent at $P(2t, t^2)$ has gradient $\frac{1}{2}(2t) = t$

and equation $y - t^2 = t(x - 2t)$

$$tx - y - t^2 = 0$$

$$(ii) \text{ At } T, \quad y = 0 \Rightarrow tx - 0 - t^2 = 0 \Rightarrow x = t$$

Hence T has coordinates $(t, 0)$, and

M is the midpoint of $P(2t, t^2)$ and $T(t, 0)$,

with coordinates $\left(\frac{2t+t}{2}, \frac{t^2+0}{2}\right) \equiv \left(\frac{3t}{2}, \frac{t^2}{2}\right)$.

$$(iii) \text{ At } M, \quad x = \frac{3t}{2} \Rightarrow t = \frac{2x}{3}$$

$$\therefore y = \frac{1}{2}t^2 = \frac{1}{2}\left(\frac{2x}{3}\right)^2 = \frac{2x^2}{9}$$

Hence the locus has equation $2x^2 = 9y$.

Question 3

(a) Outcomes Assessed: (i) P4 (ii) PE3

Marking Guidelines

Criteria	Marks
(i) • one mark for expansion and expressions for $\cos 2A$, $\sin 2A$ • one mark for simplification to obtain final expression for $\cos 3A$ in terms of $\cos A$ (ii) • one mark for expressing $2 \cos 3A$ in terms of $\left(x + \frac{1}{x}\right)$ • one mark for binomial expansion of $\left(x + \frac{1}{x}\right)^3$ • one mark for simplification to obtain final expression for $\cos 3A$ in terms of x	5

Answer:

(i)

$$\begin{aligned}
 \cos 3A &= \cos(2A + A) \\
 &= \cos 2A \cos A - \sin 2A \sin A \\
 &= (2 \cos^2 A - 1) \cos A - (2 \sin A \cos A) \sin A \\
 &= 2 \cos^3 A - \cos A - 2 \sin^2 A \cos A \\
 &= 2 \cos^3 A - \cos A - 2(1 - \cos^2 A) \cos A \\
 &= 4 \cos^3 A - 3 \cos A
 \end{aligned}$$

(ii)

$$\begin{aligned}
 2 \cos 3A &= 8 \cos^3 A - 6 \cos A \\
 &= (2 \cos A)^3 - 3(2 \cos A) \\
 &= \left(x + \frac{1}{x}\right)^3 - 3\left(x + \frac{1}{x}\right) \\
 &= x^3 + 3x + \frac{3}{x} + \left(\frac{1}{x}\right)^3 - 3x - \frac{3}{x} \\
 &= x^3 + \frac{1}{x^3}
 \end{aligned}$$

(b) Outcomes Assessed: (i) P5, HE4 (ii) P5, HE4 (iii) P4

Marking Guidelines

Criteria	Marks
(i) • one mark for finding the inverse function • one mark for the domain of the inverse function (ii) • one mark for the graph of $y = f(x)$ and intercepts • one mark for the graph of $y = f^{-1}(x)$ and intercepts • one mark for the line $y = x$ passing through the point of intersection (iii) • one mark for the equation • one mark for the coordinates of the point of intersection	7

Answer:

(i)

$$\begin{aligned}
 y &= \sqrt{x+6} && \text{Interchanging } x \text{ and } y \\
 y^2 &= x+6 && \text{gives } y = x^2 - 6 \\
 x &= y^2 - 6 && \therefore f^{-1}(x) = x^2 - 6
 \end{aligned}$$

$$\begin{aligned}
 \text{Range of } f(x) \text{ is } \{y : y \geq 0\} &\Rightarrow \text{Domain of } f^{-1}(x) \text{ is } \{x : x \geq 0\}
 \end{aligned}$$

(iii) Where $y = f(x)$, $y = f^{-1}(x)$, $y = x$ intersect,


$$f^{-1}(x) = x \Rightarrow x^2 - 6 = x$$

$$x^2 - x - 6 = 0$$

$$(x+2)(x-3) = 0$$

But $x \neq -2$ (outside domain). $\therefore x = 3$

(ii)



Hence intersection point of the curves is (3, 3).

Question 4

(a) Outcomes Assessed: HE2

Marking Guidelines

Criteria	Marks
<ul style="list-style-type: none"> • one mark for establishing the truth of $S(1)$ • one mark for $S(k)$ true $\Rightarrow 5^k + 2(11^k) = 3M$ for some integer M. • one mark for $5^{k+1} + 2(11^{k+1}) = 5(5^k) + 22(11^k)$ • one mark for deducing $S(k)$ true $\Rightarrow S(k+1)$ true • one mark for deducing $S(n)$ true for all integers $n \geq 1$ 	5

Answer:

Define the sequence of statements $S(n)$: $5^n + 2(11^n)$ is a multiple of 3, $n = 1, 2, 3, \dots$

Consider $S(1)$: $5^1 + 2(11^1) = 27 = 3 \times 9 \quad \therefore S(1)$ is true.

If $S(k)$ is true, then $5^k + 2(11^k) = 3M$ for some integer M . **

Consider $S(k+1)$: $5^{k+1} + 2(11^{k+1}) = 5(5^k) + 22(11^k) = 5\{5^k + 2(11^k)\} + 12(11^k)$
 $\therefore 5^{k+1} + 2(11^{k+1}) = 5(3M) + 12(11^k) = 3\{5M + 4(11^k)\}$ if $S(k)$ is true, using **

But M and k integral $\Rightarrow \{5M + 4(11^k)\}$ is an integer.

$\therefore S(k)$ true $\Rightarrow S(k+1)$ true, $k = 1, 2, 3, \dots$


Hence $S(1)$ is true, and if $S(k)$ is true, then $S(k+1)$ is true. $\therefore S(2)$ is true, and then $S(3)$ is true, and so on. Hence by Mathematical Induction, $S(n)$ is true for all positive integers n .

(b) Outcomes Assessed: (i) H5 (ii) P5, H2 (iii) PE3

Marking Guidelines

Criteria	Marks
(i) • one mark for areas of small circle sector and triangle OPQ • one mark for equating expression for shaded area to $\frac{1}{16}$ of large circle area • one mark for simplification to find equation in required form (ii) • one mark for showing $f(0.5), f(0.6)$ have opposite signs • one mark for using continuity of $f(x)$ to deduce $0.5 < \alpha < 0.6$ (iii) • one mark for expression for second approximation • one mark for calculation of second approximation	7

Answer:



(i)

Area of $\Delta POQ = \frac{1}{2}(4^2) \sin x$

Area small circle sector $= \frac{1}{2}(2^2)x$

\therefore shaded area $= 8 \sin x - 2x$

$\therefore 8 \sin x - 2x = \frac{1}{16} \pi (4^2) = \pi$

$8 \sin x - 2x - \pi = 0$

(ii) Let $f(x) = 8 \sin x - 2x - \pi$. Then

$f(0.5) \approx -0.31 < 0$ and $f(0.6) \approx 0.18 > 0$.

Hence, since $f(x)$ is continuous, $f(\alpha) = 0$ for some $0.5 < \alpha < 0.6$.

(iii) Taking a first approximation $\alpha \approx 0.6$,

Newton's method gives a second approximation

$$\begin{aligned} \alpha &\approx 0.6 - \frac{f(0.6)}{f'(0.6)} \\ &= 0.6 - \frac{8 \sin(0.6) - 2(0.6) - \pi}{8 \cos(0.6) - 2} \\ &\approx 0.56 \text{ to 2 decimal places.} \end{aligned}$$

Outcomes Assessed: HE6

Marking Guidelines

Criteria	Marks
<ul style="list-style-type: none"> • one mark for change of limits • one mark for change of variable • one mark for integration • one mark for evaluation 	4

Answer :

$$\text{Let } I = \int_1^{49} \frac{1}{4(x+\sqrt{x})} dx$$

$$u^2 = x, \quad u > 0$$

$$2u = \frac{dx}{du} \Rightarrow dx = 2u du$$

$$x=1 \Rightarrow u=1, \quad x=49 \Rightarrow u=7$$

$$\text{Then } I = \int_1^7 \frac{1}{4(u^2+u)} 2u du$$

$$= \int_1^7 \frac{1}{2(u+1)} du$$

$$= \frac{1}{2} [\ln(u+1)]_1^7$$

$$\therefore I = \frac{1}{2} (\ln 8 - \ln 2) = \frac{1}{2} \ln 4 = \ln 2$$

(b) Outcomes Assessed: H5

Marking Guidelines

Criteria	Marks
<ul style="list-style-type: none"> • one mark for expressing $\sin^2 x$ in terms of $\cos 2x$ • one mark for integration, including constant of integration • one mark for evaluation of $f\left(\frac{\pi}{4}\right), f\left(\frac{3\pi}{4}\right)$ • one mark for value of difference 	4

Answer:

$$\frac{dy}{dx} = \sin^2 x$$

$$= \frac{1}{2} (1 - \cos 2x)$$

$$y = \frac{1}{2} \left(x - \frac{1}{2} \sin 2x \right) + c, \quad c \text{ constant}$$

$$f(x) = \frac{1}{2} x - \frac{1}{4} \sin 2x + c$$

$$\therefore f\left(\frac{3\pi}{4}\right) - f\left(\frac{\pi}{4}\right)$$

$$= \left(\frac{3\pi}{8} - \frac{1}{4} \sin \frac{3\pi}{2} + c \right) - \left(\frac{\pi}{8} - \frac{1}{4} \sin \frac{\pi}{2} + c \right)$$

$$= \left(\frac{3\pi}{8} + \frac{1}{4} + c \right) - \left(\frac{\pi}{8} - \frac{1}{4} + c \right)$$

$$= \frac{\pi}{4} + \frac{1}{2}$$

(c) Outcomes Assessed: (i) HE3 (ii) H5, HE3

Marking Guidelines

Criteria	Marks
<ul style="list-style-type: none"> (i) • one mark for finding the period of the motion (ii) • one mark for expressing v^2 in terms of t <ul style="list-style-type: none"> • one mark for expressing v^2 in terms of x • one mark for the value of the speed. 	4

Answer:

(i) Period is $2\pi \div \frac{\pi}{2} = 4$ seconds

(ii)

$$x = 5 \cos \frac{\pi}{2} t$$

$$v = \frac{dx}{dt} = 5 \left(-\frac{\pi}{2} \sin \frac{\pi}{2} t \right)$$

$$v^2 = \left(\frac{\pi^2}{4} \right) \cdot 25 \sin^2 \frac{\pi}{2} t$$

$$v^2 = \left(\frac{\pi^2}{4} \right) \cdot 25 \left(1 - \cos^2 \frac{\pi}{2} t \right)$$

$$= \frac{\pi^2}{4} (25 - 25 \cos^2 \frac{\pi}{2} t)$$

$$v^2 = \frac{\pi^2}{4} (25 - x^2)$$

$$x = 4 \Rightarrow v^2 = \frac{\pi^2}{4} (25 - 16) = \frac{9\pi^2}{4}$$

$$\text{Speed is } \frac{3\pi}{2} \text{ ms}^{-1}$$


Question 6

(a) **Outcomes Assessed:** (i) P4, HE4 (ii) HE4, HE5 (iii) H5

Marking Guidelines	
Criteria	Marks
(i) • one mark for expression for θ (ii) • one mark for expression for $\frac{d\theta}{dx}$ • one mark for expression for $\frac{d\theta}{dt}$ (iii) • one mark for value of $\frac{d\theta}{dt}$ • one mark for value of θ	5

Answer:

(i)



$$\angle TAP = \theta$$

(alt. \angle s, parallel lines)

$$\tan \theta = \frac{200}{x}$$

$$\theta = \tan^{-1} \frac{200}{x}$$

(ii)

$$\frac{d\theta}{dx} = \frac{1}{1 + \left(\frac{200}{x}\right)^2} \left(-\frac{200}{x^2}\right) = \frac{-200}{x^2 + 40000}$$

$$\frac{d\theta}{dt} = \frac{d\theta}{dx} \cdot \frac{dx}{dt} = \frac{-200}{x^2 + 40000} (-80)$$

$$\therefore \frac{d\theta}{dt} = \frac{16000}{x^2 + 40000}$$

(iii) When $\theta = \frac{\pi}{4}$, $TP = AP \Rightarrow x = 200$, and $\frac{d\theta}{dt} = \frac{16000}{(200)^2 + 40000} = 0.2$ radians per second.

Hence θ is increasing at 11° s^{-1} (correct to the nearest degree)

(b) **Outcomes Assessed:** (i) HE5 (ii) H3, H5, HE4 (iii) HE3, HE7

Marking Guidelines	
Criteria	Marks
(i) • one mark for expression for a in terms of x (ii) • one mark for expressing t as an integral with respect to x • one mark for integration to find t in terms of x • one mark for expression for x^2 in terms of t (iii) • one mark for graph of x^2 as a function of t • one mark for limiting values of x , v , a • one mark for description of limiting behaviour in words	7

Answer:

(i)

$$v^2 = \left(\frac{32}{x} - \frac{x}{2}\right)^2 = \frac{1024}{x^2} - 32 + \frac{x^2}{4}$$

$$a = \frac{d}{dx} \left(\frac{1}{2} v^2\right) = \frac{1}{2} \frac{d}{dx} \left(\frac{1024}{x^2} - 32 + \frac{x^2}{4}\right)$$

$$\therefore a = \frac{-1024}{x^3} + \frac{x}{4}$$

(ii)

$$\frac{dx}{dt} = v = \frac{32}{x} - \frac{x}{2} = \frac{64 - x^2}{2x}$$

$$\therefore \frac{dt}{dx} = \frac{2x}{64 - x^2}$$

$$t = \int \frac{2x}{64 - x^2} dx$$


(iii)

(ii) Cont.

$$t = -\ln(64 - x^2) + c, \quad \left. \begin{matrix} t=0 \\ x=2 \end{matrix} \right\} \Rightarrow c = \ln 60$$

$$-t = \ln\left(\frac{64 - x^2}{60}\right), \quad e^{-t} = \frac{64 - x^2}{60}$$

$$\therefore x^2 = 64 - 60e^{-t}$$



$$\text{As } t \rightarrow \infty, \quad x \rightarrow 8^-, \quad v \rightarrow \frac{32}{8} - \frac{8}{2} = 0^+, \quad a \rightarrow \frac{-1024}{512} + \frac{8}{4} = 0^-$$

Hence the particle is moving right and slowing down as it approaches its limiting position 8 metres to the right of O .

Question 7

(a) **Outcomes Assessed:** (i) HE3 (ii) HE3

Marking Guidelines	
Criteria	Marks
(i) • one mark for value of probability (ii) • one mark for expression for probability of two 6's on first roll and no 6's on second • one mark for expression for probability of one 6 on first roll and one 6 on second • one mark for expression for probability of no 6's on first roll and two 6's on second • one mark for value of probability	5

Answer:

$$(i) P(\text{one 6 on first roll}) = {}^4C_1 \left(\frac{1}{6}\right)^1 \left(\frac{5}{6}\right)^3 \approx 0.39 \text{ (to 2 decimal places)}$$

$$(ii) P(\text{two 6's on first roll and no 6's on second roll}) = {}^4C_2 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right)^2 \times {}^2C_0 \left(\frac{1}{6}\right)^0 \left(\frac{5}{6}\right)^2 \approx 0.0804$$

$$P(\text{one 6 on first roll and one 6 on second roll}) = {}^4C_1 \left(\frac{1}{6}\right)^1 \left(\frac{5}{6}\right)^3 \times {}^3C_1 \left(\frac{1}{6}\right)^1 \left(\frac{5}{6}\right)^2 \approx 0.1340$$

$$P(\text{no 6's on first roll and two 6's on second roll}) = {}^4C_0 \left(\frac{1}{6}\right)^0 \left(\frac{5}{6}\right)^4 \times {}^4C_2 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right)^2 \approx 0.0558$$

$$\therefore P(\text{two 6's overall}) \approx 0.0804 + 0.1340 + 0.0558 \approx 0.27 \text{ (to 2 decimal places)}$$

(b) **Outcomes Assessed:** (i) HE3 (ii) HE3 (iii) P4, H2 (iv) P4, H2

Marking Guidelines	
Criteria	Marks
(i) • one mark for expressions for x and y in terms of θ and t (ii) • one mark for expression for y in terms of x • one mark for rearrangement as quadratic in $\tan \theta$ (iii) • one mark for discriminant in terms of X and Y • one mark for using discriminant > 0 to give required inequality (iv) • one mark for the values of the sum and product of $\tan \alpha$, $\tan \beta$ in terms of X • one mark for the value of $\alpha + \beta$	7

Answer:

(i) $x = 50 t \cos \theta$ and $y = 50 t \sin \theta - 5 t^2$

(ii) $t = \frac{x}{50 \cos \theta} \Rightarrow y = x \frac{\sin \theta}{\cos \theta} - \frac{5 x^2}{2500 \cos^2 \theta}$

$$500 y = 500 x \tan \theta - x^2 \sec^2 \theta$$

$$= 500 x \tan \theta - x^2 (1 + \tan^2 \theta)$$

$$= 500 x \tan \theta - x^2 - x^2 \tan^2 \theta$$

$$\therefore x^2 \tan^2 \theta - 500 x \tan \theta + (x^2 + 500 y) = 0$$

(iii) Projectile passes through the point (X, Y) if $\tan \theta$ satisfies the quadratic equation

$$X^2 \tan^2 \theta - 500 X \tan \theta + (X^2 + 500 Y) = 0$$

This equation has two distinct solutions for $\tan \theta$, and hence for θ , provided its discriminant $\Delta > 0$.

$$\Delta = (-500 X)^2 - 4 X^2 (X^2 + 500 Y)$$

$$= 4 X^2 (62500 - X^2 - 500 Y)$$

$$\therefore \Delta > 0 \text{ provided } 500 Y < 62500 - X^2$$

(iv) If the projectile passes through the point (X, X) where $500 X < 62500 - X^2$, then the equation

$$X^2 \tan^2 \theta - 500 X \tan \theta + (X^2 + 500 X) = 0 \text{ has two distinct real roots } \tan \alpha, \tan \beta \text{ where}$$

$$\tan \alpha + \tan \beta = \frac{500 X}{X^2} = \frac{500}{X} \quad \text{and} \quad \tan \alpha \tan \beta = \frac{X^2 + 500 X}{X^2} = 1 + \frac{500}{X}$$

$$\therefore \tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} = \frac{500}{X} + \left(-\frac{500}{X}\right) = -1$$

$$\text{Since } 0 < \alpha + \beta < \pi, \quad \alpha + \beta = \frac{3\pi}{4}.$$