



CATHOLIC SECONDARY SCHOOLS ASSOCIATION
2010 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION
MATHEMATICS EXTENSION 1

Question 1 (12 marks)

(a) (2 marks)

Outcomes assessed: PE2

Targeted Performance Bands: E2-E3

Criteria	Marks
• uses correct formula for division of interval or progress using other correct method	1
• finds correct coordinates from working	1

Sample Answer:

$A(x_1, y_1) = (-2, -1)$ and $B(x_2, y_2) = (1, 5)$; Q divides AB externally ie $m:n = 5:-2$

$$x_Q = \frac{-2 \times -2 + 5 \times 1}{5 + (-2)}$$

$$= \frac{9}{3}$$

$$= 3$$

$$y_Q = \frac{-2 \times -1 + 5 \times 5}{5 + (-2)}$$

$$= \frac{27}{3}$$

$$= 9$$

$\therefore Q$ has coordinates (3, 9)

(b) (2 marks)

Outcomes assessed: PE2

Targeted Performance Bands: E2-E3

Criteria	Marks
• correct trigonometric substitution	1
• completes the proof	1

Sample Answer:

$$\begin{aligned} \text{LHS} &= \frac{\sin 2x}{1 + \cos 2x} \\ &= \frac{2 \sin x \cos x}{1 + 2 \cos^2 x - 1} \\ &= \frac{2 \sin x \cos x}{2 \cos^2 x} \\ &= \frac{\sin x}{\cos x} \\ &= \tan x \\ &= \text{RHS} \end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (3 marks)

Outcomes assessed: PE3

Targeted Performance Bands: E2-E3

Criteria	Marks
• establishes correct quadratic or other correct significant step towards solution	1
• further progress towards solution	1
• finds correct solution	1

Sample Answer:

$$\frac{2x}{x-1} \geq 1 \quad \text{multiply by } (x-1)^2 \text{ with } x \neq 1$$

$$2x(x-1) \geq (x-1)^2$$

$$2x(x-1) - (x-1)^2 \geq 0$$

$$(x-1)(2x - (x-1)) \geq 0$$

$$(x-1)(x+1) \geq 0$$

$$\text{Solution is } x \leq -1 \text{ or } x > 1$$

(d) (2 marks)

Outcomes assessed: HE4

Targeted Performance Bands: E2-E3

Criteria	Marks
• gives correct exact trigonometric value	1
• correctly evaluates exact inverse trigonometric value	1

Sample Answer:

$$\begin{aligned} \sin^{-1}\left(\sin \frac{7\pi}{6}\right) &= \sin^{-1}\left(\frac{-1}{2}\right) \\ &= \frac{-\pi}{6} \end{aligned}$$

(e) (3 marks)

Outcomes assessed: HE6

Targeted Performance Bands: E2-E3

Criteria	Mark
• rewrites the integral using the substitution	1
• finds the correct primitive	1
• gives final result	1

Sample Answer:

$$\int \frac{dx}{x(\ln 3x)^2} = \int \frac{du}{u^2}$$

$$= -\frac{1}{u} + C$$

$$= \frac{-1}{\ln 3x} + C$$

$$u = \ln 3x$$

$$\frac{du}{dx} = \frac{1}{x} \Rightarrow du = \frac{dx}{x}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 2 (12 marks)

(a) (i) (1 mark)

Outcomes assessed: PE3**Targeted Performance Bands: E2-E3**

Criteria	Marks
• gives correct result (correct numerical equivalence)	1

Sample Answer:

$$(n-1)! = 9!$$

$$= 362880$$

(a) (ii) (2 marks)

Outcomes assessed: PE3**Targeted Performance Bands: E3-E4**

Criteria	Marks
• significant progress towards result	1
• gives correct result (correct numerical equivalence)	1

Sample Answer:

Number of arrangements without restrictions 9!

Number of arrangements if Gemma, Pasha and Ricky sit together is $3! \times 7!$

If Gemma, Pasha and Ricky sit separately then:

 $P(\text{all 3 separate}) = 1 - P(\text{all together})$

$$= 1 - \frac{3! \times 7!}{9!}$$

$$= 1 - \frac{6}{9 \times 8}$$

$$= \frac{11}{12}$$

(b) (2 marks)

Outcomes assessed: PE5, HE7**Targeted Performance Bands: E3-E4**

Criteria	Mark
• establishes correct differential relationship or progress toward result	1
• finds correct radius from working	1

Sample Answer

$$A = 4\pi r^2 \quad \Rightarrow \quad \frac{dA}{dr} = 8\pi r$$

$$\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$$

$$= 8\pi r \frac{dr}{dt} \quad \text{but} \quad \frac{dA}{dt} = \frac{dr}{dt}$$

$$\therefore 1 = 8\pi r$$

$$r = \frac{1}{8\pi} \text{ cm}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (3 marks)

Outcomes assessed: HE4

Targeted Performance Bands: E2-E3

Criteria	Marks
• uses appropriate substitution or other progress towards solution	1
• further progress towards solution	1
• establishes correct expression	1

Sample Answer:

$$\tan^{-1} x = \tan^{-1} y + \frac{\pi}{4} \Rightarrow \tan^{-1} x - \tan^{-1} y = \frac{\pi}{4}$$

$$\text{Let } \tan^{-1} x = A \text{ and } \tan^{-1} y = B \Rightarrow \text{ie } x = \tan A \text{ and } y = \tan B$$

$$\therefore A - B = \frac{\pi}{4}$$

Take the tan of both sides

$$\tan(A - B) = \tan\left(\frac{\pi}{4}\right)$$

$$\frac{\tan A - \tan B}{1 + \tan A \tan B} = \tan\left(\frac{\pi}{4}\right)$$

$$\frac{x - y}{1 + xy} = 1$$

$$x - y = 1 + xy$$

$$xy + y = x - 1$$

$$y(x + 1) = x - 1$$

$$y = \frac{x - 1}{x + 1}$$

(d) (i) (1 mark)

Outcomes assessed: HE3

Targeted Performance Bands: E2-E3

Criteria	Mark
• justifies the equation	1

Sample Answer:

$$T = 25 + 1315e^{-kt} \Rightarrow 1315e^{-kt} = T - 25$$

$$\frac{dT}{dt} = -1315ke^{-kt}$$

$$= -k(T - 25)$$

Also when $t = 0$, $T = 1340$ ie $1340 = 25 + 1315e^0$ which is true.

$\therefore T = 25 + 1315e^{-kt}$ satisfies this information

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(d) (ii) (3 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E3-E4

Criteria	Marks
• uses information to establish value for k or other progress towards solution	1
• further progress towards solution	1
• finds the correct time	1

Sample Answer:

$$\text{When } t = 12, \quad T = 1010 \quad \text{ie } 1010 = 25 + 1315e^{-12k}$$

$$985 = 1315e^{-12k}$$

$$\frac{985}{1315} = e^{-12k}$$

$$-12k = \ln\left(\frac{197}{263}\right)$$

$$k = \frac{-1}{12} \ln\left(\frac{197}{263}\right)$$

$$= 0.024079...$$

$$\text{When } T = 60; \quad 60 = 25 + 1315e^{-kt}$$

$$35 = 1315e^{-0.024t}$$

$$\frac{35}{1315} = e^{-0.024t}$$

$$-0.024t = \ln\left(\frac{7}{263}\right)$$

$$t = \frac{-1}{0.024} \ln\left(\frac{7}{263}\right)$$

$$= 150.5965769...$$

$$\therefore t = 151 \text{ minutes}$$

$$\text{OR } t = 151.09349... \text{ if using } k = 0.024$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 3 (12 marks)

(a) (2 marks)

Outcomes assessed: PE3**Targeted Performance Bands: E3-E4**

Criteria	Marks
• uses remainder theorem or other progress towards solution	1
• establishes correct conclusion	1

Sample Answer:

$$P(x) = (2x^2 + x + 3)Q(x) + (4x - 1)$$

$Q(x)$ has remainder 1 when divided by $(x + 2)$

ie $Q(-2) = 1$

$$\therefore P(-2) = (2 \times (-2)^2 + (-2) + 3) \times 1 + (4 \times (-2) - 1)$$

$$= (9) + (-9)$$


$$= 0$$

Since $P(-2) = 0$ by the Factor Theorem $(x + 2)$ is a factor of $P(x)$.

(b) (i) (1 mark)

Outcomes assessed: PE3**Targeted Performance Bands: E2-E3**

Criteria	Marks
• applies theorem correctly	1

Sample Answer:

AC is a tangent to the circle with diameter BP .

$\therefore \angle RPA = \angle RBP$ (angle between tangent and chord is equal to the angle in the alternate segment)

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (ii) (3 marks)

Outcomes assessed: PE3

Targeted Performance Bands: E2-E3

Criteria	Marks
• correctly identifies one pair of angles	1
• correctly identifies second pair of angles or other progress towards the proof	1
• completes the proof	1

Sample Answer:

In $\triangle BRP$ and $\triangle BPC$

$\angle BPC = 90^\circ$ (tangent is perpendicular to the radius drawn from the point of contact)

$\angle BRP = 90^\circ$ (angle in a semi-circle is a right angle)

$\therefore \angle BRP = \angle BPC$

$\angle RBP = \angle PBC$ (PB bisects $\angle RBC$ given $\triangle ABC$ is isosceles and $BP \perp AC$)

$\therefore \triangle BRP$ is similar to $\triangle BPC$ (equiangular)

(c) (i) (2 marks)

Outcomes assessed: PE3, PE4

Targeted Performance Bands: E2-E3

Criteria	Marks
• progress towards finding the coordinates	1
• derives correct coordinates	1

Sample Answer:

Equation of tangent at P is $y = px - ap^2$ and equation of tangent at Q is $y = qx - aq^2$

solve simultaneously $(p - q)x = a(p^2 - q^2)$

$$x = \frac{a(p - q)(p + q)}{(p - q)}$$

$$x = a(p + q)$$

Substitute for x : $y = p(a(p + q)) - ap^2$

$$y = apq$$

$\therefore T$ is $(a(p + q), apq)$

(c) (ii) (2 marks)

Outcomes assessed: PE3, PE4

Targeted Performance Bands: E3-E4

Criteria	Marks
• states the gradient of the normal	1
• equates gradients to show the result	1

Sample Answer:

The gradient of the tangent at P is $p \therefore$ gradient of normal is $-\frac{1}{p}$.

The gradient of the chord PQ is $\frac{p+q}{2} \therefore \frac{p+q}{2} = -\frac{1}{p}$

$$\text{ie } p + q + \frac{2}{p} = 0$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (iii) (2 marks)

Outcomes assessed: PE3, PE4

Targeted Performance Bands: E3-E4

Criteria	Marks
• progress towards result	1
• establishes the correct locus	1

Sample Answer:

At T $x = a(p + q)$, $y = apq$ and from (ii) $p + q = \frac{-2}{p}$

$$\therefore x = \frac{-2a}{p} \text{ ie } p = \frac{-2a}{x}$$

$$\text{also } q = \frac{x}{a} - p \text{ ie } q = \frac{x}{a} - \frac{-2a}{x} = \frac{x^2 + 2a^2}{ax}$$

$$y = a \times \frac{-2a}{x} \times \frac{x^2 + 2a^2}{ax} \\ = \frac{-2a^2x^2 - 4a^4}{ax^2}$$

$$\therefore y = \frac{-4a^3}{x^2} - 2a$$

Question 4 (12 marks)

(a) (3 marks)

Outcomes assessed: HE3, HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• progress towards solution	1
• further progress towards solution	1
• finds correct answer (correct numerical equivalence)	1

Sample Answer:

Total of 10 games – Harry wins 5 out of the first 9 and the last game

For the first 9 games consider the binomial probability of winning 5 from 9 with $p = \frac{2}{3}$

$$P(\text{winning 5}) = {}^9C_5 \left(\frac{1}{3}\right)^4 \left(\frac{2}{3}\right)^5 \\ = \frac{9!}{5!4!} \times \frac{1}{3^4} \times \frac{2^5}{3^5} \\ = \frac{9 \times 7 \times 2^6}{3^9} \\ = \frac{448}{2187}$$

Harry wins the last game \therefore probability of winning 6 games to 4 is $\frac{448}{2187} \times \frac{2}{3} = \frac{896}{6561}$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (i) (2 marks)


Outcomes assessed: PE2, HE7

Targeted Performance Bands: E2-E3

Criteria	Marks
• draws correct graph or other progress towards solution	1
• explains the conclusion	1

Sample Answer:

To solve $\sin^{-1} x + x - \frac{\pi}{2} = 0$ consider the graphs of $y = \sin^{-1} x$ and $y = -x + \frac{\pi}{2}$



There is only one point of intersection of the two graphs at a point where x is positive.

$\therefore \sin^{-1} x + x - \frac{\pi}{2} = 0$ has only one real positive root.

(b) (ii) (3 marks)

Outcomes assessed: PE3, HE7

Targeted Performance Bands: E2-E3

Criteria	Marks
• progress towards solution using Newton's Method	1
• further progress towards solution	1
• finds correct approximation (correct numerical equivalence)	1

Sample Answer:

$$f(x) = \sin^{-1} x + x - \frac{\pi}{2} \quad \therefore f'(x) = \frac{1}{\sqrt{1-x^2}} + 1$$

$$\text{For } x_1 = 0.7 \quad f(x_1) = \sin^{-1} 0.7 + 0.7 - \frac{\pi}{2} = -0.09539883...$$

$$f'(x_1) = \frac{1}{\sqrt{1-0.7^2}} + 1 = 2.40028008...$$

$$\begin{aligned} \therefore x_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} \\ &= 0.7 - \frac{-0.09539883...}{2.40028008...} \\ &= 0.73974... \\ &= 0.74 \end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (i) (2 marks)

Outcomes assessed: PE2, PE3

Targeted Performance Bands: E3-E4

Criteria	Marks
• establishes correct relationship	1
• finds correct values (correct numerical equivalence)	1

Sample Answer:

Series is geometric with $r = -\tan^2 x$, $-\frac{\pi}{2} < x < \frac{\pi}{2}$

For a limiting sum $|r| < 1$, ie consider $-1 < -\tan^2 x < 1$

If $-1 < -\tan^2 x < 1$ then $1 > \tan^2 x > -1$, ie $-1 < \tan^2 x < 1$

Since $\tan^2 x \geq 0$, solve $0 \leq \tan^2 x < 1$ for x

$\tan x$ is an increasing function for $-\frac{\pi}{2} < x < \frac{\pi}{2}$

and $\tan\left(-\frac{\pi}{4}\right) = -1$, $\tan(0) = 0$, $\tan\left(\frac{\pi}{4}\right) = 1$

ie for $-\frac{\pi}{4} < x < \frac{\pi}{4}$, $0 \leq \tan^2 x < 1$

\therefore for a limiting sum $-\frac{\pi}{4} < x < \frac{\pi}{4}$

(c) (ii) (2 marks)

Outcomes assessed: PE2, HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• applies correct formula	1
• correctly simplifies the expression	1

Sample Answer:

$$\begin{aligned} S_{\infty} &= \frac{a}{1-r} \\ &= \frac{\tan^2 x}{1 - (-\tan^2 x)} \\ &= \frac{\tan^2 x}{1 + \tan^2 x} \\ &= \frac{\tan^2 x}{\sec^2 x} \\ &= \frac{\sin^2 x}{\cos^2 x} \times \cos^2 x \\ &= \sin^2 x \end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(a) (i) (1 mark)


Outcomes assessed: PE2, HE7

Targeted Performance Bands: E2-E3

Criteria	Marks
• derives correct result	1

Sample Answer:

$$\text{In } \triangle ACE, \tan \beta = \frac{2h}{AC} \Rightarrow AC = 2h \cot \beta$$



(a) (ii) (2 marks)

Outcomes assessed: PE2, HE7

Targeted Performance Bands: E2-E3

Criteria	Marks
• derives correct result for AB	1
• derives correct result for BC	1


Sample Answer:

$$\text{In } \triangle ABD, \tan \beta = \frac{h}{AB} \Rightarrow AB = h \cot \beta$$

$$\text{Also } BC = DF$$

$$\text{In } \triangle DEF, \tan \alpha = \frac{h}{DF} \Rightarrow DF = h \cot \alpha$$

$$\therefore BC = h \cot \alpha$$



(a) (iii) (2 marks)

Outcomes assessed: PE2, HE7


Targeted Performance Bands: E3-E4

Criteria	Marks
• applies the Cosine Rule to correct triangle	1
• shows correct result	1

Sample Answer:

In $\triangle ABC$

$$\begin{aligned} \cos \theta &= \frac{AC^2 + AB^2 - BC^2}{2AC \times AB} \\ &= \frac{4h^2 \cot^2 \beta + h^2 \cot^2 \beta - h^2 \cot^2 \alpha}{4h^2 \cot^2 \beta} \\ &= \frac{5h^2 \cot^2 \beta - h^2 \cot^2 \alpha}{4h^2 \cot^2 \beta} \\ &= \frac{h^2 (5 \cot^2 \beta - \cot^2 \alpha)}{4h^2 \cot^2 \beta} \\ &= \frac{(5 \cot^2 \beta - \cot^2 \alpha)}{4 \cot^2 \beta} \end{aligned}$$



DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (3 marks)

Outcomes assessed: PE2, HE3

Targeted Performance Bands: E3-E4

Criteria	Marks
• establishes correct terms or coefficients for comparison or other progress towards result	1
• significant progress toward the result	1
• finds correct values (correct numerical equivalence)	1

Sample Answer:

Consider the 6th, 7th and 8th terms of the expansion of $(2 + bx)^{11}$

$$T_6 = {}^{11}C_5 \times 2^6 \times (bx)^5$$

$$T_7 = {}^{11}C_6 \times 2^5 \times (bx)^6$$

$$T_8 = {}^{11}C_7 \times 2^4 \times (bx)^7$$

Take coefficients of T_6 and T_8 , and compare to T_7

Consider $T_7 > T_6$ ie ${}^{11}C_6 \times 2^5 \times b^6 > {}^{11}C_5 \times 2^6 \times b^5$

$$\therefore b > \frac{{}^{11}C_5 \times 2}{{}^{11}C_6} = 2$$

Similarly for $T_7 > T_8$ ie ${}^{11}C_6 \times 2^5 \times b^6 > {}^{11}C_7 \times 2^4 \times b^7$

$$\therefore b < \frac{{}^{11}C_6 \times 2}{{}^{11}C_7} = 2.8$$

\therefore seventh term has the largest coefficient for $2 < b < 2.8$

(c) (i) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E2-E3

Criteria	Mark
• uses correct formula or progress using other correct method	1
• establishes correct result	1

Sample Answer:

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = -n^2 x$$

$$\frac{1}{2} v^2 = \frac{-n^2 x^2}{2} + C$$

at $x = a$, $v = 0$ since velocity is zero at the extremities

$$0 = \frac{-n^2 a^2}{2} + C \Rightarrow C = \frac{n^2 a^2}{2}$$

$$\frac{1}{2} v^2 = \frac{-n^2 x^2}{2} + \frac{n^2 a^2}{2} \quad \text{ie } v^2 = n^2 a^2 - n^2 x^2$$

$$\therefore v^2 = n^2 (a^2 - x^2)$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (ii) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E3-E4

Criteria	Marks
• uses correct substitutions or other progress towards result	1
• finds correct values	1

Sample Answer:

$$v = 6 \text{ when } x = 4 \Rightarrow 36 = n^2(a^2 - 16)$$

maximum velocity is at the centre of the motion

$$\therefore v = 10 \text{ when } x = 0 \Rightarrow 100 = n^2 a^2$$

$$\text{solving simultaneously } \Rightarrow 36 = 100 - 16n^2$$

$$16n^2 = 64$$

$$n^2 = 4$$

$$\text{hence } a^2 = 25$$

\therefore extremities of motion are $a = 5$ and $a = -5$

Question 6 (12 marks)

(a) (i) (1 mark)

Outcomes assessed: HE7

Targeted Performance Bands: E2-E3

Criteria	Marks
• correctly justifies the result	1

Sample Answer:

$${}^nC_k = \frac{n!}{k!(n-k)!}$$

$$\begin{aligned} {}^nC_{n-k} &= \frac{n!}{(n-k)!(n-(n-k))!} \\ &= \frac{n!}{(n-k)!k!} \end{aligned}$$

$$\therefore {}^nC_k = {}^nC_{n-k}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(a) (ii) (2 marks)

Outcomes assessed: HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• consider terms in binomial expansion or other progress towards solution	1
• establishes the result	1

Sample Answer:

Consider terms in the expansion of $(1+x)^{2n}$

$$\begin{aligned} T_{k+1} &= {}^{2n}C_k 1^{2n-k} x^k \\ &= {}^{2n}C_k x^k \end{aligned}$$

$$\begin{aligned} \text{coefficient of } x^n \text{ is: } {}^{2n}C_n &= \frac{(2n)!}{n!(2n-n)!} \\ &= \frac{(2n)!}{(n!)^2} \end{aligned}$$

Consider the expansion of $(1+x)^n (1+x)^n$

$$(1+x)^n (1+x)^n = \left[{}^nC_0 + {}^nC_1 x + {}^nC_2 x^2 + \dots + {}^nC_{n-1} x^{n-1} + {}^nC_n x^n \right] \times \left[{}^nC_0 + {}^nC_1 x + {}^nC_2 x^2 + \dots + {}^nC_{n-1} x^{n-1} + {}^nC_n x^n \right]$$

coefficients of x^n are: ${}^nC_0 {}^nC_n + {}^nC_1 {}^nC_{n-1} + {}^nC_2 {}^nC_{n-2} + \dots + {}^nC_{n-1} {}^nC_1 + {}^nC_n {}^nC_0$

$$= ({}^nC_0)^2 + ({}^nC_1)^2 + ({}^nC_2)^2 + \dots + ({}^nC_{n-1})^2 + ({}^nC_n)^2 \quad \text{using (i)}$$

$$= \sum_{k=0}^n ({}^nC_k)^2$$

$$\therefore \text{Since } (1+x)^n (1+x)^n = (1+x)^{2n} \text{ equating coefficients gives } \sum_{k=0}^n ({}^nC_k)^2 = \frac{(2n)!}{(n!)^2}$$

(b) (i) (1 mark)

Outcomes assessed: HE4

Targeted Performance Bands: E3-E4

Criteria	Marks
• differentiates or other method to explain correct conclusion	1

Sample Answer:

$$f(x) = x^3 + x + 1$$

$$f'(x) = 3x^2 + 1 > 0 \text{ for all } x \text{ since } x^2 \geq 0$$

$\therefore f(x)$ is monotonic increasing and thus has an inverse function, $f^{-1}(x)$, for all x .

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (ii) (2 marks)

Outcomes assessed: HE4

Targeted Performance Bands: E3-E4

Criteria	Marks
• identifies that curves intersect on $y = x$ or other progress towards result	1
• finds correct point of intersection	1

Sample Answer:

$f(x)$ and $f^{-1}(x)$ intersect on $y = x$

\therefore solve $x^3 + x + 1 = x$

$$x^3 + 1 = 0$$

$$x^3 = -1$$

$$x = -1$$

\therefore Point of intersection is $(-1, -1)$

(c) (i) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E2-E3

Criteria	Marks
• progress towards result	1
• shows correct result	1

Sample Answer:

$$x = vt \cos \theta$$

$$\therefore t = \frac{x}{v \cos \theta}$$

substitute into $y = vt \sin \theta - \frac{1}{2}gt^2$

$$y = \frac{x}{v \cos \theta} (v \sin \theta) - \frac{1}{2}g \left(\frac{x}{v \cos \theta} \right)^2$$

$$= x \tan \theta - \frac{gx^2}{2v^2 \cos^2 \theta}$$

$$= x \tan \theta - \frac{gx^2}{2v^2} \sec^2 \theta$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (ii) (2 marks)

Outcomes assessed: HE3

Targeted Performance Bands: E3-E4

Criteria	Marks
• progress toward solution	1
• substitutes and simplifies to obtain desired result	1

Sample Answer:

At the point P , $x = 10$, $y = 15$ and given $g = 9.8$, $v = 7\sqrt{10}$

$$\begin{aligned}\text{Using (i)} \quad 15 &= 10 \times \tan \theta - \frac{9.8(10)^2}{2(7\sqrt{10})^2} \sec^2 \theta \\ 15 &= 10 \tan \theta - \frac{9.8 \times 100}{2 \times 49 \times 10} (1 + \tan^2 \theta) \\ 15 &= 10 \tan \theta - 1 - \tan^2 \theta \\ \tan^2 \theta - 10 \tan \theta + 16 &= 0 \\ (\tan \theta - 8)(\tan \theta - 2) &= 0 \\ \therefore \tan \theta &= 8 \text{ or } \tan \theta = 2 \\ \text{since } \alpha < \beta, \tan \beta &= 8 \text{ and } \tan \alpha = 2\end{aligned}$$

(c) (iii) (2 marks)

Outcomes assessed: HE3, HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• significant progress towards solutions	1
• shows correct solution	1

Sample Answer:

Consider the two paths and find time travelled to reach P

Pebble 1: $v = 7\sqrt{10}$, $\theta = \beta$, $\tan \beta = 8$ and $x = 10$

$$\begin{aligned}t_1 &= \frac{10}{7\sqrt{10} \cos \beta} = \frac{10 \sec \beta}{7\sqrt{10}} \text{ and } \sec^2 \beta = 1 + \tan^2 \beta = 65 \\ \therefore t_1 &= \frac{10\sqrt{65}}{7\sqrt{10}} \\ &= \frac{\sqrt{650}}{7}\end{aligned}$$

Pebble 2: $v = 7\sqrt{10}$, $\theta = \alpha$, $\tan \alpha = 2$ and $x = 10$

$$\begin{aligned}t_2 &= \frac{10}{7\sqrt{10} \cos \alpha} = \frac{10 \sec \alpha}{7\sqrt{10}} \text{ and } \sec^2 \alpha = 1 + \tan^2 \alpha = 5 \\ \therefore t_2 &= \frac{\sqrt{50}}{7} \\ \therefore t_1 - t_2 &= \frac{\sqrt{650} - \sqrt{50}}{7} \text{ seconds}\end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 7 (12 marks)

(a) (3 marks)

Outcomes assessed: HE2**Targeted Performance Bands: E3-E4**

Criteria	Marks
• establishes the truth of $S(2)$	1
• establishes the correct relationship between $S(k)$ and $S(k+1)$	1
• deduces the required result	1

Sample Answer:

Let $S(n)$ be the statement $2n^2 > n^2 + n + 1$ for $n > 1$

Consider $S(2)$: $2 \times 2^2 = 8$ and $2^2 + 2 + 1 = 7$

$\therefore 2n^2 > n^2 + n + 1$ for $n = 2$ and hence $S(2)$ is true

Assume $S(k)$ is true: $2k^2 > k^2 + k + 1$ *

RTP: $S(k+1)$ is true, ie prove $2(k+1)^2 > (k+1)^2 + (k+1) + 1$

$$2(k+1)^2 = 2k^2 + 4k + 2$$

$$> k^2 + k + 1 + 4k + 2 \quad \text{if } S(k) \text{ is true using } *$$

$$= k^2 + 2k + 1 + 3k + 2$$

$$= (k+1)^2 + (k+1) + 1 + 2k$$

$$\therefore 2(k+1)^2 > (k+1)^2 + (k+1) + 1 \text{ since } k > 0$$

Hence if $S(k)$ is true then $S(k+1)$ is also true. Thus since $S(2)$ is true it follows by induction that $S(n)$ is true for positive integers $n > 1$.

(b) (i) (2 marks)

Outcomes assessed: PE2, HE7**Targeted Performance Bands: E3-E4**

Criteria	Marks
• substitutes correctly	1
• shows correct result	1

Sample Answer:

$$\cosh x = \frac{1}{2}(e^x + e^{-x}) \text{ and } \sinh x = \frac{1}{2}(e^x - e^{-x})$$

$$\text{LHS} = 2 \sinh x \cosh x$$

$$= 2 \times \frac{1}{2}(e^x + e^{-x}) \times \frac{1}{2}(e^x - e^{-x})$$

$$= \frac{1}{2}((e^x)^2 - (e^{-x})^2)$$

$$= \frac{1}{2}(e^{2x} - e^{-2x})$$

$$= \sinh(2x)$$

$$= \text{RHS}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (ii) (2 marks)

Outcomes assessed: PE2, HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• establishes correct equation using given substitutions	1
• shows correct result	1

Sample Answer:

$$p \cosh x + q \sinh x = r \text{ and } \cosh x = \frac{1}{2}(e^x + e^{-x}) \text{ and } \sinh x = \frac{1}{2}(e^x - e^{-x})$$

$$p \times \frac{1}{2}(e^x + e^{-x}) + q \times \frac{1}{2}(e^x - e^{-x}) = r$$

$$\frac{pe^x}{2} + \frac{pe^{-x}}{2} + \frac{qe^x}{2} - \frac{qe^{-x}}{2} = r$$

$$\frac{e^x}{2}(p+q) + \frac{1}{2e^x}(p-q) = r$$

$$e^{2x}(p+q) + (p-q) = 2re^x$$

$$(p+q)e^{2x} - 2re^x + (p-q) = 0$$

(b) (iii) (3 marks)

Outcomes assessed: PE2, HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• recognises that the equation is a quadratic or other progress towards the solution	1
• uses the discriminant or other progress towards the solution	1
• establishes correct conclusion	1

Sample Answer:

From (ii) the equation $p \cosh x + q \sinh x = r$ is equivalent to

$$(p+q)e^{2x} - 2re^x + (p-q) = 0, \text{ which is a quadratic in } e^x$$

$$\Delta = 4r^2 - 4(p+q)(p-q)$$

$$= 4r^2 - 4(p^2 - q^2)$$

$$= 4(r^2 - p^2 + q^2)$$

$$= 0 \text{ since } p^2 = q^2 + r^2$$

\therefore the equation has only one solution

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (iv) (2 marks)

Outcomes assessed: PE2, HE7

Targeted Performance Bands: E3-E4

Criteria	Marks
• establishes the correct equation or other progress towards solution	1
• finds the correct solution	1

Sample Answer:

For the equation $13\cosh x + 5\sinh x = 12 \Rightarrow p = 13, q = 5, r = 12$

$$\therefore (p+q)e^{2x} - 2re^x + (p-q) = 0 \text{ becomes } 18e^{2x} - 24e^x + 8 = 0$$

$$\text{ie solve } 9e^{2x} - 12e^x + 4 = 0$$

$$(3e^x - 2)^2 = 0$$

$$3e^x = 2$$

$$e^x = \frac{2}{3}$$

$$\therefore x = \ln\left(\frac{2}{3}\right)$$

OR

Let $e^{2x} = y$ ie solve $9y^2 - 12y + 4 = 0$

$$(3y - 2)^2 = 0$$

$$y = \frac{2}{3} \Rightarrow e^x = \frac{2}{3}$$

$$\therefore x = \ln\left(\frac{2}{3}\right)$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

BLANK PAGE

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 2 (12 marks) Use a SEPARATE writing booklet.


- (a) Ten students are seated around a circular table.
- (i) In how many ways can they be arranged? 1
- (ii) What is the probability that three particular students, Gemma, Pasha and Ricky, are not sitting together when the seats are randomly assigned. 2
- (b) A ball in the shape of a sphere has radius r centimetres at time t seconds. The surface area is changing as the radius changes over time. At a particular time, t seconds, the rate of change of the surface area is equal to the rate of change of the radius. 2
- Find the exact radius at this time.
- (c) Find an expression for y in terms of x if for $x > 0$ and $y > 0$, 3
- $$\tan^{-1} x = \tan^{-1} y + \frac{\pi}{4}.$$
- (d) A heated metal bar has a temperature of 1340°C when it is removed from a furnace. Its temperature T after t minutes in a room with a constant temperature of 25°C satisfies the equation $\frac{dT}{dt} = -k(T - 25)$, where k is a constant.
- (i) Show that the equation $T = 25 + 1315e^{-kt}$ satisfies this information. 1
- (ii) The metal bar cools to 1010°C after 12 minutes. 3
- Find how long it will take for the bar to cool to 60°C , giving your answer correct to the nearest minute.

Question 3 (12 marks) Use a SEPARATE writing booklet.

- (a) The polynomial $P(x)$ is expressed as $P(x) = (2x^2 + x + 3)Q(x) + (4x - 1)$. **2**

If $Q(x)$ leaves a remainder of 1 when divided by $(x + 2)$, show that $(x + 2)$ is a factor of $P(x)$.

- (b) The diagram shows an isosceles triangle ABC , with $AB = BC$. The point P lies on AC and the point O lies on BP . A circle with centre O passes through B and P and cuts AB at R .



Copy or trace the diagram into your writing booklet.

- (i) Explain why $\angle RPA = \angle RBP$. **1**
- (ii) Hence, or otherwise, prove that $\triangle BRP$ is similar to $\triangle BPC$. **3**

Question 3 continues on page 5