SECTION I - CORE

PART A (15 marks)

Each question is worth 1 mark. Attempt all questions.

Select the alternative A, B, C or D that best answers the question. Mark your answers in pencil on the Answer Sheet provided.

- Emission from radioisotopes can be detected by a --
- A. pH meter
 B. thermomete
 C. smoke dete
 D. photograph
- thermometer smoke detector
- photographic film
- PVC (polyvinyl chloride) is an addition polymer made from vinyl chloride (chloroethene) ٨i

The most likely structure of a section of PVC is

ū

Student Number:

HIGHER SCHOOL CERTIFICATE Sample Examination Paper

CHEMISTRY

General Instructions

Reading time - 5 minutes Working time - 3 hours

- · Board approved calculators may be
- Write your Student Number on each Answer Book.
- provided in the Answer Books. Anything written in the Question Book will NOT be marked. Answer all questions in the spaces

Section I - Core

Attempt ALL questions.

Answer this part in the Part B Mark your answers in pencil on the Answer Sheet provided.

• Part B Other questions with a total mark value of 60.

Part A 15 multiple -choice questions,

each worth 1 mark

part of the Answer Book.

- Attempt ONE question only.
 Each question is worth 25 marks. Section II - Electives
 - Answer the question in a separate
- · You may ask for an extra Elective Elective Answer Booklet.
 - Answer Book if you need one.
- provided at the back of this paper. A Data Sheet and Periodic Table are

Directions to School or College

To ensure auxiliams confidentially and security, estantiation papers must NOT be removed from the examination room. Examination papers may and be retained to students or used for revision purposes till Spiceaber 2001. These examination papers are applied Copyright Free, as such the parchaner may photocopy and/or cut and paste name for educational purposes while the examinas of the action of oreflect.

All care has been taken to ensure that this canaliantion paper is error free and that it follows the style, format and material content of the sew Higher School Cerifficies pitabon. Candidates are advised that authors of this examination paper cannot any way generate that the 2001 HSC Estanisation will have a similar content or format.

- Which of the following is a dilute strong acid?
- A. 5.00 mol L⁻¹ HCl B. 5.00 mol L⁻¹ CH₂COOH C. 0.05 mol L⁻¹ HCl D. 0.05 mol L⁻¹ CH₂COOH

- The table shows the pH of some naturally occurring substances:

		,	
substance	approximate pH	substance	approximate pH
stomach acid	2	tornato juice	4
lemon juice	3	sea water	œ

next table shows the properties of four acid-base indicators

THE THE STATE STAT	or and properties of	in the more arrows are properties of four activities indicators.	Carolia.
Name	Colour in		p.H. range in which
	Hq wol	th pH	colour changes
phenolphthalein	colouriess	pink	8.3 - 10.0
phenol red	yellow	1 00	6.8 - 8.4
methyl red	pink	yellow	4.4 - 6.2
methyl vellow	red	vellow	2.4 – 4.0


Which mixture has the correct colour next to it?

- A. Iemon juice + phenolphthalein → pink
 B. saliva + phenol red → yellow
 C. sea water + methyl red → pink
 D. stomach acid + methyl yellow → yellow
- stomach acid + methyl yellow → yellow
- The pHs of 0.2 molar solutions of citric acid, acetic acid and hydrochloric acid are 2.0, 2.7 and 0.7 respectively. œ

This means that the hydrogen ion concentration of

- A. hydrochloric acid is about 100 times that of acetic acid. B. acetic acid is about 1/3 more than that of citric acid. C. citric acid is about 3 times that of hydrochloric acid. D. acetic acid is about ½ that of hydrochloric acid.
- In preparing a 0.1 mol L-1 primary standard NaHCO, solution, ο.
- solid NaHCO, should be dried in the oven before being weighed. the volumetric flask should be rinsed with some 0.1 mol L $^{-1}$ NaHCO, 100 mL of 0.100 mol L $^{-1}$ solution should be made up to exactly 1.000 L. A. 0.100 moles of NaHCO, should be added to 1.000 litre of water. B. solid NaHCO, should be dried in the oven before being weighed. C. the volumetric flask should be rinsed with some 0.1 mol L⁻¹ NaH. D. 100 mL of 0.100 mol L⁻¹ solution should be made up to exactly 1

- The process of catalytic cracking
- changes a long chain alkane to two short chain alkanes.
 - involves reactions on the surface of inorganic catalysts.
- cracks the solid catalyst to fragments to increase its surface area is used to convert six-carbon glucose to carbon dioxide and ethanol. 4 E U D
- When glucose polymerises to form cellulose, the other product formed is 4
- water
- ethanol
- methane 水色じ口
- carbon dioxide



In the above galvanic cell electrons flow in the wire from one metal to the other, and anions and cations travel in the salt bridge.

Which set of arrows correctly shows the direction of the movement of the electrons (solid arrow) in the wire and the direction of the movement of the anions (broken arrow) in the salt bridge?

- 1 Ŷ ተ
- 11 4ほじひ

15. In which of the following is the coordinate covalent bond correctly indicated by the arrows?

10. HCO3 is an amphiprotic species as shown by the equations

$$HCO_{3}^{-}(\omega_{0} + HCl_{(\omega_{0})} \rightarrow H_{2}CO_{3(\omega_{0})} + Cl_{-}(\omega_{0})$$

 $HCO_{3}^{-}(\omega_{0}) + OH_{-}(\omega_{0}) \rightarrow CO_{3}^{-}(\omega_{0}) + H_{2}O_{(1)}$

The conjugate base of HCO3 [44] is

- A. OH B. H;CO; C. H;O D. CO;

- 11. The products of the combustion of petrol (mainly octane) are determined partly by the fuel to air ratio and partly by the homogeneity (evenness) of the fuel-air mixture. These products, apart from water, rang contribute to global warming, smoky air and a reduction of the oxygen supply to body cells.

Which of the following reactions would cause the greatest amount of smoke?

A.
$$C_6H_{18} + 7O_7 \Rightarrow 9H_7O + 4C + CO_7 + 3CO$$

B. $C_6H_{18} + 9O_7 \Rightarrow 9H_7O + 3CO + 2C + 3CO_7$
C. $C_6H_{18} + 7O_7 \Rightarrow 9H_2O + 5CO + 3C$
D. $C_6H_{18} + 9O_7 \Rightarrow 9H_7O + CO_7 + 7CO$


- 12. Which solution is used to detect the presence of chloride ions?
- A. H₂SO₄ B. Ba(OH)₂ C. AgNO₃ D. NH₃
- 13. A chief cause of algal bloom in waterways is an increase in the water's

- A. acidity
 B. hardness
 C. temperature
 D. phosphate concentration
- II. 2,4-dichlorohexane IV. 2,2,3,3-tetrachloropentane 14. I. 2,4-dichloropentane III 2,3-dichlorohexane

Which two of the above chloroalkanes are isomers?

- A. I and II
 B. I and IV
 C. II and III
 D. III and IV

20. The diagram shows the set-up for the electrolytic refining of copper.



a. Name a suitable electrolyte.

Write the half equation for the reaction which occurs where this electrolyte is in contact with the pure copper.

The impure copper acts as the anode in this electrolytic cell. What process takes place at the anode of any cell? ڻ

21. In water there is an equilibrium between gaseous and dissolved carbon dioxide, according to the equation

The dissolving process is exothermic.

A further series of equilibrium reactions occurs as the dissolved carbon dioxide reacts with water.

$$CO_{Meg} + H_2O_m = H_2CO_{Heg}$$
 $H_2CO_{Meg} + H_2O_m = H_3O^+_{leg} + HCO_3^-_{leg}$
 $HCO_3^-_{leg} + H_2O_m = H_3O^+_{leg} + CO_3^{-2}_{leg}$

Explain in terms of Le Chatelier's Principle why

- a fizzing occurs when a bottle of soft drink is opened?
- the fizzing is less if the bottle of soft drink was kept in the refrigerator than if it was kept at room temperature.
- c. a "flat" soft drink (i.e. from which all the bubbles escaped) has a higher pH than a newly opened bottle.

PART B (60 marks)

Attempt all questions.

The value of each question is indicated next to the question.

In questions requiring calculations show all working.

Put the appropriate chemical symbol, mass number or atomic number into the boxes to represent a balanced nuclear reaction producing a transuranic element. <u>1</u>9

Mark

-{	‡ *
Н	۲
1	•
ູເ	<u> </u>
₽ +	\sqcup
8	8

17. Name a radioisotope used in medicine, and describe what it is used for.

Biopolymers are naturally occurring polymers.

a. Give an example of a biopolymer.

Give two reason for the need to replace polymers made from petroleum products with biopolymers.

19. a. Write equations for the

i. production of ethanol by fermentation.

ii. dehydration of ethanol.

In both cases indicate the catalyst used

b. Explain why ethanol is a fairly good solvent for most water soluble compounds. 2

c. The table lists some of the properties of octane (the main component of petrol) and of ethanol (which can be used as an alternative to car fuel).

fuel	molar mass (g)	mass of 1 litre (g)	heat released during combustion (kJ mol ⁻¹)
cthanol	46.1	785	1367
octane	114.2	869	5470

Assuming all other variables to be equal, 1 litre of which firel would allow a car to travel further? (Calculate the energy obtained by the complete combustion of 1 litre of each of these fuels.)

- 25. a. Explain in words and with equations why
- i. HCl is an Arrhenius acid when it dissolves in water.
- ii. NH4Cl is a Bronsted-Lowry acid when it reacts with NaOH.
- b. Draw Lewis diagrams for each of the substances in the following equation BF₃ + NH₃ → BF₃NH₃

Indicate which species is the Lewis acid.

26. The synthesis of ammonia gas is represented by the equation.

$$N_{2(g)} + 3H_{2(g)} = 2NH_{3(g)}$$

- Explain why the industrial process uses high pressures but moderate (not too high and not too low) temperatures. æ
- A catalyst allows this same reaction to proceed at a lower temperature.
 - i. Name the catalyst which is used in the Haber process of ammonia manufacture.
- ii. Explain why the use of the catalyst can lower the temperature that is needed for this reaction.
- 27. The nitrogen content of a fertiliser was determined by heating 0.95 g of the finely ground fertiliser with 100.0 mL of 0.15 mol L⁻¹ of sodium hydroxide solution, known to be in excess. This converted all the ammonium ions in the fertiliser to ammonia gas, which left the mixture.

$$NH_4^+$$
 (eq. + $OH_{(eq)} \rightarrow NH_{3(g)} + H_2O_{(f)}$

The sodium hydroxide left over after the reaction stopped needed 23.1 mL of 0.23 mol L^{-1} hydrochloric acid for neutralisation.

$$NaOH_{(eq)} + HCl_{(eq)} \rightarrow H_2O_{(f)} + NaCl_{(eq)}$$

How many moles of sodium hydroxide reacted with the ammonium ions?

3

- How many moles of nitrogen were in 0.95 g of fertiliser? (Assume that all the nitrogen in the fertiliser is present as ammonium ions.) نم
- c. What was the percentage of nitrogen, by mass, in the fertiliser?

22. One of the half equations which occurs in a vanadium cell is

$$2H^7 + VO_2^4 + \epsilon^7 \rightarrow VO^{2+} + H_2O$$

- a. Is this an oxidation or a reduction half-equation? Justify your answer.
- What is the oxidation state of vanadium in VO2+?
- 23. a. To extract metal from sulfide ores, the ore is "roasted" in air. A by-product of the reaction is the poisonous gas sulfur dioxide.

Write the equation for the roasting of the ore copper pyrites (copper (II) sulfide). 1

- Calculate the volume of suffur dioxide gas produced (measured at 101.3 kPa and 25°C) from the roasting of 1 tonne (= 1000 kg = 10^6 g) of copper pyrites. ئع
- If sulfur dioxide is not removed, it combines with moisture in the air, producing ن

Write an equation for the reaction between sulfur dioxide and water.

The acid in part (c) can further oxidise, giving acid rain.

Write an equation between acid rain (assume it to be sulfuric acid) and marble statues (assume marble to be calcium carbonate).

- a. 0.1 mole of proparoic acid was refluxed with 0.1 mole of ethanol and a small amount (1 mL) of concentrated sulfuric acid catalyst.
- i. Complete the equation for the reaction

CH3CH2CH2COOH0+CH3CH2OH0

ii. Name the organic product.

- b. Although this is an equilibrium reaction, it was forced to go to completion, so there was no propanoic acid left. To find the amount of sulfuric acid catalyst used, the resultant mixture was titrated with a 0.5 mol L⁻¹ sodium hydroxide
- i. What was the pH of the sodium hydroxide solution?
- ii. It required 68 mL of the sodium hydroxide solution to reach the end point of

$$H_2SO_{4(eq)} + 2NaOH_{(eq)} \rightarrow 2H_2O_{(f)} + Na_2SO_{4(eq)}$$

How many moles of sulfuric acid were used?

SECTION II — OPTIONS

(25 marks)

Candidates should attempt ONE option from this section.

Answer the question in a separate Elective Answer Book.

In questions involving calculations, you are advised to show working, as marks may be awarded for relevant working.


OPTION 1	INDUSTRIAL CHEMISTRY13
OPTION 2.	SHIPWRECKS AND SALVAGE17
OPTION 3.	BIOCHEMISTRY OF MOVEMENT20
OPTION 4.	CHEMISTRY OF ART
OPTION 5.	FORENSIC CHEMISTRY 24

- 29. Explain why it is dangerous to
- a. decrease the concentration of ozone in the upper atmosphere (stratosphere).
 - b. increase the concentration of ozone in the lower atmosphere (troposphere).
- 30. The river water downstream from a factory was suspected of having lead ion and barium ion contamination.
- a. Analysis of a sample of this water by atomic absorption spectroscopy found its absorbance value at the main lead wavelength to be 0.078. The absorbance value of a standard laboratory sample, known to have a concentration of 5.85 ppm (parts per million) of lead, was 1.087.
- Calculate the concentration of lead, in ppm, in the river water sample. (Note: absorbance is proportional to concentration.)
- b. A 100.0 mL sample of the river water was mixed with 10.0 mL of 2.05 mol L⁻¹ sulfuric acid. The precipitate that formed was filtered, dried and weighed. Its mass was 0.27 g.

Ba²⁺ (rep + SO₄ 2- (req) → BaSO₄₍₆₎

- i. How many moles of barium sulfate precipitated?
- ii. How can you be sure that there was enough sulfuric acid used?
- iii. What was the mass of barium (ions) per litre of river water?

4. In which container will the rate of electrolysis of the same electrolyte be the fastest?



Key. [52] 0.1 mol L⁻¹ ■ 1.0 mol L⁻¹ a. Explain with the help of a labelled diagram how the hull of a steel ship is protected by a "sacrificial anode" (cathodic protection).

. Apart from ships, where is cathodic protection used?

c. i. Identify another way of preventing rust formation.

ii. Explain how this method prevents rusting

6. a. What are anacrobic bacteria?

How do anaerobic bacteria speed up the corrosion of deep sea shipwrecks?

7. Sketch graphs to show the relationship between

a. the solubility of salts as a function of water temperature.

the solubility of gases as a function of water temperature.

the solubility of gases as a function of water depth.

82

OPTION 2. SHIPWRECKS AND SALVAGE

Marks

Identify the people associated with the following:

 The generation of electricity by creating a circuit of two dissimilar metals and a fresh frog muscle. The first scientific use of electricity to decompose compounds. (The first person to prepare samples of potassium and sodium.)

 c. The discovery that the mass of a substance formed at an electrode during electrolysis is proportional to the quantity of electricity that passes through the electrolytic cell.

 Although aluminium is a much more reactive metal than iron, aluminium objects do not deteriorate readily.

Explain why aluminium objects resist deterioration.

Below are the reduction potentials of some standard half cells.

	E (V)
Mg' + 2e - Mg(s)	-2.35
2H ₂ O ₀₀ + 2e ⁻ ⇒ H _{2(g)} + 2OH	-0.83
$Pb^{2+} + 2e^- + Pb_{(i)}$	-0.13
Cu ²⁺ + 2e = Cu ₍₀₎	0.34
l _{2(mq)} + 2e ⁻ → 2T	0.54
$O_{2(g)} + 4H^{\dagger} + 4e^{-} = 2H_2O_{(f)}$	1.23
Cl _{2(eq)} + 2e ⁻ = 2Cl ⁻	1.36

a. What is the voltage of a standard Cu/Cu²⁺ || Pb²⁺/Pb cell?

b. What will be discharged initially at the anode and at the cathode when 0.1 mol L⁻¹ solutions of the following compounds are electrolysed?

i. copper (II) iodide

ii. magnesium chloride

OPTION 3. BIOCHEMISTRY OF MOVEMENT

a. Which cell organelle is the site of aerobic respiration? _:

Marks

b. What is the function of the ATP produced?

c. The diagram shows a shorthand notation for adenosine monophosphate.

Complete the copy of this diagram in your answer book changing it into ATP.

a. The structural formula of glycerol is 7

Give its systematic name.

Explain glycerol's ض

high viscosity miscibility with water. .**.**: : ::

c. The structure of triacylglycerol (TAG) which may be found in coconut oil is

Why is coconut oil insoluble in water?

CH₃(CH₂)₁₂COOCH₂

This molecule is a special type of ester. Write the formula of the functional group which qualifies it to be called an ester. ... :≓

Why are TAGs important to humans?

8

~

Why is the simple drying of soaked leather and wood artefacts recovered from shipwrecks a bad practice? œ

 Write the neutral species equation for the reaction between silver coins (sunk with a ship) and hydrogen sulfide (produced by anaerobic bacteria). ف

Write the reduction half equation for the restoration of these silver coins by electrolysis, using the silver sulfide covered coins (one at a time) as the cathode.

iii. Write the ionic equation for the process by which limestone and coral encrusted copper artefacts are chemically cleaned.