

Sydney Boys High School

4 unit mathematics

Trial HSC Examination 1994


1. (a) Find $\int \frac{2x}{1+x^4} dx$
(b) Evaluate
(i) $\int_1^2 \frac{dx}{x^2-x+1}$
(ii) $\int_0^{\frac{\pi}{2}} \frac{dx}{2+\cos x}$
(iii) $\int_0^1 e^{\sqrt{x}} dx$
(c) Prove that $\int_a^b f(mx) dx = \frac{1}{m} \int_{ma}^{mb} f(x) dx \quad (m \neq 0)$
2. (a) (i) If $z = 3 - 4i$, express z^2 and $\frac{1}{z}$ in the form $a + ib$ where a and b are real numbers. Represent them on an Argand diagram.
(ii) If $w^2 = z$, express the two values of w in the form $a + ib$.
(b) Illustrate on the Argand diagram the region $\{z : 0 \leq \arg(z+4) \leq \frac{2\pi}{3} \wedge |z+4| \leq 4\}$
(c) Let z_1, z_2 and z_3 be three complex numbers represented by Z_1, Z_2 and Z_3 respectively where $z_1 \times z_3 = (z_2)^2$. Show that OZ_2 bisects $\angle Z_1 O Z_3$.
(d) A sequence z_1, z_2, z_3, \dots satisfy $z_{m+1} = z_n^2 + z_1$ for all $n \geq 1$. If $z_1 = i$, find the distinct values that occur in the sequence.
3. (a) A hyperbola has equation $9x^2 - 16y^2 = 144$
(i) Prove that the eccentricity is $\frac{5}{4}$.
(ii) Find the coordinates of the foci.
(iii) Find the equations of the asymptotes.
(iv) Find the length of the latus rectum. (A latus rectum is a focal chord parallel to the directrix.)
(v) Find the equation of the normal to the hyperbola at the point $(\frac{4\sqrt{10}}{3}, 1)$
(b) The tangents at the two points P and Q with parameters θ and ϕ respectively on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ intersect at the point T . Show that
$$T = \left(\frac{a(\sin \theta - \sin \phi)}{\sin(\theta - \phi)}, \frac{b(\cos \phi - \cos \theta)}{\sin(\theta - \phi)} \right).$$

(c) Find the condition for the line $px + qy + r = 0$ to be a tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.
4. (a) Sketch, on separate diagrams, the curves
(i) $y = \frac{x-4}{x}$ (ii) $y = \frac{x^2-16}{x^2}$.


(b) (i) A particle P is projected, from a point O on the horizontal ground, with speed V at an angle θ above the horizontal, where $\tan \theta = \frac{1}{3}$. The particle passes through the point with coordinates $(3a, \frac{3a}{4})$ relative to the horizontal and vertical axes at O in the plane of motion. Show that $v^2 = 20ga$.

(ii) A particle Q is projected, from a point O at the instant when P is moving horizontally. It strikes the ground at the same place and at the same instant as P . Show that the speed of projected of Q is $\sqrt{\frac{145ga}{2}}$ and find the tangent of the angle of projection.

8. (a)



Mr Keating's crystal ball rests on a solid stand which is in the shape of a square based frustum as shown.



The stand is constructed such that the crystal ball of radius R fits snugly inside and just touches the centre of the square base. The sides, EC , of the base slope so that, if extended, they would pass through the top-most point on the ball at A and make an angle θ with the vertical AD .

(i) Show that $OB = R \cos 2\theta$

(ii) Consider a slice PQR of thickness Δx as shown taken perpendicular to AD such that $OQ = x$ units. Draw a neat sketch of the slice, determine its dimensions and show that it has a volume ΔV given by $\Delta V \approx [4 \tan^2 \theta (R + x)^2 - \pi(R^2 - x^2)] \Delta x$

(iii) Find the volume of such a solid where the angle $\theta = \frac{\pi}{6}$.