



SAINT IGNATIUS' COLLEGE

## **Trial Higher School Certificate**

**2004**

# **MATHEMATICS EXTENSION 1**

**1:25pm – 3:30 pm  
Thursday 19th August 2004**

### **Directions to Students**

- Reading Time : 5 minutes
- Working Time : 2 hours
- Write using blue or black pen. (sketches in pencil).
- Board approved calculators may be used.
- A table of standard integrals is provided at the back of this paper.
- All necessary working should be shown in every question.
- **Answer each question in the booklets provided and clearly label your name and teacher's name.**
- Total Marks 84
- Attempt Question 1 – 7
- All questions are of equal value

*This paper has been prepared independently of the Board of Studies NSW to provide additional exam preparation for students. Although references have been reproduced with permission of the Board of Studies NSW, the publication is in no way connected with or endorsed by the Board of Studies NSW.*

**Total marks (84)**

**Attempt Questions 1 – 7**

**All questions are of equal value**

Answer each question in a SEPARATE writing booklet.

---

**QUESTION 1 (12 Marks) Use a SEPARATE writing booklet. Marks**

(a) Solve  $\frac{5}{2x-1} < 3$ . 3

(b) Find the acute angle between the lines  $2x - y + 1 = 0$  and  $x + 3y - 4 = 0$ . 3  
Give answer to the nearest degree.

(c) Find the coordinates of the point that divides the interval joining  $(-2, 5)$  and  $(8, -9)$  internally in the ratio  $2 : 3$ . 2

(d) If  $\alpha, \beta, \gamma$  are the roots of the equation  $x^3 - 5x^2 - 3x + 2 = 0$ , 2  
find the value of  $\frac{1}{\alpha\beta} + \frac{1}{\beta\gamma} + \frac{1}{\gamma\alpha}$ .

(e) Write down the general solution, in terms of  $\pi$ , of the equation 2  
 $\cos \theta = -\frac{1}{2}$ .

**QUESTION 2** (12 Marks) Use a SEPARATE writing booklet.

**Marks**

- (a) Use the substitution  $x = u^2 + 1$ , for  $u > 0$ , to evaluate

**4**

$$\int_1^5 (x+1)\sqrt{x-1} \, dx.$$

- (b) Evaluate  $\int_0^{\frac{\pi}{4}} \sin^2\left(\frac{1}{2}x\right) dx$ .

**3**

- (c) Prove, using the principle of mathematical induction, that  $9^{n+2} - 4^n$  is divisible by 5, for  $n$  a positive integer.

**5**

**QUESTION 3 (12 Marks)** Use a SEPARATE writing booklet.

**Marks**


- (a) Find the exact value of  $\tan\left(2\sin^{-1}\frac{3}{4}\right)$ . **3**
- (b) Consider the function  $f(x) = \sin^{-1}(x+1) + \frac{\pi}{2}$ .
- (i) What is the domain of  $f(x)$ ? **1**
- (ii) Sketch the graph of  $y = f(x)$ . **2**
- (c) Consider the function  $f(x) = \log_e(2x+1)$ .
- (i) Write down the domain of  $f(x)$ . **1**
- (ii) Find the inverse function of  $f(x)$ , and write it in the form  $f^{-1}(x) = \dots\dots$  **2**
- (iii) Find the gradients of the graphs of  $y = f(x)$  and  $y = f^{-1}(x)$  at the origin. **1**
- (iv) On the same diagrams, draw the graphs of  $y = f(x)$  and  $y = f^{-1}(x)$ . **2**

**QUESTION 4 (12 Marks)** Use a SEPARATE writing booklet.

**Marks**

- (a) Find the coefficient of  $x^3$  in the expansion of  $(2 - x)(1 + x)^5$ . 3

(b)



A swimming pool is 20 metres long, 8 metres wide, 4 metres deep at one end, and zero depth at the other end. The floor of the pool is a plane rectangular surface.


- (i) When the depth of water at the deeper end is  $h$  metres, show that the volume ( $V \text{ m}^3$ ) of water in the pool is given by  $V = 20h^2$ . 2
- (ii) If water is being poured into the pool at the rate of  $2 \text{ m}^3/\text{minute}$ , find the rate at which the depth of the water is increasing at the deepest end, when the depth is 1 metre. 2
- (c) The value of a home business,  $\$V$ , is increasing at a rate proportional to the amount by which the value is less than  $\$4000$ .  
i.e.  $\frac{dV}{dt} = k(4000 - V)$   
Initially, the value of the business was  $\$2000$  and after 5 years it was  $\$3000$ .
- (i) Show that  $V = 4000 - Ae^{-kt}$  satisfies this equation. 1
- (ii) Find the value of  $A$  and the value of  $k$  to 4 decimal places. 2
- (iii) Find the number of years for the value of the business to grow to  $\$3800$ . 2

**QUESTION 5** (12 Marks) Use a SEPARATE writing booklet.

**Marks**

- (a) (i) Show that the derivative of  $x^2 e^{-x}$  is  $xe^{-x}(2-x)$ . 1
- (ii) Show that  $x^2 e^{-x} = 0.4$  has a root between  $x = 1$  and  $x = 2$ . 1
- (iii) Use Newton's approximation to find an approximation to the root of  $x^2 e^{-x} = 0.4$ , taking  $x = 1$  as a first approximation. 3

(b)



$ABCD$  is a cyclic quadrilateral in which  $AB = AC$ , and  $CD$  is produced to  $E$ .  
Prove that  $AD$  bisects the angle  $BDE$ .

3

- (c) In the expansion of  $(3+2x)^8$ ,  $c_r$  is the coefficient of  $x^r$ .

- (i) Show that  $\frac{c_r}{c_{r-1}} = \frac{18-2r}{3r}$ . 2
- (ii) Hence or otherwise find the largest coefficient in the expansion of  $(3+2x)^8$ . 2

**QUESTION 6 (12 Marks)** Use a SEPARATE writing booklet.

**Marks**

- (a) The position of a particle at time  $t$  is given by:

$$x = 3 \sin 2t - 4 \cos 2t.$$

- |       |   |   |
|-------|---|---|
| (i)   | Show that this equation satisfies $\ddot{x} = -n^2 x$ . | 2 |
| (ii)  | What is the initial velocity of the particle?           | 1 |
| (iii) | At what time does the particle first come to rest?      | 3 |

- (b) The acceleration of a particle at position  $x$  is given by:

$$\ddot{x} = -\frac{1}{4x^3}.$$

Initially the particle is at  $x = 1$  moving with a velocity of  $\frac{1}{2}$  unit in the positive direction.


- |     |  |   |
|-----|--|---|
| (i) | Prove that the velocity of the particle at position $x$ is given by: | 3 |
|-----|--|---|

$$v = \frac{1}{2x}.$$

- |      |   |   |
|------|---|---|
| (ii) | Hence find the position of the particle at time $t$ . | 3 |
|------|---|---|

**QUESTION 7** (12 Marks) Use a SEPARATE writing booklet.

**Marks**



An object is thrown from ground level with a speed of 40 m/s at an angle of  $60^\circ$  to the horizontal. Assume acceleration due to gravity is  $10 \text{ m/s}^2$  and neglect air resistance.

- (a) Find equations for  $x$  and  $y$  in terms of time  $t$  seconds, starting from the acceleration equations  $\ddot{x} = 0$  and  $\ddot{y} = -10$ , and hence show that: **4**

$$y = \sqrt{3}x - \frac{x^2}{80}.$$

- (b) The object is thrown up a slope with a gradient of  $\frac{1}{4}$ . **2**  
Show that the horizontal distance travelled by the object when it lands on the slope is given by:

$$x = 80\sqrt{3} - 20.$$

- (c) Hence find the distance  $OA$  (to the nearest metre) up the slope from the point of projection to the point of landing. **2**

- (d) Show that the maximum height reached by the object above the slope is **4**  
 $(61.25 - 10\sqrt{3})$  metres.

**End of paper**