

Student Number:

2006
HIGHER SCHOOL CERTIFICATE
Sample Examination Paper

MATHEMATICS

Extension 1

General Instructions

- Reading Time – 5 minutes
- Working Time – 2 hours
- Write using blue or black pen
- Write your student number at the top of this page

Total marks – 84

- Attempt ALL questions.
- Show all necessary working, marks may be deducted for careless or untidy work.
- Board-approved calculators may be used.
- Additional Answer Booklets are available.

Directions to school or college

To ensure maximum confidentiality and security, examination papers used for trial examinations must NOT be removed from the examination room or used with students for revision purposes until Monday 4 September 2006.

The purchasing educational institution and its staff are permitted to photocopy and/or cut and paste examination papers for educational purposes, within the confines of the educational institution, provided that: 1. the number of copies does not exceed the number reasonably required by the educational institution to satisfy their teaching purposes; 2. copies are not sold or lent.

All care has been taken to ensure that this sample examination paper is error free and that it follows the style, format and material content of the current NSW syllabus. Candidates are advised that the authors of this examination paper cannot in any way guarantee that the actual Board Of Studies Examination will have a similar content or format.

QUESTION 1 (Start a new booklet)


Marks

- (a) Find the exact value of $\int_0^1 \left(\frac{1}{1+x} + e^{-x} + \frac{1}{\sqrt{1-x^2}} \right) dx$ **4**
- (b) How many numbers greater than 5000 can be formed with the digits 4, 5, 6, 7 and 8 if no digit is used more than once in a number? **3**
- (c) Solve for x if $\log_{10}(10^x + 2) = 2x + 1$ **3**
- (d) Show that $\frac{1 - \cos 2x}{\sin 2x} = \tan x$ and hence express $\tan 15^\circ$ in simplest surd form. **2**

QUESTION 2 (Start a new booklet)

Marks

- (a) The figure shows a circle centre O and a radius 2 units. It is divided into 12 sectors and perpendiculars have been drawn as indicated. The points A, B, C , etc are shown as well as the distances P_1, P_2, P_3, \dots




- (i) Find the value of P_1 by considering $\triangle OBA$. **1**
- (ii) Show that the distances P_1, P_2, P_3, \dots form a geometric series. **3**
- (iii) Find the exact length of one spiral. ($P_1 + P_2 + P_3 + \dots + P_{12}$) **2**
- (iv) If the spiral is continued indefinitely, show that its total length will not exceed 7.5 units. **1**
- (b) Using the substitution $u = e^x$, find $\int \frac{e^x}{1 + e^{2x}} dx$. **2**
- (c) Use Newton's method once to find a better root of the following equation $\cos x = \log_e x$ given that it has a root near $x = 1$. **3**

QUESTION 3 (Start a new booklet)

Marks

- (a) TA is a tangent to the circle at A , and AB is a chord.
 P is a point on AB such that $TA = PT$. The secant TB cuts the circle at C and AC and PC are joined.




- (i) Prove that $TAPC$ is a cyclic quadrilateral. 3
- (ii) Prove that TP is a tangent to circle PCB at P . 2
- (b) Differentiate with respect to x : $\cos^{-1}\left(\frac{1}{x}\right)$ 2
- (c) A bag contains twice as many white marbles as blue marbles. If a single marble is chosen at random, what is the probability that it is
- (i) white? 1
- (ii) blue? 1
- (iii) If two marbles are chosen with replacement, find the probability that they will be of different colours. 1
- (d) A particle is moving along the x -axis. Its velocity V at position x is given by $V = \sqrt{8x - x^2}$. Find the acceleration when $x = 3$. 2

QUESTION 4 (Start a new booklet)

Marks

- (a) The diagram represent the graphs of $f(x) = x^2$ and $g(x) = 7x - 6$.



- (i) What is the maximum value of AB ? **3**
- (ii) Show that the line AB at its maximum value divides the area between the curve and the straight line in half. **4**
- (b) Given that $f(x) = x^2 + bx + c$ and $g(x) = c + x + bx^2$ and if both $f(x)$ and $g(x)$ are divided by $x - t$, the remainder is 2, prove that $t = \frac{2-c}{b+1}$ **2**
- (c) (i) Express $\sin x + \sqrt{3} \cos x$ in the form $A \sin(x + \alpha)$ where α is in radians and $A > 0$. **2**
- (ii) Hence or otherwise, sketch the graph of $y = \sin x + \sqrt{3} \cos x$ for $0 \leq x \leq \pi$. **1**

QUESTION 5 (Start a new booklet)

Marks

- (a) (i) Draw graphs of $y = |\sin x|$ and $y = -\cos x$ for $0 \leq x \leq 360^\circ$. **1**
- (ii) For what values of x will $|\sin x| + \cos x = 0$? **1**
- (iii) For what values of x will $|\sin x| \cdot \cos x < 0$? **1**
- (b) A bowl of hot soup at temperature $T^\circ\text{C}$, when placed in a cooler environment, loses heat according to the law
- $$\frac{dT}{dt} = k(T - T_0)$$
- where t is the time elapsed in minutes and T_0 is the temperature of the environment in degrees Celsius.
- (i) A bowl of soup at 96°C is left to stand in a room at a temperature of 18°C . After 3 minutes the soup cools down to 75°C . Calculate the value of k to 4 decimal places. **2**
- (ii) Susan wishes to enjoy her soup at a temperature of 60°C . How long should she wait? **2**
- (c) On certain days of constant weather the variation of temperature each day follows a pattern of simple harmonic motion. If it is 13°C at 5 a.m. and 23°C at 5 p.m., at what daylight times would the temperature be
- (i) 18°C ? **3**
- (ii) 15°C ? **1**
- (iii) What would be the expected temperature at 1 p.m.? **1**

QUESTION 6 (Start a new booklet)**Marks**

- (a) Differentiate $(x^2 + 2x + 2)e^{-x}$ and hence evaluate $\int_1^2 x^2 e^{-x} dx$ correct to 3 decimal places. **2**
- (b) $P(2ap, ap^2)$ is a point on the parabola $x^2 = 4ay$.
- (i) Show that the equation of the normal at P is $x + py = 2ap + ap^3$. **2**
- (ii) This normal cuts the y -axis at R . State the coordinates of R . **1**
- (iii) From P a line PT is drawn perpendicular to the directrix meeting it in T . State the coordinates of T . **1**
- (iv) If M is the midpoint of RT , find the coordinates of M . **1**
- (v) Find the locus of M and show that it is a parabola with vertex at the focus of the original parabola. **3**
- (c) A point P is moving on the curve $y = 2x^3$ in such a way that its x -coordinate is changing at a constant rate of 0.5 units/s. At what rate is the gradient changing when $x = 1$? **2**

QUESTION 7 (Start a new booklet)

Marks

- (a) A particle is projected with a speed of 20 m/s and passes through a point P whose horizontal distance from the point of projection is 30 m and whose vertical height above the point of projection is $8\frac{3}{4}$ m. Taking $g = 10 \text{ m/s}^2$ and θ to be the angle of elevation, **8**
- (i) Prove that $x = 20t \cos \theta$ and $y = -5t^2 + 20t \sin \theta$.
- (ii) Find the angle of elevation of θ and the time taken for the particle to reach P .
- (b) In the expansion $(1 + x + kx^2)^9$ in ascending powers of x , the coefficient of x^2 is zero. Find the value of k . **4**

End of paper

STANDARD INTEGRALS

$\int x^n dx$	$= \frac{1}{n+1} x^{n+1}, n \neq -1; x \neq 0, \text{ if } n < 0$
$\int \frac{1}{x} dx$	$= \ln x, x > 0$
$\int e^{ax} dx$	$= \frac{1}{a} e^{ax}, a \neq 0$
$\int \cos ax dx$	$= \frac{1}{a} \sin ax, a \neq 0$
$\int \sin ax dx$	$= -\frac{1}{a} \cos ax, a \neq 0$
$\int \sec^2 ax dx$	$= \frac{1}{a} \tan ax, a \neq 0$
$\int \sec ax \tan ax dx$	$= \frac{1}{a} \sec ax, a \neq 0$
$\int \frac{1}{a^2 + x^2} dx$	$= \frac{1}{a} \tan^{-1} \frac{x}{a}, a \neq 0$
$\int \frac{1}{\sqrt{a^2 - x^2}} dx$	$= \sin^{-1} \frac{x}{a}, a > 0, -a < x < a$
$\int \frac{1}{\sqrt{x^2 - a^2}} dx$	$= \ln \left(x + \sqrt{x^2 - a^2} \right), x > a > 0$
$\int \frac{1}{\sqrt{x^2 + a^2}} dx$	$= \ln \left(x + \sqrt{x^2 + a^2} \right)$
NOTE: $\ln x = \log_e x, x > 0$	

This page has been left intentionally blank

2006 MATHEMATICS EXTENSION 1 HSC TRIAL
Examination Mapping Grid

Question	Marks	Content	Syllabus Outcomes	Targeted Performance Bands
1(a)	4	Integration of various functions	HE4	E2 – E3
1(b)	3	Permutations and combinations	PE3	E2 – E3
1(c)	3	Logarithmic functions	H3	E2 – E3
1(d)	2	Trigonometric functions	PE2 H5	E2 – E4
2(a)	7	Sequences and series	H5	E2 – E3
2(b)	2	Integration by substitution	HE6	E2 – E4
2(c)	3	Polynomials	PE3	E2 – E4
3(a)	5	Circle Geometry	PE3	E2 – E4
3(b)	2	Differentiation of inverse trigonometric functions	HE4/5	E2 – E3
3(c)	3	Probability	H5	E2 – E3
3(d)	2	Applications of calculus to the physical world	HE5	E2 – E3
4(a)	7	Calculus	HE1 HE4 PE6	E2 – E3
4(b)	2	Real functions of a real variable	PE3	E2 – E4
4(c)	3	Trigonometry	HE1 H9	E2 – E3
5(a)	3	Trigonometry	PE2 H5	E2 – E4
5(b)	4	Applications of calculus to the physical world	HE3	E2 – E3
5(c)	5	Applications of calculus to the physical world	HE3	E2 – E3
6(a)	2	Calculus	HE1	E2 – E4
6(b)	8	Parametric representation	PE3	E2 – E4
6(c)	2	Calculus	HE5	E2 – E4
7(a)	8	Applications of calculus to the physical world	HE3	E2 – E4
7(b)	4	Binomial theorem	HE3	E2 – E4

SOLUTIONS**QUESTION 1**

$$\begin{aligned}
 \text{(a)} \quad & \int_0^1 \left(\frac{1}{1+x} + e^{-x} + \frac{1}{\sqrt{1-x^2}} \right) dx \\
 &= \left[\log(1+x) - e^{-x} + \sin^{-1} x \right]_0^1 \\
 &= \log 2 - e^{-1} + \sin^{-1} 1 - (\log 1 - e^0 + \sin^{-1} 0) \\
 &= \log 2 - \frac{1}{e} + \frac{\pi}{2} + 1
 \end{aligned}$$

(b) If all 5 digits are used: number of permutations = $5!$
 $= 120$

If only 4 digits are used

4	4	3	2
---	---	---	---

Number of permutations = $4 \times 4 \times 3 \times 2 = 96$

$\therefore \text{Total} = 216$

(c) $\log_{10}(10^x + 2) = 2x + 1$
 $10^x + 2 = 10^{2x+1}$
 $10^x + 2 = 10(10^{2x})$

Let $10^x = a$

$\therefore 10a^2 - a - 2 = 0$

$(5a + 2)(2a - 1) = 0$

$a = -\frac{2}{5}, a = \frac{1}{2}$

$10^x = -\frac{2}{5}; 10^x = \frac{1}{2}$

No solution; $x = \log_{10} \frac{1}{2}$

$$\begin{aligned} \text{(d)} \quad \frac{1 - \cos 2x}{\sin 2x} &= \frac{1 - (1 - 2\sin^2 x)}{2\sin x \cos x} \\ &= \frac{2\sin^2 x}{2\sin x \cos x} \\ &= \frac{\sin x}{\cos x} \\ &= \tan x \\ \tan 15^\circ &= \frac{1 - \cos 30^\circ}{\sin 30^\circ} \\ &= \frac{1 - \frac{\sqrt{3}}{2}}{\frac{1}{2}} \\ &= 2 - \sqrt{3} \end{aligned}$$

QUESTION 2

(a) (i) $OA = 2$ $\angle AOB = 30^\circ$ $\therefore P_1 = 1$ and $OB = \sqrt{3}$

(ii) $OB = \sqrt{3}$ $\angle BOC = 30^\circ$ $\therefore P_2 = \sqrt{3} \sin 30^\circ = \frac{\sqrt{3}}{2}$

and $OC = OB \cos 30^\circ = \sqrt{3} \cdot \frac{\sqrt{3}}{2} = \frac{3}{2}$

$OC = \frac{3}{2}$, $\angle COD = 30^\circ$, $\therefore P_3 = \frac{3}{2} \sin 30^\circ = \frac{3}{4}$

and $OD = OC \cos 30^\circ = \frac{3}{2} \cdot \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{4}$

$$\frac{P_3}{P_2} = \frac{\frac{3}{4}}{\frac{\sqrt{3}}{2}} = \frac{3}{4} \times \frac{2}{\sqrt{3}} = \frac{\sqrt{3}}{2}$$

$$\frac{P_2}{P_1} = \frac{\frac{\sqrt{3}}{2}}{1} = \frac{\sqrt{3}}{2}$$

$\therefore P_1, P_2, P_3, \dots$ form a G.P.

(iii) Length of 1 spiral $S_{12} = \frac{a(r^{12} - 1)}{r - 1} = \frac{a(1 - r^{12})}{1 - r}$

$$= \frac{1 \left[1 - \left(\frac{\sqrt{3}}{2} \right)^{12} \right]}{1 - \frac{\sqrt{3}}{2}}$$

$$= \frac{\left[1 - \frac{729}{4096} \right]}{\frac{2 - \sqrt{3}}{2}}$$

$$= \frac{(3367) \times 2}{4096(2 - \sqrt{3})}$$

$$= \frac{3367}{2048(2 - \sqrt{3})}$$


$$\begin{aligned}
 \text{(iv) } S_{\infty} &= \frac{a}{1-r} \\
 &= \frac{1}{1-\frac{\sqrt{3}}{2}} \\
 &= \frac{2}{2-\sqrt{3}} \\
 &\approx 7.464 < 7.5
 \end{aligned}$$

$$\begin{aligned}
 \text{(b) } u &= e^x \\
 du &= e^x dx \\
 \therefore \int \frac{du}{1+u^2} &= \tan^{-1} u + c \\
 &= \tan^{-1}(e^x) + c
 \end{aligned}$$

$$\begin{aligned}
 \text{(c) Let } f(x) &= \cos x - \log x \\
 f'(x) &= -\sin x - \frac{1}{x} \\
 x_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} \\
 &= 1 - \frac{\cos 1 - \log 1}{-\sin 1 - 1} \\
 &\approx 1.29
 \end{aligned}$$

QUESTION 3

(a) (i)



$$\angle PAT = \angle TPA \quad (TA = TP)$$

$$\therefore \angle PAC + \angle TAC = \angle TPA$$

$$\angle TPA = \angle ABT + \angle PTB \quad (\text{exterior angle of } \triangle PBT)$$

$$\therefore \angle ABT + \angle PTB = \angle PAC + \angle TAC$$

But $\angle TAC = \angle TBA$ (angle between tangent and chord)

$$\therefore \angle PTB = \angle PAC$$

$\therefore TACP$ is a cyclic quadrilateral (angles subtended by arc PC are equal).

(ii) $\angle TAC = \angle TPC$ ($TAPC$ is a cyclic quadrilateral)

But $\angle TAC = \angle ABC$ (angle between tangent and chord)


$$\therefore \angle TPC = \angle ABC$$

$\therefore PT$ is a tangent to circle PCB at P .

(b) Let $u = \frac{1}{x}$

$$\begin{aligned} y &= \cos^{-1} u \\ \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} = -\frac{1}{\sqrt{1-u^2}} \left(-\frac{1}{x^2} \right) \\ &= -\frac{1}{\sqrt{1-\frac{1}{x^2}}} \left(-\frac{1}{x^2} \right) \\ &= \frac{-x}{\sqrt{x^2-1}} \left(-\frac{1}{x^2} \right) \\ &= \frac{1}{x\sqrt{x^2-1}} \end{aligned}$$

(c)



(i) $P(E) = \frac{2}{3}$

(ii) $P(E) = \frac{1}{3}$

(iii)
$$P(E) = \frac{2}{3} \left(\frac{1}{3} \right) + \frac{1}{3} \left(\frac{2}{3} \right)$$

$$= \frac{4}{9}$$

(d) $v = \sqrt{8x - x^2}$

$$\therefore v^2 = 8x - x^2$$

$$\frac{1}{2}v^2 = 4x - \frac{x^2}{2}$$


$$a = \frac{d}{dx} \left(4x - \frac{x^2}{2} \right)$$

$$a = 4 - x$$

When $x = 3$, $a = 1$

QUESTION 4

(a) (i)



Let distance $AB = s$

$$s = 7x - 6 - x^2$$

$$\frac{ds}{dx} = 7 - 2x = 0 \text{ for a maximum or minimum}$$

$$x = \frac{7}{2}$$

$$\frac{d^2s}{dx^2} = -2 \therefore \text{maximum}$$

$$\begin{aligned} s \text{ maximum} &= 7 \cdot \frac{7}{2} - 6 - \left(\frac{7}{2}\right)^2 \\ &= \frac{49}{2} - 6 - \frac{49}{4} \\ &= \frac{25}{4} \end{aligned}$$

(ii) The parabola and the line meet when $x^2 = 7x - 6$

$$x^2 - 7x + 6 = 0$$

$$(x - 6)(x - 1) = 0$$

$$x = 6, x = 1$$

Area between the parabola and the line is

$$\begin{aligned} &\int_1^6 (7x - 6 - x^2) dx \\ &= \left[\frac{7x^2}{2} - 6x - \frac{x^3}{3} \right]_1^6 \\ &= 126 - 36 - 72 - \left(\frac{7}{2} - 6 - \frac{1}{3} \right) \\ &= \frac{125}{6} \end{aligned}$$

Area between parabola and line and line $x = \frac{7}{2}$ is

$$\begin{aligned}\int_1^{\frac{7}{2}} (7x - 6 - x^2) dx &= \left[\frac{7x^2}{2} - 6x - \frac{x^3}{3} \right]_1^{\frac{7}{2}} \\ &= \frac{343}{8} - 21 - \frac{343}{24} - \left(\frac{7}{2} - 6 - \frac{1}{3} \right) \\ &= \frac{125}{12}\end{aligned}$$

\therefore Line AB divides the area in half.

(b) $f(x) = x^2 + bx + c$ $f(t) = t^2 + tb + c = 2 \dots (1)$

$g(x) = c + x + bx^2$ $g(t) = c + t + bt^2 = 2 \dots (2)$

From (1) $t^2 = 2 - tb - c$

Sub in (2) $c + t + b(2 - tb - c) = 2$

$$c + t + 2b - tb^2 - bc = 2$$

$$t(1 - b^2) = 2 - c - 2b + bc$$

$$= 2 - c - b(2 - c)$$

$$t(1 - b)(1 + b) = (2 - c)(1 - b)$$

$$t(1 + b) = 2 - c$$


$$t = \frac{2 - c}{1 + b}$$

(c) (i) $A = \sqrt{1+3} = 2$


$$\tan \alpha = \frac{\sqrt{3}}{1}$$

$$\therefore \alpha = \frac{\pi}{3}$$

$$\therefore y = 2 \sin\left(x + \frac{\pi}{3}\right)$$



QUESTION 5



(ii) The solutions are found where the graphs intersect (see A and B)

$$|\sin x| = -\cos x$$

when $x = 135^\circ$ or 225°

(iii) $\cos x |\sin x| < 0$

when $-\cos x |\sin x| > 0$

i.e. when $90^\circ < x < 180^\circ$ and

$180^\circ < x < 270^\circ$

(b) (i) The solution of the differential equation is

$$T = T_0 + Ae^{-kt} \text{ where } A \text{ is a constant}$$

When $t = 0$, $T = 96$, $T_0 = 18$ so that $A = 78$

$$T = 18 + 78e^{-kt}$$

When $t = 3$, $T = 75^\circ$

$$75 = 18 + 78e^{-3k}$$

$$57 = 78e^{-3k}$$

$$e^{-3k} = \frac{57}{78}$$

$$-3k = \ln \frac{57}{78}$$

$$k = \frac{\ln \frac{57}{78}}{-3}$$

$$\approx 0.1046$$

$$(ii) \quad 60 = 18 + 78e^{-0.1046t}$$

$$42 = 78e^{-0.1046t}$$


$$\frac{42}{78} = e^{-0.1046t}$$

$$-0.1046t = \ln \frac{42}{78}$$

$$t = \frac{\ln \frac{42}{78}}{-0.1046}$$

$$\approx 5.92 \text{ minutes}$$

- (c) Let x be the number of degrees by which the temperature differs from the mean temperature at time t after 5 a.m. (This places the origin of the temperature at mean temperature and the origin of time at 5 a.m.)



The motion is simple harmonic with amplitude 5°C and period 24 hours

$$\therefore T = \frac{2\pi}{n} \text{ and } n = \frac{\pi}{12}$$

$$\therefore x = -5 \cos \frac{\pi}{12} t$$

- (i) When temperature = 18°C

$$x = 0$$

$$\therefore \cos \frac{\pi t}{12} = 0$$

$$\frac{\pi t}{12} = \frac{\pi}{2}$$

$$t = 6 \text{ h}$$

\therefore Time would be 11 a.m.

$$(ii) \quad -5 \cos \frac{\pi t}{12} = -3$$

$$\cos \frac{\pi t}{12} = 0.6$$

$$\frac{\pi t}{12} = 0.927$$

$$t = 3 \text{ h } 32 \text{ mins}$$

\therefore Time would be 8.32 a.m.

(iii) At 1 p.m. $t = 8$

$$x = -5 \cos \frac{\pi t}{12}$$

$$x = -5 \cos \frac{8\pi}{12}$$

$$= -5 \cos \frac{2\pi}{3}$$


$$= -5 \left(-\frac{1}{2} \right)$$

$$= \frac{5}{2}$$

\therefore Temperature is $18 + 2.5$
 $= 20.5^\circ\text{C}$

QUESTION 6

$$\begin{aligned}
 \text{(a)} \quad \frac{d}{dx} [x^2 + 2x + 2] e^{-x} &= (2x + 2)e^{-x} - e^{-x}(x^2 + 2x + 2) \\
 &= e^{-x}(2x + 2 - x^2 - 2x - 2) \\
 &= e^{-x}(-x^2) \\
 \therefore \int_1^2 x^2 e^{-x} dx &= \left[\frac{x^2 + 2x + 2}{e^x} \right]_1^2 \\
 &= \frac{5}{e} - \frac{10}{e^2} \\
 &\approx 0.486
 \end{aligned}$$



(i) $x^2 = 4ay$

$$y = \frac{x^2}{4a}$$

$$\frac{dy}{dx} = \frac{x}{2a}$$

$$\text{At } x = 2ap, \frac{dy}{dx} = \frac{2ap}{2a} = p$$

$$\therefore \text{Gradient of normal at } P \text{ is } -\frac{1}{p}$$

\therefore Equation of normal at P is

$$y - ap^2 = -\frac{1}{p}(x - 2ap)$$

$$py - ap^3 = -x + 2ap$$

$$\text{i.e. } x + py = ap^3 + 2ap$$

(ii) Normal cuts y -axis when $x = 0$

$$\text{i.e. } py = ap^3 + 2ap$$

$$\text{i.e. } y = ap^2 + 2a$$

$$\therefore \text{Coordinates of } R(0, ap^2 + 2a)$$

(iii) Coordinates of $T(2ap, -a)$

(iv) Coordinates of $M\left(ap, \frac{ap^2 + a}{2}\right)$

(v) $x = ap$

$$y = \frac{ap^2 + a}{2}$$

$$\therefore p = \frac{x}{a}$$

$$y = \frac{a\left(\frac{x}{a}\right)^2 + a}{2}$$

$$2y = \frac{x^2}{a} + a$$

$$2ay = x^2 + a^2$$

$$x^2 = 2ay - a^2$$

$x^2 = 2a(y - a)$ which is a parabola with vertex at $(0, a)$, the focus of the original parabola

(c) Gradient $= m = \frac{dy}{dx}$
 $= 6x^2$

$$\begin{aligned} \frac{dm}{dt} &= \frac{dm}{dx} \cdot \frac{dx}{dt} \\ &= 12x(0.5) \\ &= 12(1)(0.5) \\ &= 6 \text{ u / s} \end{aligned}$$

QUESTION 7

(a) (i) $\ddot{x} = 0, \dot{x} = 20 \cos \theta, x = 20t \cos \theta$

$$\ddot{y} = -10, \dot{y} = -10t + 20 \sin \theta, y = -5t^2 + 20t \sin \theta$$

(ii) At P $30 = 20t \cos \theta$

$$3 = 2t \cos \theta \quad \therefore t = \frac{3}{2 \cos \theta}$$

$$\frac{35}{4} = -5t^2 + 20t \sin \theta$$

$$35 = -20t^2 + 80t \sin \theta$$

$$7 = -4t^2 + 16t \sin \theta$$

$$\therefore 7 = -4 \left(\frac{3}{2 \cos \theta} \right)^2 + 16 \left(\frac{3}{2 \cos \theta} \right) \sin \theta$$

$$= -9 \sec^2 \theta + 24 \tan \theta$$

$$= -9(\tan^2 \theta + 1) + 24 \tan \theta$$

$$\therefore 9 \tan^2 \theta - 24 \tan \theta + 16 = (3 \tan \theta - 4)^2$$

$$= 0$$

$$\theta = \tan^{-1} \frac{4}{3} \quad \text{and} \quad \cos \theta = \frac{3}{5}$$

$$= 53^\circ 08'$$

$$t = \frac{3}{2 \left(\frac{3}{5} \right)}$$

$$= 2.5 \text{ s}$$

(b) $(1 + x + kx^2)^9 = 1 + \binom{9}{1}(x + kx^2) + \binom{9}{2}(x + kx^2)^2 + \dots$

$$= 1 + 9x + 9kx^2 + 36(x + 2kx^3 + \dots) + \dots$$

$$= 1 + 9x + (9k + 36)x^2 + \dots$$

$$\therefore 9k + 36 = 0$$

$$k = -4$$

2006 Mathematics Extension 1 HSC Trial
Marking Guidelines

Questions			Marks	Criteria
1	(a)		3	Integration
			1	Evaluation
1	(b)		1	All digits
			1	Four digits
			1	Total
1	(c)		1	Index form
			1	Quadratic equation
			1	Solution
1	(d)		1	Identity
			1	Substitution
2	(a)	(i)	1	Values of P_1
		(ii)	3	Ratio constant
		(iii)	2	Length of one spiral
		(iv)	1	Evaluation
2	(b)		1	Substitution
			1	Integration
2	(c)		1	First derivative
			1	Correct substitution into formulae
			1	Evaluation
3	(a)	(i)	3	Statements with reasons
		(ii)	2	Statements with reasons
3	(b)		1	Chain rule
			1	Algebra
3	(c)	(i)	1	Correct answer
		(ii)	1	Correct answer
		(iii)	1	Correct answer
3	(d)		1	Acceleration
			1	Evaluation
4	(a)	(i)	1	Expression for AB
			1	Differentiation
			1	Maximum value
		(ii)	1	Coordinates of intersection
			1	First area
			1	Second area
			1	Conclusion
4	(b)		1	$f(t)$ and $g(t)$
			1	Various methods of obtaining result
4	(c)	(i)	1	Value of A
			1	Value of α
		(ii)	1	Graph
5	(a)	(i)	1	Graphs
			1	Answer
			1	Answer
5	(b)	(i)	1	Solution of differential equation
			1	Value of k
		(ii)	1	Correct substitution

Questions			Marks	Criteria
			1	Answer
5	(c)	(i)	1	Value of n
			1	Trigonometric equation
			1	Time at 18°C
		(ii)	1	Time at 16°C
		(iii)	1	Temperature at 1 o'clock
6	(a)		1	Differentiation
			1	Integration
6	(b)	(i)	1	Gradient of normal
			1	Equation of normal
		(ii)	1	Coordinates of R
		(iii)	1	Coordinates of T
		(iv)	1	Coordinates of M
		(v)	3	Locus of M and vertex
	(c)		1	Gradient
			1	Rate
7	(a)	(i)	1	Horizontal motion
			1	Vertical motion
		(ii)	1	T in terms of θ
			1	Quadratic equation of t
			1	Quadratic equation of $\tan \theta$
			1	Factorisation and solution
			1	θ
			1	Time
7	(b)		1	Expansion
			1	Terms containing x^2
			1	Equation
			1	Solution