

Question 1

a. Outcomes assessed : H5

Marking Guidelines

Criteria	Marks
• writes primitive function	1
• evaluates by substitution of limits	1

Answer

$$\int_0^{\frac{\pi}{6}} \sec 2x \tan 2x \, dx = \frac{1}{2} [\sec 2x]_0^{\frac{\pi}{6}} = \frac{1}{2} (\sec \frac{\pi}{3} - \sec 0) = \frac{1}{2} (2 - 1) = \frac{1}{2}$$

b. Outcomes assessed : P4

Marking Guidelines

Criteria	Marks
• substitutes gradients into expression for $\tan \theta$	1
• calculates θ to required accuracy	1

Answer

$$3x - y - 2 = 0$$

$$y = 3x - 2$$

Gradient is 3

$$x + 2y - 3 = 0$$

$$y = -\frac{1}{2}x + \frac{3}{2}$$

Gradient is $-\frac{1}{2}$

Acute angle θ between the lines is given by

$$\tan \theta = \left| \frac{3 - (-\frac{1}{2})}{1 + 3(-\frac{1}{2})} \right| = 7.$$

$\therefore \theta \approx 82^\circ$ (to the nearest degree)

c. Outcomes assessed : PE3, P4

Marking Guidelines

Criteria	Marks
i • shows $P(1)=0$ by substitution	1
ii • deduces that equation $P(x)=0$ has 3 real roots provided $x^2 + kx + 1 = 0$ has real roots.	1
• finds discriminant of this quadratic in terms of k and realizes $\Delta \geq 0$ for real roots	1
• states values of k	1

Answer

i. $P(1) = 1 + (k-1) + (1-k) - 1 = 0.$

ii. Equation $P(x)=0$ has 3 real roots if equation $x^2 + kx + 1 = 0$ has two real roots.

For this quadratic equation, $\Delta = k^2 - 4 \geq 0$ for $k^2 \geq 4.$

Hence $P(x)=0$ has 3 real roots for $k \leq -2$ or $k \geq 2.$

DISCLAIMER


The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

d. Outcomes assessed : PE2, PE3

Marking Guidelines

Criteria	Marks
ii • quotes the alternate segment theorem in circle APB	1
iii • quotes theorem about angles standing on the same arc (or chord) in circle AQB	1
iv • writes a sequence of deductions leading to a test for $BCDP$ to be cyclic	1
• supports these deductions with reasons	1

Answer



ii. In circle APB , angle between tangent DP and chord PA is equal to the angle subtended by PA in the alternate segment at B .
Hence $\angle DPA = \angle PBA$.

iii. In circle AQB , angles subtended by the same arc CA at points B and Q on the circumference are equal.
Hence $\angle CQA = \angle CBA$.

iv. $\angle QDP + \angle DPQ + \angle DQP = 180^\circ$ (Angle sum of $\triangle QPD$ is 180°)
But $\angle QDP = \angle CDP$, $\angle DPQ = \angle DPA$, $\angle DQP = \angle CQA$ (Q, C, D collinear; P, A, Q collinear)
Hence $\angle CDP + \angle DPA + \angle CQA = 180^\circ$.
 $\therefore \angle CDP + \angle PBA + \angle CBA = 180^\circ$ ($\angle DPA = \angle PBA$, $\angle CQA = \angle CBA$ shown above)
But $\angle PBA + \angle CBA = \angle PBC$ (by addition of adjacent angles)
 $\therefore \angle CDP + \angle PBC = 180^\circ$
Hence $BCDP$ is a cyclic quadrilateral (one pair of opposite angles supplementary)

Question 2

a. Outcomes assessed : H5

Marking Guidelines

Criteria	Marks
• uses the equivalence of expressions 3^x and $e^{x \ln 3}$	1
• derives the equivalent exponential function with base e .	1

Answer

$$3^x = e^{\ln 3^x} = e^{x \ln 3} \quad \text{Hence} \quad \frac{d}{dx} 3^x = \frac{d}{dx} e^{x \ln 3} = \ln 3 e^{x \ln 3} = 3^x \ln 3$$

b. Outcomes assessed : P4

Marking Guidelines

Criteria	Marks
• applies an appropriate process to determine the coordinates	1
• calculates both coordinates correctly	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.
No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Answer

$$\begin{array}{ccc}
 A(-3, 7) & & B(4, -2) \\
 & \swarrow \quad \searrow & \\
 & 3 \quad : \quad 2 & \\
 \hline
 P\left(\frac{3 \times 4 + 2 \times (-3)}{3+2}, \frac{3 \times (-2) + 2 \times 7}{3+2}\right)
 \end{array}$$

Hence the point of internal division is $P\left(\frac{6}{5}, \frac{8}{5}\right)$.

c. Outcomes assessed : H5

Marking Guidelines

Criteria	Marks
• uses double angle identities for sine and cosine	1
• rearranges and factorises resulting equation	1
• solves $\cos x = 0$ in required domain	1
• solves $\tan x = 1$ in required domain	1

Answer

$$\begin{aligned}
 1 + \cos 2x &= \sin 2x, \quad 0 \leq x \leq 2\pi & \therefore \cos x = 0 & \text{ or } \cos x = \sin x \\
 2 \cos^2 x &= 2 \sin x \cos x & & 1 = \tan x \\
 \cos x (\cos x - \sin x) &= 0 & \therefore x = \frac{\pi}{2}, \frac{3\pi}{2} & \text{ or } x = \frac{\pi}{4}, \frac{5\pi}{4} \\
 & & \therefore x = \frac{\pi}{4}, \frac{\pi}{2}, \frac{5\pi}{4}, \frac{3\pi}{2}
 \end{aligned}$$

d. Outcomes assessed : P4, PE3

Marking Guidelines

Criteria	Marks
i • finds gradients of OP and OQ and sets product equal to -1	1
ii • uses appropriate rectangle property to find the coordinates of R .	1
iii • writes y coordinate of R in terms of sum and product of p and q .	1
• substitutes for sum and product of p and q to find Cartesian equation.	1

Answer

- i Gradient $OP = \frac{ap^2}{2ap} = \frac{1}{2}p$. Similarly gradient $OQ = \frac{1}{2}q$.
 $\therefore OP \perp OQ \Rightarrow \frac{1}{2}p \cdot \frac{1}{2}q = -1 \quad \therefore pq = -4$
- ii The diagonals of a rectangle bisect each other. Hence M is the midpoint of OR .
Hence at R , $\frac{1}{2}(x+0) = a(p+q)$ and $\frac{1}{2}(y+0) = \frac{1}{2}a(p^2+q^2)$.
 $\therefore x = 2a(p+q)$ and $y = a(p^2+q^2)$
- iii At R , $y = a\{[p+q]^2 - 2pq\} = a\left[\left(\frac{x}{2a}\right)^2 + 8\right]$
Hence locus of R has equation $x^2 = 4a(y-8a)$.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 3

a. Outcomes assessed : P5, P8, H6, HE4

Marking Guidelines

Criteria	Marks
i • shows $f(-x) = f(x)$ to deduce function f is even	1
ii • shows formally that required limit is 1	1
iii • finds the first derivative, showing it is zero at the origin	1
• shows the origin is a maximum turning point by applying first or second derivative test	1
iv • shows the two vertical asymptotes and the central branch of the curve	1
• shows the horizontal asymptote and the remaining branches of the curve	1
v • makes x the subject, interchanges x and y to obtain equation for the inverse g^{-1}	1
• writes the domain of the inverse function	1

Answer

i. $f(-x) = \frac{(-x)^2}{(-x)^2 - 1} = \frac{x^2}{x^2 - 1} = f(x)$,
 $x \neq \pm 1$. Hence f is an even function.

ii. $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{1}{1 - \frac{1}{x^2}} = \frac{1}{1 - 0} = 1$


Curve has horizontal asymptote $y = 1$ as $x \rightarrow \pm\infty$

iii. $\frac{dy}{dx} = \frac{2x(x^2 - 1) - x^2 \cdot 2x}{(x^2 - 1)^2}$
 $= \frac{-2x}{(x^2 - 1)^2}$

$\therefore \frac{dy}{dx} = 0$ when $x = 0$

Sign of $\frac{dy}{dx}$ $\begin{array}{c|c|c|c} + & + & 0 & - \\ -1 & 0 & 1 & \end{array}$ x
 Curve \nearrow \mid \searrow \nearrow \mid \searrow

iv.



Hence $(0, 0)$ is a maximum turning point.

v. $y = \frac{x^2}{x^2 - 1}$, $x \geq 0$

$y(x^2 - 1) = x^2$

$yx^2 - y = x^2$

$yx^2 - x^2 = y$

$x^2(y - 1) = y$

$x^2 = \frac{y}{y - 1}$

\therefore for the function g , $x = \sqrt{\frac{y}{y - 1}}$, since $x \geq 0$.

Interchanging x and y , $g^{-1}(x) = \sqrt{\frac{x}{x - 1}}$.

Inspection of the graph of $y = f(x)$ shows that the range of the function g is $\{y : y \leq 0 \text{ or } y > 1\}$.

Hence the domain of the inverse function g^{-1} is $\{x : x \leq 0 \text{ or } x > 1\}$.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

b. Outcomes assessed : HE2

Marking Guidelines

Criteria	Marks
• verifies that statement true for $n = 1$	1
• writes LHS of $(k + 1)^{\text{th}}$ statement in terms of RHS of k^{th} statement (assumed true)	1
• rearranges resulting expression into form of RHS of $(k + 1)^{\text{th}}$ statement	1
• deduces the required result, showing understanding of the process of mathematical induction	1

Answer

Let $S(n)$, $n = 1, 2, 3, \dots$ be the sequence of statements $\sum_{r=1}^n \frac{r+2}{r(r+1)2^r} = 1 - \frac{1}{(n+1)2^n}$, $n = 1, 2, 3, \dots$

Consider $S(1)$: $LHS = \frac{3}{1 \times 2 \times 2} = \frac{4-1}{2 \times 2^1} = 1 - \frac{1}{2 \times 2^1} = RHS$. $\therefore S(1)$ is true.

If $S(k)$ is true: $\sum_{r=1}^k \frac{r+2}{r(r+1)2^r} = 1 - \frac{1}{(k+1)2^k}$ **

Consider $S(k+1)$: $LHS = \sum_{r=1}^{k+1} \frac{r+2}{r(r+1)2^r}$

$$= \sum_{r=1}^k \frac{r+2}{r(r+1)2^r} + \frac{(k+1)+2}{(k+1)(k+2)2^{k+1}}$$

$$= 1 - \frac{1}{(k+1)2^k} + \frac{(k+1)+2}{(k+1)(k+2)2^{k+1}} \quad \text{if } S(k) \text{ is true, using **}$$

$$= 1 - \frac{2(k+2) - (k+3)}{(k+1)(k+2)2^{k+1}}$$

$$= 1 - \frac{k+1}{(k+1)(k+2)2^{k+1}}$$

$$= 1 - \frac{1}{(k+2)2^{k+1}}$$

$$= RHS$$

Hence if $S(k)$ is true, then $S(k+1)$ is true. But $S(1)$ is true, hence $S(2)$ is true, and then $S(3)$ is true and so on. Hence by mathematical induction $S(n)$ is true for all positive integers $n \geq 1$.

Question 4

a. Outcomes assessed : HE4

Marking Guidelines

Criteria	Marks
• makes x the subject of the equation of the curve	1
• expresses the volume as a definite integral with respect to y with integrand $\tan^2\left(\frac{1}{2}y\right)$	1
• uses an appropriate trig. identity to find the primitive function	1
• substitutes the limits to evaluate the exact volume	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Answer

$$y = 2 \tan^{-1} x$$

$$\frac{1}{2} y = \tan^{-1} x$$

$$\tan\left(\frac{1}{2} y\right) = x$$

Hence volume is V cubic units where

$$V = p \int_0^{\frac{p}{2}} \tan^2\left(\frac{1}{2} y\right) dy$$

$$\begin{aligned} V &= p \int_0^{\frac{p}{2}} \left\{ \sec^2\left(\frac{1}{2} y\right) - 1 \right\} dy \\ &= p \left[2 \tan\left(\frac{1}{2} y\right) - y \right]_0^{\frac{p}{2}} \\ &= p \left\{ 2 \left(\tan \frac{p}{4} - \tan 0 \right) - \left(\frac{p}{2} - 0 \right) \right\} \\ &= p \left\{ 2 - \frac{p}{2} \right\} \end{aligned}$$

Hence volume is $\frac{1}{2} p (4 - p)$ cubic units.

b. Outcomes assessed : H5, PE3**Marking Guidelines**

Criteria	Marks
i • writes equation using expressions for areas of segment and sector	1
• simplifies to obtain required equation	1
ii • writes second approximation in terms of $f(2)$, $f'(2)$ where $f(q) = q - 2 \sin q$	1
• evaluates expression for second approximation correct to 2 decimal places	1

Answer

i. $\text{area segment} = \frac{1}{2} \text{area sector}$

$$\frac{1}{2} r^2 q - \frac{1}{2} r^2 \sin q = \frac{1}{4} r^2 q$$

$$\frac{1}{4} r^2 q - \frac{1}{2} r^2 \sin q = 0$$

$$r^2 (q - 2 \sin q) = 0$$

$$\therefore r \neq 0 \Rightarrow q - 2 \sin q = 0$$

ii. Let $f(q) = q - 2 \sin q$

Then $f'(q) = 1 - 2 \cos q$

Using Newton's method with $q_1 = 2$,

$$q_2 = 2 - \frac{f(2)}{f'(2)} \approx 2 - \frac{0.1814}{1.8323}$$

Hence second approximation is 1.90 (to 2 dec. pl.)

c. Outcomes assessed : HE3**Marking Guidelines**

Criteria	Marks
i • writes numerical expression for required probability	1
• evaluates probability as a fraction	1
ii • writes numerical expression for required probability	1
• evaluates probability as a fraction	1

Answer

Probability distribution is Binomial with $n = 6$, $p = \frac{1}{3}$, $q = \frac{2}{3}$.

i. $P(\text{exactly 2 correct}) = {}^6C_2 \left(\frac{1}{3}\right)^2 \left(\frac{2}{3}\right)^4 = 15 \times \frac{16}{729} = \frac{80}{243}$

ii. $P(\text{exactly 1 correct out of first 5, then 6th correct}) = {}^5C_1 \left(\frac{1}{3}\right)^1 \left(\frac{2}{3}\right)^4 \times \frac{1}{3} = 5 \times \frac{16}{243} \times \frac{1}{3} = \frac{80}{729}$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 5

a. Outcomes assessed : HE6

Marking Guidelines

Criteria	Marks
• writes dx in terms of du and converts x limits to u limits	1
• writes integrand in terms of u	1
• finds primitive function in terms of u	1
• evaluates integral in simplest exact form by substitution of limits	1

Answer

$$\begin{aligned}
 u &= x - 1 \\
 du &= dx \\
 x = 0.5 &\Rightarrow u = -0.5 \\
 x = 1.5 &\Rightarrow u = 0.5 \\
 2x - x^2 &= 2(u+1) - (u^2 + 2u + 1) \\
 &= 1 - u^2
 \end{aligned}$$

$$\begin{aligned}
 \int_{0.5}^{1.5} \frac{1}{\sqrt{2x-x^2}} dx &= \int_{-0.5}^{0.5} \frac{1}{\sqrt{1-u^2}} du \\
 &= \left[\sin^{-1} u \right]_{-0.5}^{0.5} \\
 &= \frac{\pi}{6} - \left(-\frac{\pi}{6} \right) \\
 &= \frac{\pi}{3}
 \end{aligned}$$

b. Outcomes assessed : P4, HE5, HE7

Marking Guidelines

Criteria	Marks
i • uses similar triangles or tangent ratio to write r in terms of h	1
ii • writes $\frac{dr}{dt}$ in terms of $\frac{dh}{dt}$	1
• substitutes values of h and $\frac{dh}{dt}$	1
• finds required rate	1

Answer

- i. The ray of light from P makes equal angles with the horizontal in both right triangles. Corresponding sides in these similar triangles are in proportion.

$$\therefore \frac{r}{6} = \frac{10}{h} \text{ and hence } r = \frac{60}{h}$$

$$\text{ii. } \frac{dr}{dt} = \frac{dr}{dh} \cdot \frac{dh}{dt} = -\frac{60}{h^2} \times \frac{dh}{dt}$$

$$\text{But } \frac{dh}{dt} = -0.1. \text{ Hence when } h = 5,$$

$$\frac{dr}{dt} = \frac{60}{25} \times 0.1 = 0.24$$

Hence r is increasing at a rate of 0.24 cm s^{-1} .

c. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • differentiates $\frac{1}{2}v^2$ to find a in terms of x	1
ii • states the centre of the motion	1
• states the amplitude of the motion	1
iii • finds the maximum speed	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Answer

$$\text{i. } a = \frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \frac{d}{dx} (16 + 4x - 2x^2) \quad \therefore a = 4 - 4x$$

$$\begin{aligned} \text{ii. } v^2 &= 4(-x^2 + 2x + 8) \\ \therefore v^2 &= 4(x+2)(4-x) \\ v^2 &\geq 0 \Rightarrow -2 \leq x \leq 4 \end{aligned}$$

The midpoint of this interval is $x = 1$.
Hence centre of motion is 1 m to the right of O
and the amplitude is 3 m.

iii. Maximum speed occurs at the centre of the motion.

$$x = 1 \Rightarrow v^2 = 36. \quad \text{Hence maximum speed is } 6 \text{ ms}^{-1}$$

Question 6

a. Outcomes assessed : H5

Marking Guidelines

Criteria	Marks
i • states maximum rate of flow	1
ii • expresses total amount of water as a definite integral	1
• uses an appropriate trig. identity to find the primitive	1
• evaluates by substitution of limits, giving answer to nearest litre	1

Answer

$$\begin{aligned} \text{i. } 0 &\leq \sin^2 t \leq 1 \\ \therefore 0 &\leq R \leq 4 \\ \text{Maximum rate of flow is } 4 \text{ kL/min,} \\ \text{since } R &= 4 \text{ when } t = \frac{\pi}{2}. \end{aligned}$$

$$\begin{aligned} \text{ii. } \int_0^p 4 \sin^2 t \, dt &= 2 \int_0^p (1 - \cos 2t) \, dt \\ &= [2t - \sin 2t]_0^p \\ &= 2(p - 0) - (\sin 2p - \sin 0) \\ &= 2p \\ \therefore 2p \text{ kL} &\approx 6 \cdot 283 \text{ kL (to the nearest L) flows into the tank.} \end{aligned}$$

b. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • substitutes one pair of N, t values to obtain one equation in A and B	1
• similarly obtains a second equation in A and B	1
• solves simultaneously to evaluate A and B	1
ii • states limiting value of N .	1

Answer

$$\begin{aligned} \text{i. } N &= A + B e^{-t} \\ 60 &= A + B e^{-\ln 2} & 36 &= A + B e^{-\ln 5} & \text{By subtraction, } 3A &= 60 \\ &= A + B e^{\ln \frac{1}{2}} & &= A + B e^{\ln \frac{1}{5}} & \therefore A &= 20 \text{ and } B = 80 \\ &= A + \frac{1}{2} B & &= A + \frac{1}{5} B \\ \therefore 120 &= 2A + B & \text{and} & 180 &= 5A + B \end{aligned}$$

$$\begin{aligned} \text{ii. As } t &\rightarrow \infty, N \rightarrow A + B \times 0 = 20 \\ \text{Hence limiting population size is } 20. \end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

c. Outcomes assessed : P4, HE5

Marking Guidelines

Criteria	Marks
i • establishes result algebraically	1
ii • writes $\frac{dt}{dx}$ as sum of two algebraic fractions using (i)	1
• integrates and evaluates constant to find t as a function of x	1
• rearranges to find x as a function of t	1

Answer

$$i. \frac{1}{x} + \frac{1}{2-x} = \frac{(2-x)+x}{x(2-x)} = \frac{2}{x(2-x)}$$

ii. Initially particle is at $x=1$ moving right with $v=\frac{1}{2}$.

But $v = \frac{x(2-x)}{2}$ and $a = v \frac{dv}{dx}$. Hence if particle reaches $x=2$,

$v=a=0$ and particle will remain at rest at this point. Hence $1 \leq x \leq 2$.

$$\frac{dx}{dt} = \frac{x(2-x)}{2}$$

$$\frac{dt}{dx} = \frac{2}{x(2-x)}$$

$$= \frac{1}{x} + \frac{1}{2-x}$$

$$t = \ln x - \ln(2-x) + c$$

$$= \ln\left(\frac{x}{2-x}\right) + c \quad (c \text{ constant})$$

$$\left. \begin{matrix} t=0 \\ x=1 \end{matrix} \right\} \Rightarrow \ln 1 + c = 0$$

$$\therefore c = 0$$

$$\therefore t = \ln\left(\frac{x}{2-x}\right)$$

$$-t = \ln\left(\frac{2-x}{x}\right)$$

$$e^{-t} = \frac{2-x}{x}$$

$$e^{-t} = \frac{2}{x} - 1$$

$$1 + e^{-t} = \frac{2}{x}$$

$$\therefore x = \frac{2}{1+e^{-t}}$$

Question 7

a. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • uses integration to find expression for x	1
• uses integration to find expression for y	1
ii • substitutes given values to write two equations in V and a	1
• finds exact value of V	1
• finds required approximate value of a to required accuracy	1
iii • finds horizontal and vertical components of impact velocity	1
• finds speed of impact to required accuracy	1
• finds angle of impact to required accuracy	1

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Answer

i. *Horizontal component*

$$\ddot{x} = 0$$

$$\dot{x} = c_1, \quad c_1 \text{ const.}$$

$$t = 0 \quad \left\{ \begin{array}{l} \dot{x} = V \cos q \\ x = V \cos q \end{array} \right\} \Rightarrow c_1 = V \cos q \quad \therefore \dot{x} = V \cos q$$

$$x = (V \cos q)t + c_2, \quad c_2 \text{ const}$$

$$t = 0 \quad \left\{ \begin{array}{l} x = 0 \end{array} \right\} \Rightarrow c_2 = 0 \quad \therefore x = (V \cos q)t$$

Vertical component

$$\ddot{y} = -10$$

$$\dot{y} = -10t + c_3, \quad c_3 \text{ const.}$$

$$t = 0 \quad \left\{ \begin{array}{l} \dot{y} = V \sin q \\ y = V \sin q \end{array} \right\} \Rightarrow c_3 = V \sin q \quad \therefore \dot{y} = -10t + V \sin q$$

$$y = -5t^2 + (V \sin q)t + c_4, \quad c_4 \text{ const}$$

$$t = 0 \quad \left\{ \begin{array}{l} y = 0 \end{array} \right\} \Rightarrow c_4 = 0 \quad \therefore y = (V \sin q)t - 5t^2$$

ii. When $t = 4$, $x = 64$ and $y = -32$

$$\left\{ \begin{array}{l} 4V \cos q = 64 \\ 4V \sin q - 80 = -32 \end{array} \right\} \quad \therefore \left\{ \begin{array}{l} V \cos q = 16 \\ V \sin q = 12 \end{array} \right\}$$

$$\therefore V^2 (\cos^2 q + \sin^2 q) = 16^2 + 12^2$$


$$\therefore V^2 = 4^2 (4^2 + 3^2)$$

$$\text{Also } \cos q = \frac{4}{5} \text{ and } \sin q = \frac{3}{5}$$

$$\therefore V = 20, \quad q \approx 36^\circ 52'$$

iii. When $t = 4$,

$$\dot{x} = V \cos q = 16 \text{ and } \dot{y} = -40 + V \sin q = -28$$



$$v^2 = 16^2 + 28^2 \Rightarrow v \approx 32 \cdot 2$$

$$\tan a = \frac{7}{4} \Rightarrow a \approx 60^\circ 15'$$

Speed of impact is 32 ms^{-1} (to nearest integer)

Angle of impact with beach is $60^\circ 15'$ (nearest minute).

b. Outcomes assessed : H9, HE3

Marking Guidelines

Criteria	Marks
i • writes expansion as required	1
ii • differentiates both sides with respect to x	1
• substitutes $x = 1$	1
• rearranges to obtain required identity	1

Answer

i. $x(1+x)^n \equiv x + {}^nC_1 x^2 + {}^nC_2 x^3 + \dots + {}^nC_{n-1} x^n + x^{n+1}$

ii. Differentiation with respect to x gives

$$(1+x)^n + nx(1+x)^{n-1} \equiv 1 + 2 {}^nC_1 x + 3 {}^nC_2 x^2 + \dots + n {}^nC_{n-1} x^{n-1} + (n+1)x^n$$

$$\text{Substituting } x = 1, \quad 2^n + n \cdot 2^{n-1} = 1 + 2 {}^nC_1 + 3 {}^nC_2 + \dots + n {}^nC_{n-1} + (n+1)$$

$$\therefore 2 {}^nC_1 + 3 {}^nC_2 + \dots + n {}^nC_{n-1} = (n+2)2^{n-1} - (n+2) = (n+2)(2^{n-1} - 1)$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.