



**CATHOLIC SECONDARY SCHOOLS ASSOCIATION OF NSW**  
**2012 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION**  
**MATHEMATICS EXTENSION 1**

**Section I**  
**10 marks**

**Questions 1–10** (1 mark each)

**Question 1** (1 mark)

**Outcomes Assessed:** PE2

**Targeted Performance Bands:** E2

Solution	Answer	Mark
$P(-1) = -1$	A	1

**Question 2** (1 mark)

**Outcomes Assessed:** H5, HE7

**Targeted Performance Bands:** E2

Solution	Answer	Mark
$\left  \frac{2-m}{1+2m} \right  = \tan 45^\circ, m > 0$ $m = \frac{1}{3}$	A	1

**Question 3** (1 mark)

**Outcomes Assessed:** H5

**Targeted Performance Bands:** E2

Solution	Answer	Mark
$\sin \theta = \frac{2t}{1+t^2}$ and $\cos \theta = \frac{1-t^2}{1+t^2}$ $\sin \theta + \cos \theta = \frac{1+2t-t^2}{1+t^2}$	C	1

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

**Question 4 (1 mark)****Outcomes Assessed: P4, PE2****Targeted Performance Bands: E2**

Solution	Answer	Mark
$(x, y) = \left( \frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right)$ $= \left( \frac{3 \times 3 + (-2) \times (-2)}{3 + (-2)}, \frac{3 \times (-4) + (-2) \times 3}{3 + (-2)} \right)$ $= (13, -18)$	D	1

**Question 5 (1 mark)****Outcomes Assessed: PE3****Targeted Performance Bands: E2**

Solution	Answer	Mark
$\binom{6}{3} \times \binom{4}{2} = 120$	B	1

**Question 6 (1 mark)****Outcomes Assessed: HE4****Targeted Performance Bands: E3**

Solution	Answer	Mark
$f^{-1}(3) = x$ $f(x) = 3$ $\frac{2x}{x+1} = 3$ $x = -3$	A	1

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

**Question 7 (1 mark)****Outcomes assessed: PE3, PE4****Targeted Performance Bands: E2**

Solution	Answer	Mark
$x + py = 2ap + ap^3$ When $x = 0, y = a$ : $ap = 2ap + ap^3$ $ap(1 + p^2) = 0$ $p = 0$ is the only solution. Therefore there exists only one solution.	B	1

**Question 8 (1 mark)****Outcomes Assessed: HE3, HE7****Targeted Performance Bands: E3**

Solution	Answer	Mark
Let $p = 0.8$ and $q = 0.2$ $P(X \geq 5) = P(X = 5) + P(X = 6)$ $= \binom{6}{5}(0.2)(0.8)^5 + (0.8)^6$ $= (0.8)^5(6 \times 0.2 + 0.8)$ $= 2(0.8)^5$	B	1

**Question 9 (1 mark)****Outcomes Assessed: HE3****Targeted Performance Bands: E3**

Solution	Answer	Mark
$T_{k+1} = \binom{7}{k}(x^2)^{7-k}\left(\frac{2}{x}\right)^k$ $= \binom{7}{k}2^k x^{14-3k}$  When $14 - 3k = 2, k = 4$ . The coefficient of $x^2 = \binom{7}{4}2^4$ $= 560$ .	D	1

**Question 10 (1 mark)****Outcomes Assessed: HE3****Targeted Performance Bands: E3, E4**

Solution	Answer	Mark
The FALSE statement is C	C	1

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

**Section II**  
**60 marks**

**Question 11 (15 marks)**

(a) (1 mark)

*Outcomes assessed: H5*

*Targeted Performance Bands: E2*

Criteria	Mark
• Correct answer	1

*Sample answer:*

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin 2x}{x} &= 2 \lim_{x \rightarrow 0} \frac{\sin 2x}{2x} \\ &= 2\end{aligned}$$

(b) (2 marks)

*Outcomes assessed: HE6*

*Targeted Performance Bands: E2*

Criteria	Marks
• Correct answer	2
• Recognising the correct primitive	1

*Sample answer:*

$$\begin{aligned}\int_0^4 \frac{dx}{\sqrt{x^2 + 9}} &= \left[ \ln \left( x + \sqrt{x^2 + 9} \right) \right]_0^4 \\ &= \ln 9 - \ln 3 \\ &= \ln 3\end{aligned}$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (3 marks)

**Outcomes assessed: HE4, HE6**

**Targeted Performance Bands: E2–E3**

Criteria	Marks
• Correct answer	3
• Significant progress towards solution	2
• Correct integral with correct limits, after applying given substitution	1

**Sample answer:**

$$\text{Let } u = x - 8$$

$$du = dx$$

$$x = 8.5 \Rightarrow u = 0.5 \text{ and } x = 8 \Rightarrow u = 0.$$

Upon substituting:

$$\begin{aligned} & \int_0^{0.5} \frac{du}{\sqrt{(7-(u+8))(u+8-9)}} \\ &= \int_0^{0.5} \frac{du}{\sqrt{1-u^2}} \\ &= [\sin^{-1}u]_0^{0.5} \\ &= \frac{\pi}{6} \end{aligned}$$

(d) (3 marks)

**Outcomes assessed: PE3**

**Targeted Performance Bands: E2–E3**

Criteria	Marks
• Correct range of solutions given	3
• Significant progress towards solution (e.g. inequality if using solution presented below or points of intersection if using graphical techniques)	2
• Some progress towards answer (considering technique used e.g. graphical or algebraic)	1

**Sample answer:**

$$\begin{aligned} \frac{2t}{1-t} &\geq t & t &\neq 1 \\ 2t(1-t) &\geq t(1-t)^2 \\ t(1-t)(1+t) &\geq 0 \\ \therefore t &\leq -1, 0 \leq t < 1 \end{aligned}$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(e) (i) (2 marks)

**Outcomes assessed: PE2, PE3**

**Targeted Performance Bands: E2**

Criteria	Marks
• Correct answer	2
• Recognises the relationship between the tangent and the intercepts from an external point	1

**Sample answer:**

$$\begin{aligned}PC^2 &= AP \cdot BP \\12^2 &= (x + 7) \cdot x \\x &= 9 \quad (\text{since } x > 0)\end{aligned}$$

(e) (ii) (1 mark)

**Outcomes assessed: PE3**

**Targeted Performance Bands: E2**

Criteria	Mark
• Correct answer	1

**Sample answer:**

$\angle BPC = 90^\circ$  since  $BC$  is the diameter of the circle passing through  $P$ ,  $B$  and  $C$ .  
 $\therefore \triangle BPC$  is a right-angled triangle.

$$\begin{aligned}BC^2 &= BP^2 + PC^2 \\BC^2 &= 9^2 + 12^2 \\BC &= 15\end{aligned}$$

(f) (3 marks)

**Outcomes assessed: PE2, PE6, HE7**

**Targeted Performance Bands: E3**

Criteria	Marks
• Correct proof logically presented	3
• Progress towards solution	2
• Correctly identifying a relevant trigonometric relationship in one of the triangles	1

**Sample answer:**

Let  $CD = h$

$$\begin{aligned}\text{In } \triangle ADC, \tan \alpha &= \frac{h}{AC} \\ \text{In } \triangle BDC, \tan \beta &= \frac{h}{BC} \\ \therefore \frac{\tan \alpha}{\tan \beta} &= \frac{BC}{AC}\end{aligned}$$

$\triangle ABC$  is right-angled at  $C$  and  $\angle CAB = 90^\circ - 60^\circ = 30^\circ$  (from the bearings given), hence

$$\begin{aligned}\frac{BC}{AC} &= \tan 30^\circ = \frac{1}{\sqrt{3}} \\ \therefore \frac{\tan \alpha}{\tan \beta} &= \frac{1}{\sqrt{3}}\end{aligned}$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

**Question 12 (15 marks)**

(a) (i) (1 mark)

**Outcomes assessed: PE3****Targeted Performance Bands: E2**

Criteria	Mark
• Correct answer	1

**Sample answer:**

$$\alpha + \beta + \gamma = \frac{-b}{a}$$

(a) (ii) (2 marks)

**Outcomes assessed: H5, HE7****Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correctly using the result in part (i) to obtain the result	2
• Solves $f''(x) = 0$	1

**Sample answer:**

$$\begin{aligned} f'(x) &= 3ax^2 + 2bx + c \\ f''(x) &= 6ax + 2b \end{aligned}$$

$$\begin{aligned} \text{Solving } f''(x) &= 0 \\ 6ax + 2b &= 0 \\ x &= \frac{-b}{3a} \\ &= \frac{\alpha + \beta + \gamma}{3} \end{aligned}$$

(a) (iii) (1 mark)

**Outcomes assessed: HE7****Targeted Performance Bands: E3**

Criteria	Mark
• Correct answer	1

**Sample answer:**

$$\begin{aligned} x &= \frac{\alpha + \beta + \gamma}{3} \\ &= \frac{-1 + 3 + 4}{3} \\ &= 2 \end{aligned}$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (i) (2 marks)

**Outcomes assessed: HE4, HE7**

**Targeted Performance Bands: E3**

Criteria	Marks
• Correct value of $x$	2
• Correct value of $\alpha$	1

**Sample answer:**

$$\begin{aligned}\sin^{-1}x &= \alpha, \cos^{-1}x = \frac{\pi}{2} - \alpha \\ \text{Now, } \sin^{-1}x &= \cos^{-1}x \\ \text{i.e. } \alpha &= \frac{\pi}{2} - \alpha \\ \alpha &= \frac{\pi}{4} \\ \therefore \sin^{-1}x &= \frac{\pi}{4} \\ x &= \sin \frac{\pi}{4} \\ \therefore \text{The solution is } x &= \frac{\sqrt{2}}{2}.\end{aligned}$$


(b) (ii) (1 marks)

**Outcomes assessed: HE4, HE7**

**Targeted Performance Bands: E2**

Criteria	Marks
• Sketches curves correctly, showing the point of intersection	1

**Sample answer:**



**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.



(b) (iii) (2 marks)

**Outcomes assessed: HE2**

**Targeted Performance Bands: E3**

Criteria	Mark
• Correct explanation	2
• Reference to symmetry of region	1

**Sample answer:**

The region is symmetrical about the line  $y = \frac{\pi}{4}$ .

Therefore the volume of the rotated region bounded by  $y = \sin^{-1} x$  between  $y = 0$  and  $y = \frac{\pi}{4}$  (given by  $\pi \int_0^{\frac{\pi}{4}} \sin^2 y \, dy$ ) is equal to the volume of the rotated region bounded by  $y = \cos^{-1} x$  between  $y = \frac{\pi}{4}$  and  $y = \frac{\pi}{2}$  (given by  $\pi \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cos^2 y \, dy$ ).

$$\begin{aligned}\therefore V &= \pi \int_0^{\frac{\pi}{4}} \sin^2 y \, dy + \pi \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cos^2 y \, dy \\ &= 2\pi \int_0^{\frac{\pi}{4}} \sin^2 y \, dy\end{aligned}$$

(b) (iv) (2 marks)

**Outcomes assessed: H8, HE6**

**Targeted Performance Bands: E3**

Criteria	Marks
• Correct answer	2
• Substantial progress towards integrating $\sin^2 y$	1

**Sample answer:**

$$\begin{aligned}V &= 2\pi \int_0^{\frac{\pi}{4}} \sin^2 y \, dy \\ &= \pi \int_0^{\frac{\pi}{4}} (1 - \cos 2y) \, dy \\ &= \pi \left[ y - \frac{\sin 2y}{2} \right]_0^{\frac{\pi}{4}} \\ &= \frac{\pi}{4} (\pi - 2) \\ \therefore \text{The volume is } \frac{\pi}{4} (\pi - 2) \text{ units}^3.\end{aligned}$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (i) (3 marks)

**Outcomes assessed: HE2, HE7**

**Targeted Performance Bands: E3**

Criteria	Marks
• Correct solution	3
• Establishes the induction step	2
• Verifies the result for $n=2$	1

**Sample answer:**

$$\begin{aligned}
 (1) \text{ When } n = 2: \quad \text{LHS} &= \left(1 - \frac{1}{2^2}\right) \\
 &= \frac{3}{4} \\
 \text{RHS} &= \frac{2+1}{2 \times 2} \\
 &= \frac{3}{4} \\
 &= \text{LHS}
 \end{aligned}$$

$$\text{Assume true for } n = k: \left(1 - \frac{1}{2^2}\right) \times \left(1 - \frac{1}{3^2}\right) \times \left(1 - \frac{1}{4^2}\right) \times \dots \times \left(1 - \frac{1}{k^2}\right) = \frac{k+1}{2k}$$

Prove true for  $n = k + 1$ :

$$\text{i.e. } \left(1 - \frac{1}{2^2}\right) \times \left(1 - \frac{1}{3^2}\right) \times \left(1 - \frac{1}{4^2}\right) \times \dots \times \left(1 - \frac{1}{k^2}\right) \times \left(1 - \frac{1}{(k+1)^2}\right) = \frac{k+2}{2(k+1)}$$

$$\text{Now LHS} = \frac{k+1}{2k} \times \left(1 - \frac{1}{(k+1)^2}\right) \quad (\text{using assumption})$$

$$= \frac{k+1}{2k} \times \frac{(k+1)^2 - 1}{(k+1)^2}$$

$$= \frac{k^2 + 2k}{2k(k+1)}$$

$$= \frac{k+2}{2(k+1)}$$

$$= \text{RHS}$$

Hence, by the Principle of Mathematical Induction, the result holds true for all integers  $n \geq 2$ .

(c) (ii) (1 mark)

**Outcomes assessed: HE2, HE7**

**Targeted Performance Bands: E2**

Criteria	Mark
• Correct answer	1

**Sample answer:**

$$n = 100, \text{ hence the product equals } \frac{101}{200}.$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

**Question 13** (15 marks)

(a) (i) (1 mark)

**Outcomes assessed:** HE3, HE7**Targeted Performance Bands:** E2–E3

Criteria	Mark
• Correct proof	1

**Sample answer:**

$$\begin{aligned}
 T &= 100 - Ae^{-0.2t} \\
 \frac{dT}{dt} &= 0.2Ae^{-0.2t} \\
 &= 0.2(100 - T) \quad \text{as } Ae^{-0.2t} = 100 - T.
 \end{aligned}$$

(a) (ii) (1 mark)

**Outcomes assessed:** HE3, HE7**Targeted Performance Bands:** E2–E3

Criteria	Mark
• Correct value for A	1

**Sample answer:**

$$\begin{aligned}
 &\text{When } t = 0, T = 4. \\
 &4 = 100 - Ae^{-0.2 \times 0} \\
 &A = 96
 \end{aligned}$$

(a) (iii) (1 mark)

**Outcomes assessed:** HE3, HE7**Targeted Performance Bands:** E2–E3

Criteria	Mark
• Correct value for temperature	1

**Sample answer:**

$$\begin{aligned}
 T &= 100 - 96e^{-0.2t} \\
 &\text{Substitute } t = 4.5 \\
 T &= 100 - 96e^{-0.2 \times 4.5} \\
 T &= 60.9693 \dots
 \end{aligned}$$

Therefore, the temperature correct to three significant figures is 61.0°C.

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(a) (iv) (2 marks)

**Outcomes assessed: HE3, HE7**

**Targeted Performance Bands: E2–E3**

Criteria	Marks
• Correct answer	2
• Correct value of $t$ for an egg boiled from room temperature	1

**Sample answer:**

$$T = 100 - 79e^{-0.2t}$$

$$\text{Substituting } T = 60.9693 \dots$$

$$60.9693 \dots = 100 - 79e^{-0.2t}$$

$$t = \ln\left(\frac{100-60.9693\dots}{79}\right) \div -0.2$$

$$= 3.525498 \dots \approx 3.5 \text{ minutes}$$

Therefore, the egg at room temperature will cook approximately one minute faster than an egg taken from the refrigerator.

(b) (i) (3 marks)

**Outcomes assessed: HE3, HE7**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correct answers for $A$ and $\alpha$	3
• Correct value for either $A$ or $\alpha$	2
• Significant progress towards correct equations involving $A$ and $\alpha$	1

**Sample answer:**

$$x = A\cos(2t + \alpha)$$

$$\text{When } t = 0, x = 4: \quad A\cos \alpha = 4 \quad (\text{i})$$

$$\text{Now} \quad \frac{dx}{dt} = -2A\sin(2t + \alpha)$$

$$\text{When } t = 0, \frac{dx}{dt} = -8\sqrt{3} \quad A\sin \alpha = 4\sqrt{3} \quad (\text{ii})$$

$$\text{From (i) and (ii) above:} \quad \frac{A\sin \alpha}{A\cos \alpha} = \frac{4\sqrt{3}}{4}$$

$$\tan \alpha = \sqrt{3}$$

$$\alpha = \frac{\pi}{3}$$

$$\text{Substitute } \alpha = \frac{\pi}{3} \text{ into (i):} \quad A\cos \frac{\pi}{3} = 4$$

$$A = 8$$

$$\text{Therefore, } A = 8 \text{ and } \alpha = \frac{\pi}{3}.$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (ii) (2 marks)

**Outcomes assessed: HE3, HE7**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correct answer	2
• Correct equation with ONE unknown, $t$	1

**Sample answer:**

Particle is at the centre of motion when  $x = 0$ :

$$\begin{aligned}8 \cos\left(2t + \frac{\pi}{3}\right) &= 0 \\2t + \frac{\pi}{3} &= \frac{\pi}{2}, \text{ for the first time.} \\t &= \frac{\pi}{12}\end{aligned}$$

Therefore, the particle arrives at the origin after  $\frac{\pi}{12}$  seconds.

(c) (i) (3 marks)

**Outcomes assessed: HE5, HE7**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correctly shows the result	3
• Correct expression for $v^2$ , including the value of the constant	2
• Progress towards expressing the result in terms of $v$ and $x$ , e.g. using the correction version of acceleration $\frac{d}{dx}\left(\frac{1}{2}v^2\right)$	1

**Sample answer:**

$$\frac{d}{dx}\left(\frac{1}{2}v^2\right) = 2x^3 + 18x$$

$$\frac{1}{2}v^2 = \frac{1}{2}x^4 + 9x^2 + c$$

$$\text{When } x = 0, v = 9: \quad c = \frac{81}{2}$$

$$\text{Therefore,} \quad \frac{1}{2}v^2 = \frac{1}{2}x^4 + 9x^2 + \frac{81}{2}$$

$$v^2 = x^4 + 18x^2 + 81$$

$$v^2 = (x^2 + 9)^2$$

$$v = \pm(x^2 + 9)$$

However, since initially the particle has a positive velocity and there are no solutions to  $v = 0$ , the particle must always travel in a positive direction.

Hence,  $v = x^2 + 9$ .

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(c) (ii) (2 marks)

**Outcomes assessed: HE5, HE7**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correct equation for $x$ in terms of $t$	2
• Progress towards an expression for $t$ in terms of $x$	1

**Sample answer:**

$$\frac{dx}{dt} = x^2 + 9$$

$$\frac{dt}{dx} = \frac{1}{x^2 + 9}$$

$$t = \frac{1}{3} \tan^{-1} \left( \frac{x}{3} \right) + c$$

When  $x = 0$ ,  $t = 0$ , therefore  $c = 0$ .

$$t = \frac{1}{3} \tan^{-1} \left( \frac{x}{3} \right)$$

$$x = 3 \tan(3t)$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

**Question 14** (15 marks)

(a) (3 marks)

**Outcomes assessed:** H5, HE7**Targeted Performance Bands:** E2–E3

Criteria	Marks
• Correct answer	3
• Correct substitution into the formula of Newton's Method	2
• Establishing an appropriate function and determining its derivative	1

**Sample answer:**

$$3 \sin x = \ln x$$

$$\text{Let } f(x) = 3 \sin x - \ln x$$

$$f'(x) = 3 \cos x - \frac{1}{x}$$

$$\begin{aligned} x_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} \\ &= 3 - \frac{3 \sin 3 - \ln 3}{3 \cos 3 - \frac{1}{3}} \\ &= 2.79558 \dots \\ &= 2.80 \text{ (2 d.p.)} \end{aligned}$$

(b) (i) (1 mark)

**Outcomes assessed:** H5**Targeted Performance Bands:** E3, E4

Criteria	Mark
• Correct justification for result	1

**Sample answer:**

For the gradient of  $MB$ : applying Pythagoras' Theorem, rise =  $\sqrt{L^2 - x^2}$ ; run =  $x$ .

$$\text{The gradient of } MB = \frac{-\text{rise}}{\text{run}} = \frac{-\sqrt{L^2 - x^2}}{x}.$$

Since the rope is always tangent to the curve, the line  $MB$  is a tangent to  $y = f(x)$

$$\therefore \frac{dy}{dx} = \frac{-\sqrt{L^2 - x^2}}{x}.$$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies.

No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(b) (ii) (2 marks)

**Outcomes assessed: HE5, HE7**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correct answer	2
• Substantial progress towards expressing $\frac{dx}{dt}$ as a function of $x$	1

**Sample answer:**

$$\begin{aligned}\frac{dx}{dt} &= \frac{dx}{dy} \times \frac{dy}{dt} \\ &= \frac{-x}{\sqrt{L^2 - x^2}} \times 3 \\ &= \frac{-\frac{L}{2}}{\sqrt{L^2 - \frac{L^2}{4}}} \times 3 \quad \text{when } x = \frac{L}{2} \\ &= -\sqrt{3}\end{aligned}$$

Therefore, the boat approaches the pier at a rate of  $\sqrt{3} \text{ ms}^{-1}$ .

(c) (3 marks)

**Outcomes assessed: PE3, PE4, HE2**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correct proof	3
• Substantial progress towards determining both required distances or both required angles	2
• Recognising the gradient at $P$ is given by $\frac{1}{p}$ and progress towards determining a relevant distance or a relevant angle	1

**Sample answer:**

The gradient of the tangent at  $P = \frac{2a}{2ap} = \frac{1}{p}$ .

The equation of the tangent at  $P$  is given by  $y - 2ap = \frac{1}{p}(x - ap^2)$

Let the tangent at  $P$  intersect the  $x$ -axis at  $G$ .

$\therefore G$  has coordinates  $(-ap^2, 0)$ .  $P$  has coordinates  $(ap^2, 2ap)$ .  $S$  has coordinates  $(a, 0)$ .

$$SG = ap^2 + a$$

$$\begin{aligned}SP &= \sqrt{(ap^2 - a)^2 + (2ap)^2} \\ &= \sqrt{a^2p^4 + 2a^2p^2 + a^2} \\ &= \sqrt{(ap^2 + a)^2} \\ &= ap^2 + a\end{aligned}$$

Since  $SG = SP$ ,  $\triangle SPG$  is isosceles.

$\therefore \angle SGP = \angle SPG$  (angles opposite equal sides of an isosceles triangle are equal), thus the angle of inclination to the axis of the parabola equals the angle between the tangent and the focal chord.

Hence the tangent to the parabola at  $P$  is equally inclined to the axis of the parabola and the focal chord through  $P$ .

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.



(d) (i) (1 mark)

**Outcomes assessed: HE2, HE7,**

**Targeted Performance Bands: E3**

Criteria	Mark
• Correct proof	1

**Sample answer:**

The coefficient of  $x^r$  in the expansion  $(1+x)^n$  is  $\binom{n}{r}$ .

The coefficient of  $x^r$  in the expansion  $(x+1)^n$  is  $\binom{n}{n-r}$ .

Since  $(1+x)^n = (x+1)^n$ , the expansions are the same. Therefore,  $\binom{n}{n-r} = \binom{n}{r}$ .

(d) (ii) (3 marks)

**Outcomes assessed: HE2, HE7**

**Targeted Performance Bands: E4**

Criteria	Marks
• Correct proof	3
• Correctly identifies and matches coefficients of $x^{n-r}$ in the two binomial expansions	2
• Expansion of ONE relevant binomial expression	1

**Sample answer:**

$$f(r) = \binom{n}{0}\binom{n}{r} + \binom{n}{1}\binom{n}{r+1} + \cdots + \binom{n}{n-r}\binom{n}{n}.$$

Therefore,  $f(r) = \binom{n}{0}\binom{n}{n-r} + \binom{n}{1}\binom{n}{n-r-1} + \cdots + \binom{n}{n-r}\binom{n}{0}$ , using the result in part (i).

Now, consider  $(1+x)^n(1+x)^n = (1+x)^{2n}$

$$(1+x)^n(1+x)^n = \left( \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \cdots + \binom{n}{n}x^n \right) \left( \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \cdots + \binom{n}{n}x^n \right)$$

The coefficient of  $x^{n-r}$  in this expansion is:

$$\binom{n}{0}\binom{n}{n-r} + \binom{n}{1}\binom{n}{n-r-1} + \binom{n}{2}\binom{n}{n-r-2} + \cdots + \binom{n}{n-r}\binom{n}{0} = f(r) \text{ from above}$$

But the coefficient of  $x^{n-r}$  in the expansion of  $(1+x)^{2n}$  is  $\binom{2n}{n-r}$ .

Therefore,  $f(r) = \binom{2n}{n-r}$ , for  $r = 0, 1, 2, \dots, n$

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

(d) (iii) (2 marks)

**Outcomes assessed: HE2, HE7**

**Targeted Performance Bands: E3–E4**

Criteria	Marks
• Correct proof	2
• Correctly identifies and matches coefficients in relevant binomial expansions	1

**Sample answer:**

Consider  $(1+x)^{3n} = (1+x)^{2n}(1+x)^n$

$$(1+x)^{2n}(1+x)^n = \left( \binom{2n}{0} + \binom{2n}{1}x + \binom{2n}{2}x^2 + \dots + \binom{2n}{2n}x^{2n} \right) \left( \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots + \binom{n}{n}x^n \right)$$

The coefficient of  $x^n$  in this expansion is:

$$\begin{aligned} & \binom{2n}{0}\binom{n}{n} + \binom{2n}{1}\binom{n}{n-1} + \binom{2n}{2}\binom{n}{n-2} + \dots + \binom{2n}{n}\binom{n}{0} \\ &= f(n)\binom{n}{n} + f(n-1)\binom{n}{n-1} + f(n-2)\binom{n}{n-2} + \dots + f(0)\binom{n}{0}, \text{ as } f(r) = \binom{2n}{n-r} \text{ for } r = 0, 1, 2, \dots, n \end{aligned}$$

But the coefficient of  $x^n$  in the expansion of  $(1+x)^{3n}$  is  $\binom{3n}{n}$ .

Therefore,  $\binom{n}{0}f(0) + \binom{n}{1}f(1) + \dots + \binom{n}{n}f(n) = \binom{3n}{n}$ .

**DISCLAIMER**

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee or warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.