



Student Number:

2003
HIGHER SCHOOL CERTIFICATE
Sample Examination Paper

MATHEMATICS

Extension 1

General Instructions

Reading time - 5 minutes
Working time - 2 hours

- Attempt ALL questions
- Show all necessary working, marks may be deducted for careless or untidy work
- Standard integrals are printed on the last page
- Board-approved calculators may be used
- Additional Answer Booklets are available

Directions to School or College

To ensure maximum confidentiality and security, examination papers must NOT be removed from the examination room. Examination papers may not be returned to students or used for revision purposes till September 2003. These examination papers are supplied Copyright Free, as such the purchaser may photocopy and/or cut and paste same for educational purposes within the confines of the school or college.


All care has been taken to ensure that this examination paper is error free and that it follows the style, format and material content of previous Higher School Certificate Examination. Candidates are advised that authors of this examination paper cannot in any way guarantee that the 2003 HSC Examination will have a similar content or format.

Question 1.

- (a) Solve for x : $3^{x+1} = 2$ expressing the answer correct to two decimal places. 2
- (b) Find the exact value of $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} (1 + \sin^2 x) dx$ 3
- (c) State the domain and range of the function $g(x) = \frac{1}{2} \cos^{-1} \frac{x}{2}$ 2
- (d) Use the remainder theorem to fully factorise $6x^3 + 17x^2 - 4x - 3$ 3
- (e) (i) Find the general solution of the equation $\tan \alpha = -\frac{1}{\sqrt{3}}$ 1
expressing your answer in terms of π .
- (ii) Hence generate a value of α such that $-\frac{3\pi}{2} < \alpha < -\pi$ 1

Question 2

- (a) Use the substitution $u = 2 - x^2$ to find $\int \frac{x}{(2-x^2)^3} dx$ 2
- (b) 3



Two circles meet in points A and B . CAD is a double chord and BTP is a chord of the larger circle. Prove that $CT \parallel PD$.

- (c) Solve the inequality $\frac{2x-5}{x-4} \geq x$ 3
- (d) Find $\frac{d}{dx} (3^{\sqrt{x+1}})$ 2
- (e) A particle moves along the x -axis starting at $x = 1$ at time $t = 0$. 2
The velocity of the particle is $v = \frac{1}{x+3}$. Find the value of t when $x = 5$.

Question 3

- (a) Use the method of mathematical induction to prove that $2(n-3) + (n-4) + \dots + 3 + 2 + 1 = \binom{n}{2} - n$ for $n \geq 4$. 4
- (b) Find the roots of the following equation $4x^3 - 4x^2 - 29x + 15 = 0$ given that one root is the difference between the other two roots. 3
- (c) The tangent to the point $P(2ap, ap^2)$ on the parabola $x^2 = 4ay$ cuts the x -axis in A and the y -axis in B .
- (i) Find the coordinates of M , the midpoint of A and B in terms of P . 2
- (ii) Show that the locus of M is a parabola. 2
- (iii) Find the coordinates of the focus of this parabola and the equation of its directrix. 1

Question 4

- (a) (i) How many 11 letter 'words' can be formed from the letters of the word 'PROBABILITY'? 1
- (ii) In how many of these does the word BABY appear? 1
- (b) A surveyor observes two towers, one due north of height 80m, and the other on a bearing of θ° ($< 90^\circ$) of height 120m. The angles of elevation of the two towers are 40° and 36° respectively. If the towers are 150m apart on a horizontal plane, calculate the value of θ to the nearest minute. 6
- (c) By considering the expansion of $x(1+x)^n$ or otherwise, show that 4

$$\sum_{r=0}^n (r+1)^n C_r = 2^n \left(\frac{n}{2} + 1 \right)$$

Question 5

- (a) A golf ball is to be struck so as to clear a tree 20m away and 6m high on level ground. If the selected club produces an angle of elevation of 40° , (take $g = 10\text{m/s}^2$)
- (i) Write down an expression for y , the vertical distance travelled. 1
 - (ii) Write down an expression for x , the horizontal distance travelled. 1
 - (iii) Hence derive the cartesian equation of the flight path. 1
 - (iv) Calculate the speed at which the ball must leave the ground in order to just clear the obstacle. 2
- (b) Given $3x^2 - 5x = -\frac{k}{4}$ calculate value(s) of k if
- (i) the real roots are real 1
 - (ii) the roots are rational and k is a positive integer. 2
- (c) To promote the sale of Studebaker cars, a dealer offers a special deal in which no interest is charged for the first 3 months and then interest rates are left at 1% per month. Lam Lai buys a 6-cylinder car for \$30 000, pays \$10 000 in cash and agrees to pay the loan plus interest monthly over 3 years. After 20 months, he wins \$10 150 as part of a lotto syndicate. Show that this win is just sufficient to pay off the loan at that time. 4

Question 6

- (a) Tap water at 24°C is placed in a fridge-freezer maintained at a temperature of -11°C . After t minutes the rate of change of temperature T of the water is given by $\frac{dT}{dt} = -k(T + 11)$
- (i) Show that $T = Ae^{-kt} - 11$ is a solution of the above equation, where A is a constant. 1
- (ii) Find the value of A 1
- (iii) After 15 minutes the temperature of the water falls to 10°C . Find to the nearest minute the time taken for the water to start freezing. (Freezing point of water = 0°C) 2
- (b) Evaluate $\int_{\frac{1}{3}}^{\frac{1}{2}} \frac{dx}{1+9x^2}$ 3
- (c) In the game of craps 2 dice are thrown and the sum of the dice is noted. The most likely outcome is a total of 7. If two dice are rolled 20 times,
- (i) What is the most probable number of sevens thrown? 3
- (ii) Calculate the probability that this number of sevens does indeed occur. 2

Question 7

- (a) A particle moves in a straight line and its position at any time is given by $x = 4.8 \cos 2t + 5.5 \sin 2t$. Show that the motion is simple harmonic and calculate its greatest speed. 3
- (b) Evaluate $\sin \left[\cos^{-1} \frac{2}{3} + \tan^{-1} \left(-\frac{3}{4} \right) \right]$ giving its exact value. 3
- (c) Consider the function $y = x \sec x$.
- (i) Find $\frac{dy}{dx}$ 1
- (ii) By drawing two graphs, show that the function has one stationary point in the domain $\frac{\pi}{2} < x < \frac{3\pi}{2}$. 2
- (iii) Prove that the stationary point lies between $x = 2.5$ and $x = 3.0$. 1
- (iv) Use halving the interval method twice to find a closer approximation of the stationary point. 2

Solutions

Question 1

(a) $3^{x+1} = 2$

$\log 3^{x+1} = \log 2$

$(x+1) \log 3 = \log 2$

$x+1 = \frac{\log 2}{\log 3}$

$x = \frac{\log 2}{\log 3} - 1$

$= -0.37$ (2 d.p.)

(b) $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} (1 + \sin^2 x) dx = \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} (1 + \frac{1}{2}(1 - \cos 2x)) dx$

$= \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} (\frac{3}{2} - \frac{\cos 2x}{2}) dx$

$= [\frac{3}{2}x - \frac{\sin 2x}{2}]_{\frac{\pi}{4}}^{\frac{\pi}{2}}$

$= \frac{\pi}{2} - \frac{\sqrt{3}}{2} - (\frac{3\pi}{8} - \frac{1}{4})$

$= \frac{\pi}{4} - \frac{\sqrt{3}}{2} + \frac{1}{4}$

$= \frac{\pi}{4} - \frac{\sqrt{3}-1}{2}$

(c) Domain $-1 \leq \frac{x}{2} \leq 1, -2 \leq x \leq 2$. Range $0 \leq y \leq \frac{\pi}{2}$.

(d) $P(x) = 6x^3 + 17x^2 - 4x - 3$

$P(-3) = -162 + 153 + 12 - 3 = 0$
 $\therefore x+3$ is a factor.

$$\frac{6x^3 - x - 1}{x+3} = \frac{6x^3 + 17x^2 - 4x - 3}{6x^3 + 18x^2}$$

$$\frac{-x^2 - 4x}{-x^2 - 3x}$$

$$\frac{-x-3}{-x-3}$$

Factors are $(x+3)(6x^2 - x - 1) = (x+3)(2x-1)(3x+1)$

(e) (i) $\tan \alpha = -\frac{1}{\sqrt{3}}$

$\tan \alpha = \tan(-\frac{\pi}{6})$ or $\tan \alpha = \tan(\frac{5\pi}{6})$

$\therefore \alpha = -\frac{\pi}{6} + n\pi$ or $\alpha = \frac{5\pi}{6} + n\pi$

(ii) $\alpha = -\frac{\pi}{6} - \pi$ or $\alpha = \frac{5\pi}{6} - 2\pi$
 $= -\frac{7\pi}{6}$ or $= -\frac{7\pi}{6}$

Question 2

(a) $\int (2-x^2)^{-1/2} dx$


$u = 2 - x^2$
 $du = -2x dx$
 $\therefore x dx = -\frac{du}{2}$

$\therefore \int = -\frac{1}{2} \int \frac{1}{u^{1/2}} du$

$= -\frac{1}{2} \frac{u^{-1/2}}{-1/2} + C$

$= \sqrt{2-x^2} + C$

(b)



Join AB:

$\angle ACT = \angle ABT$ (angles standing on the same arc in smaller circle)
 But $\angle ABT = \angle ADP$ (angles standing on the same arc in larger circle)

$\therefore \angle ACT = \angle ADP$

$\therefore CT \parallel PD$ (alternate angles are equal)

(c)

$\frac{2x-5}{x-4} \geq x$
 $(2x-5)(x-4) \geq x(x-4)^2$

$(2x-5)(x-4) - x(x-4)^2 \geq 0$

$(x-4)[2x-5-x(x-4)] \geq 0$

$(x-4)(2x-5-x^2+4x) \geq 0$

$(x-4)(-x^2+6x-5) \geq 0$

$(x-4)(-5+x)(1-x) \geq 0$

Test $x = 0$ valid $\therefore x \leq 1$ or $4 < x \leq 5$ (d) Let $y = 3^{\sqrt{x+1}}$ Put $u = \sqrt{x+1}$. $y = 3^u$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = \ln 3 (3^u) \times \frac{1}{2} (x+1)^{-\frac{1}{2}} = \frac{\ln 3 (3^{\sqrt{x+1}})}{2\sqrt{x+1}}$$

(e) $v = \frac{1}{x+3}$ i.e., $\frac{dv}{dx} = -\frac{1}{(x+3)^2}$ $\frac{dv}{dx} = x+3$

$$t = \int (x+3) dx = \frac{x^2}{2} + 3x + c$$

When $t = 0$, $x = 1$; $0 = \frac{1}{2} + 3 + c \therefore c = -\frac{7}{2}$

$$\therefore t = \frac{x^2}{2} + 3x - \frac{7}{2}$$

When $x = 5$, $t = \frac{25}{2} + 15 - \frac{7}{2} = 24$

Question 3

(a) Test $n = 4$: LHS = $2(4-3) = 2$

$$\text{RHS} = \binom{5}{2} - 4 = 6 - 4 = 2$$

 \therefore true for $n = 4$.Suppose that an integer k exists for which the result is true i.e., $2(k-3) + (k-4) + \dots + 3 + 2 + 1 = \binom{k}{2} - k$ Consider when $n = k+1$.

$$\therefore \text{RHS} = 2(k-2) + (k-3) + (k-4) + \dots + 3 + 2 + 1$$

$$= 2(k-2) + 2(k-3) + (k-4) + \dots + 3 + 2 + 1 - (k-3)$$

$$= \binom{k}{2} - k + 2(k-2) - (k-3)$$

$$= \binom{k}{2} - k + k - 1$$

$$= \binom{k}{2} - 1$$

$$\text{But } \binom{k}{2} = \binom{k+1}{2} - \binom{k}{2-1}$$

$$\therefore \binom{k}{2} = \binom{k+1}{2} - \binom{k}{1}$$

$$\therefore \text{LHS} = \binom{k+1}{2} - \binom{k}{1} - 1$$

$$= \binom{k+1}{2} - k - 1$$

$$= \binom{k+1}{2} - (k+1)$$

So the result is true for $n = k+1$.But it is true for $n = 4$. \therefore it is true for $n = 4 + 1 = 5$ and by mathematical induction it is true for all positive integers $n \geq 4$.(b) Let the roots of the equation be α, β and $\alpha - \beta$.Sum of roots = $\alpha + \beta + \alpha - \beta = 2\alpha = -\frac{1}{4} = \frac{1}{4} \therefore \alpha = \frac{1}{8}$ Product of all 3 roots = $\alpha\beta(\alpha - \beta) = -\frac{1}{8}$

$$\frac{1}{8}\beta\left(\frac{1}{8} - \beta\right) = -\frac{1}{8}$$

$$\frac{1}{8}\beta - \frac{1}{8}\beta^2 + \frac{1}{8} = 0$$

$$2\beta^2 - \beta - 15 = 0$$


$$(2\beta + 5)(\beta - 3) = 0$$

$$2\beta + 5 = 0 \text{ or } \beta = 3$$

$$\beta = -\frac{5}{2}, \beta = 3$$

$$\therefore \text{roots are } \frac{1}{8}, 3, -\frac{5}{2}$$

(c)

(i) Equation of tangent at P : $y = px - ay^2$ It cuts the x -axis when $y = 0$, i.e., $x = ap$; $A = (ap, 0)$ It cuts the y -axis when $x = 0$, i.e., $y = -ap$; $B = (0, -ap^2)$

$$\therefore M = \left(\frac{ap}{2}, -\frac{ap^2}{2}\right)$$

(ii) From $x = \frac{ap}{2}$, $p = \frac{2x}{a}$

$$\text{Sub in } y = -\frac{ap^2}{2}$$

$$y = -\frac{a}{2} \left(\frac{2x}{a}\right)^2$$

$$y = -\frac{2x^2}{a}$$

i.e., $x^2 = -\frac{1}{2}ay$ which is the equation of a parabola.

$$\text{(iii) } x^2 = -\frac{1}{2}ay$$



$$= 4\left(-\frac{1}{8}a\right)y$$

Its focal length is $\frac{a}{8}$

\therefore coordinates of focus are $(0, -\frac{1}{2}a)$
Equation of directrix: $y = \frac{1}{2}a$

Question 4


- (a) (i) $N = \frac{11}{21} = 9\ 979\ 200$
(ii) Consider the letters BABY to be one item, so $N = \frac{8!}{2!} = 20\ 160$
- (b) In a vertical plane:



$$\begin{aligned}\tan 40^\circ &= \frac{80}{a} \\ a &= \frac{80}{\tan 40^\circ}\end{aligned}$$

$$\begin{aligned}\tan 36^\circ &= \frac{120}{b} \\ b &= \frac{120}{\tan 36^\circ}\end{aligned}$$

On a horizontal plane:



$$\begin{aligned}\cos \theta &= \frac{(80)^2 + (120)^2 - (150)^2}{2(80)(120)} \\ \theta &= 63^\circ 52'\end{aligned}$$

- (c) $x(1+x)^n = x({}^nC_0 + {}^nC_1x + {}^nC_2x^2 + \dots + {}^nC_nx^n)$
Differentiating both sides of the equation gives
 $(1+x)^n + nx(1+x)^{n-1} = {}^nC_0 + {}^nC_1x + {}^nC_2x^2 + \dots + {}^nC_nx^n$
 $+ x({}^nC_1 + 2{}^nC_2x + \dots + n{}^nC_nx^{n-1})$
 $= {}^nC_0 + 2x{}^nC_1 + 3x^2{}^nC_2 + \dots + (n+1)x^n{}^nC_n$
Put $x = 1$

$$\therefore 2^n + n(2^{n-1}) = {}^nC_0 + 2{}^nC_1 + 3{}^nC_2 + \dots + (n+1){}^nC_n$$

$$2^n + \frac{n(2^n)}{2} = \sum_{r=0}^n (r+1) \cdot {}^nC_r$$

i.e., $2^n(\frac{3}{2} + 1) = \sum_{r=0}^n (r+1) \cdot {}^nC_r$

Question 5

- (c) (i) $y = vt \sin 40^\circ - 5t^2$

(ii) $x = vt \cos 40^\circ$

(iii) From (ii) $t = \frac{x}{v \cos 40^\circ}$
 $y = \frac{v \sin 40^\circ x}{v \cos 40^\circ} - 5 \left(\frac{x}{v \cos 40^\circ} \right)^2$
 $= x \tan 40^\circ - \frac{5x^2}{v^2 \cos^2 40^\circ}$
 $\therefore y = x \tan 40^\circ - \frac{5x^2}{v^2} (1 + \tan^2 40^\circ)$

(iv) $x = 20\text{m}, y = 6\text{m}$
 $\therefore 6 = 20 \tan 40^\circ - \frac{5(20)^2}{v^2} (1 + \tan^2 40^\circ)$

$$\begin{aligned}\therefore v^2 &= \frac{5(20)^2(1 + \tan^2 40^\circ)}{20 \tan 40^\circ - 6} \\ \therefore v &= 17.8\text{m/s}\end{aligned}$$

(b) $3x^2 - 5x = -\frac{1}{3}$
 $12x^2 - 20x + k = 0$

(i) For real roots $\Delta \geq 0$
 $b^2 - 4ac \geq 0$
 $400 - 4(12)k \geq 0$
 $25 - 3k \geq 0$
 $k \leq \frac{25}{3}$

(ii) For rational roots Δ is a perfect square i.e., $25 - 3k$ is a perfect square. Since k is a positive integer, less than $\frac{25}{3}$, the only possible values of $25 - 3k$ are 0, 1, 4, 9, 16, 25, yielding solutions $k = 3, 7$ and 8.

- (c) Let P = monthly repayment.
 A_n = amount owing after n months.

$$\begin{aligned}
 A_1 &= 20\,000 - P \\
 A_2 &= 20\,000 - 2P \\
 A_3 &= 20\,000 - 3P \\
 A_4 &= A_3(1.01) - P \\
 &= (20\,000 - 3P)(1.01) - P \\
 A_5 &= [(20\,000 - 3P)(1.01) - P](1.01) - P \\
 &= (20\,000 - 3P)(1.01)^2 - P(1 + 1.01) \\
 A_6 &= (20\,000 - 3P)(1.01)^3 - P(1 + 1.01 + 1.01^2) \\
 A_{36} &= (20\,000 - 3P)(1.01)^{36} - P(1 + 1.01 + 1.01^2 + \dots + 1.01^{35}) \\
 0 &= (20\,000 - 3P)(1.01)^{36} - P \frac{1.01^{36} - 1}{1.01 - 1} \\
 (1.01^{36} - 1)P &= 0.01(1.01)^{36} 20\,000 - 3P(0.01)(1.01)^{36} \\
 P[1.01^{36} - 1 + 3(0.01)(1.01)^{36}] &= 0.01(1.01)^{36} \times 20\,000 \\
 P &= \frac{0.01(1.01)^{36} \times 20\,000}{1.01^{36} - 1 + 3(0.01)(1.01)^{36}} \\
 &= \frac{0.01(1.01)^{36} \times 20\,000}{1.01^{36}(1 + 0.03) - 1} \\
 &\approx \$645.38 \\
 A_{30} &= (20\,000 - 3P)(1.01)^{30} - \frac{P(1.01^{30} - 1)}{1.01 - 1} \\
 \text{Sub } P &= 645.38 \\
 A_{30} &= (20\,000 - 3(645.38))(1.01)^{30} - \frac{645.38(1.01^{30} - 1)}{0.01} \\
 &= 10\,146.70
 \end{aligned}$$

\therefore \$10 150 is just sufficient to pay off the loan at 20 months.

Question 6


- (a) (i) $T = Ae^{-kt} - 11$ ($Ae^{-kt} = T + 11$)
 $\frac{dT}{dt} = -kAe^{-kt}$
 $= -k(T + 11)$
- (ii) When $t = 0$, $T = 24^\circ$
 $\therefore T = Ae^{-kt} - 11$
 $24 = Ae^0 - 11$
 $A = 35$
- (iii) $T = 35e^{-kt} - 11$
 When $t = 15$ min, $T = 10$
 $\therefore 10 = 35e^{-15k} - 11$
 $e^{-15k} = \frac{21}{35} = 0.6$
 $-15k = \ln(0.6)$
 $k = \frac{\ln(0.6)}{-15} \approx 0.034$

$$\begin{aligned}
 \therefore T &= 35e^{-0.034t} - 11 \\
 \text{When } T &= 0 \\
 0 &= 35e^{-0.034t} - 11 \\
 e^{-0.034t} &= \frac{11}{35} \\
 t &= \frac{\ln(0.6)}{-0.034} \\
 &\approx 33.98
 \end{aligned}$$

≈ 34 min to the nearest minute.

(b) $\int_0^{\frac{1}{2}} \frac{dx}{1+x^2} = \int_0^{\frac{1}{2}} \frac{1}{1+x^2} dx$
 $= \frac{1}{2} \int_0^{\frac{1}{2}} \frac{1}{1+x^2} dx$
 $= \frac{1}{2} [\tan^{-1} x]_0^{\frac{1}{2}}$
 $= \frac{1}{2} (\tan^{-1} \frac{1}{2} - \tan^{-1} 0)$
 $= \frac{1}{2} (\frac{\pi}{6} - 0) = \frac{\pi}{12}$

(c)



- (i) Consider $(q + p)^{20}$ i.e., $(\frac{7}{8} + \frac{1}{8})^{20}$
 To find greatest coefficient in this expansion,
 $\frac{C_r^{20}}{C_{r-1}^{20}} = \frac{20-r+1}{r} \cdot \frac{1}{7} \geq 1$
 $\frac{21-r}{r} \cdot \frac{1}{7} \geq 1$
 $21 - r \geq 7r$
 $6r \leq 20$
 $r \leq 3$

i.e., most probable number of total of 7 is 3.

- (ii) Consider $(q + p)^{20} = \sum_{r=0}^{20} {}^{20}C_r q^r p^{20-r}$
 $r = 3$ produces
 $P(E) = {}^{20}C_3 (\frac{1}{8})^3 (\frac{7}{8})^{17}$
 ≈ 0.238

Question 7

(a) $x = 4.8 \cos 2t + 5.5 \sin 2t = A \sin(2t + \alpha)$

$A = \sqrt{(4.8)^2 + (5.5)^2} = 7.3$

$\tan \alpha = \frac{4.8}{5.5} \quad (0 < \alpha < \frac{\pi}{2})$

$\alpha = 0.72^\circ$

$\therefore x = 7.3 \sin(2t + 0.72)$

$\dot{x} = 2(7.3) \cos(2t + 0.72)$

$\ddot{x} = -2^2(7.3) \sin(2t + 0.72)$

$= -2^2 x$

Since $\ddot{x} = -n^2 x$, the motion is simple harmonic.

OR: $x = 4.8 \cos 2t + 5.5 \sin 2t$

$\dot{x} = -2(4.8) \sin 2t + 2(5.5) \cos 2t$

$\ddot{x} = -2^2(4.8) \cos 2t - 2^2(5.5) \sin 2t$

$= -2^2(4.8 \cos 2t + 5.5 \sin 2t)$

$= -2^2 x$

The speed is greatest when $\ddot{x} = 0$

i.e., when $-4(7.3) \sin(2t + 0.72) = 0$

$2t + 0.72 = 0, \pi, 2\pi, \dots$

The smallest positive value of t for which the speed is a maximum

is given by $2t + 0.72 = \pi$

$\dot{x} = 2(7.3) \cos \pi$

$= -14.6$

The maximum speed has magnitude 14.6.



(b) $\sin[\cos^{-1} \frac{3}{5} + \tan^{-1}(-\frac{3}{4})]$

Let $\cos^{-1} \frac{3}{5} = \alpha, 0 \leq \alpha \leq \pi$

and $\tan^{-1}(-\frac{3}{4}) = \beta, -\frac{\pi}{2} < \beta < \frac{\pi}{2}$

$\therefore \cos \alpha = \frac{3}{5}$ and $\tan \beta = -\frac{3}{4}$

α may be represented as an angle in the first question and β may be represented as an angle in the fourth quadrant.



$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$

$= \frac{\sqrt{5}}{5} \cdot \frac{4}{5} + \frac{3}{5}(-\frac{3}{5})$

$= \frac{4\sqrt{5}-9}{25}$

(c) (i) $y = x \sec x$

$\frac{dy}{dx} = \sec x + x \sec x \tan x$


(ii) $\sec x + x \sec x \tan x = 0$ for stationary points

$$\sec x(1 + x \tan x) = 0$$

$$\sec x \neq 0, 1 + x \tan x = 0$$

$$x \tan x = -1$$

$$\tan x = -\frac{1}{x}$$



(iii) Let $f(x) = 1 + x \tan x$

$$f(2.5) = -0.867... < 0$$

$$f(3.0) = 0.572... > 0$$

\therefore change in sign between 2.5 and 3 so that stationary point lies between 2.5 and 3.

(iv) Consider $x = 2.75$

$$f(2.75) = -0.1355... < 0$$

\therefore stationary point lies between 2.75 and 3.

Consider $x = 2.875$

$$f(2.875) = 0.2148... > 0$$

\therefore stationary point lies between 2.75 and 2.875

\therefore closer approximation of the stationary point is $\frac{2.75+2.875}{2} = 2.8125$

Marking Guidelines

1 (a)	1	Log form
1 (b)	1	Correct answer, 2d.p.
1 (c)	1	Conversion to multiple angle
1 (d)	1	Integration
1 (e)	1	Exact value
2 (a)	1	Domain
2 (b)	1	Range
2 (c)	1	First factor
2 (d)	1	Second factors by long division
2 (e)	1	Factorisation
3 (a)	1	General solution
3 (b)	1	Specific value
3 (c)	1	Substitution
3 (d)	1	Integration
3 (e)	1	Statement with reason
4 (a)	1	Second statement and reason
4 (b)	1	Conclusion
4 (c)	1	Various methods
4 (d)	1	Testing a value or graphical method
4 (e)	1	Solution
5 (a)	1	Chain rule
5 (b)	1	Answer
5 (c)	1	1 in terms of x
5 (d)	1	Substitution and answer
6 (a)	1	Test $m=4$
6 (b)	1	Integer k
6 (c)	1	$m \neq 1$ and algebra
6 (d)	1	Conclusion
6 (e)	1	Sum of roots
7 (a)	1	Product of roots
7 (b)	1	Solution
7 (c)	1	Coordinates of A and B
7 (d)	1	Coordinates of M
7 (e)	1	Elimination of parameter
8 (a)	1	Conclusion
8 (b)	1	Focal length
8 (c)	1	Correct answer
8 (d)	1	Correct answer
8 (e)	2	Triangles in vertical plane
9 (a)	1	Triangle on horizontal plane
9 (b)	1	Expressions for a and b
9 (c)	1	Cosine rule
9 (d)	1	Correct angle (nearest minute)
9 (e)	1	Expansion
10 (a)	1	Differentiation
10 (b)	1	$x=1$
10 (c)	1	Algebra and answer

5 (a)	(i)	1	Vertical distance
	(iii)	1	Horizontal distance
	(iii)	1	Derivation
	(iv)	1	Substitution
		1	Correct answer
(b)	(i)	1	k for real roots
	(ii)	1	k for rational roots
		1	Integral values
(c)		1	A_1-A_3
		1	A_36
		1	Value of instantant
		1	A_20 and conclusion
6 (a)	(i)	1	Differentiation
	(ii)	1	Value of A
	(iii)	1	Value of k
		1	Time taken
(b)		1	Adjustment
		1	Integration
		1	Evaluation
(c) (i)		1	P(total of 7)
		1	Greatest coefficient
		1	Answer
	(ii)	2	Binomial probability
7 (a)		1	Transformation
		1	Differentiation and conclusion
(b)		1	Maximum speed
		2	Two triangles
		1	Compound angle
(c) (i)		1	Differentiation
	(ii)	1	Equation
		1	Graphs
(iii)		1	Proof
(iv)		2	Halving the interval twice

Mathematics Extension 1

HSC TRIAL EXAMINATION MAPPING GRID

Question/Metric	Content	Syllabus Outcome	Targeted Performance Bands
1(a)	2 Logarithmic and Exponential functions	PS, H9	E2-E3
1(b)	3 Integration; Trigonometric functions	H8	E2-E3
1(c)	2 Inverse Trigonometric functions	HE4	E2-E3
1(d)	3 Polynomials	PE3	E2-E3
1(e)(i)	1 Trigonometric functions	P4	E2-E3
(ii)	1 Trigonometric functions	P4	E2-E3
2(a)	2 Integration	HE3	E2-E3
(b)	3 Circle Geometry	PE2, PE3	E2-E3
(c)	3 Inequalities	PE3	E2-E3
(d)	2 Derivative of a function	P7, H5	E2-E3
(e)	2 Applications of Calculus	HE3	E3-E4
3(a)	4 Mathematical Induction	HE2	E3-E4
(b)	3 Polynomials	PE3	E2-E3
(c)(i)	2 Parametric representation	PE4	E2-E3
(c)(iii)	2 Parametric representation	PE4	E2-E3
(c)(iii)	1 Parametric representation	PE4	E2-E3
4(a)(i)	1 Permutations and Combinations	HE3	E2-E3
(ii)	1 Permutations and Combinations	HE3	E2-E3
(b)	6 Trigonometric functions	PE3	E2-E3
(c)	4 Binomial Theorem	PE3, PE6, HE7	E2-E4
5(a)(i)	1 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
(ii)	1 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
(iii)	1 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
(iv)	2 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
5(b)(i)	1 Quadratic theory	PE3	E2-E3
(ii)	2 Quadratic theory	PE3	E2-E4
(c)	4 Series and Applications	H5, H9	E2-E4
6(a)(i)	1 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
(ii)	1 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
(iii)	2 Applications of Calculus to the Physical World	HE3, HE7	E2-E3
(b)	3 Integration; Inverse Trigonometric functions	H8	E3-E4
(c)(i)	3 Binomial Theorem	PE3, PE6	E2-E4
(ii)	2 Binomial probability	HE7	E2-E4
7(a)	3 Applications of Calculus to the Physical World	HE3, HE7	E2-E4
(b)	3 Inverse Trigonometric functions	H8	E3-E4
(c)(i)	1 Differentiation of trigonometric functions	P4, H5	E2-E3
(ii)	2 Differentiation of trigonometric functions	PE2	E3-E4
(iii)	1 Differentiation	PE2	E2-E3
(iv)	2 Polynomials	PE3	E3-E4