Candidate's Number. Math's teacher

## SYDNEY TECHNICAL HIGH SCHOOL (Est. 1911)



# 2000 TRIAL HIGHER SCHOOL CERTIFICATE

## MATHEMATICS

#### 2 UNIT

Time Allowed - Three hours (plus 5 minutes reading time)

## DIRECTION TO CANDIDATES:

- All questions may be attempted
- All questions are of equal value

Approximate marks are shown

- All working should be shown in every question.
- Full marks may not be awarded for careless of badly arranged
- Standard Integrals are printed on the last page of this examination
- Approved calculators may be used
- name and class on the top right hand side of the page clearly marked with the number of the question and your Each question attempted is to be started ON A NEW PAGE

conclusion of the examination This question paper must be handed in with your answers at the


QUESTION	Mark
Question 1	
Question 2	
Question 3	
Question 4	
Question 5	
Question 6	
Question 7	
Question 8	
Question 9	
Question 10	
FOTAL	

- a) Factorise  $4x^2 25$
- 1 b) Find log<sub>3</sub>27
- 2 c) Find sin 1.3 radians correct to 3 decimal places.
- 2 d) Simplify  $\frac{x}{3} \frac{3x 1}{10}$
- 2 e) If  $\sqrt{45} + \sqrt{A} = 5\sqrt{5}$  find A.
- 2. f) Solve 4-3x < 9
- 2 g) For the parabola  $x^2 = -8(y+2)$  find
- i) co-ordinates of the vertex.
- ii) focal length

#### QUESTION 2

- 6 a) Differentiate the following
- i)  $(x^2 6)^4$
- ii) xtanx
- iii)  $\sqrt[3]{x}$   $\begin{cases} 6 \\ \sqrt{x-1} \\ \sqrt{x} \end{cases} dX \text{ (exact value)}$
- s c) i) Write down the discriminant of  $2x^2 kx + k$
- ii) For what values of k does  $2x^2 kx + k$  have real roots?


- A , B and C are the points ( 1,0 ), ( 3,2 ) and ( -1,8 ) respectively
- i) Draw a diagram to show the points A, B and C.
- ii) Find the length BC.
- iii) Show that the equation of the line BC is 3x + 2y - 13 = 0
- iv) Find the perpendicular distance from A to the line BC
- v) Find the area of the triangle ABC.
- **₹**. If ABCD is a parallelogram write down the co-ordinates of D.
- ¥: BD meet. Find the co-ordinates of the point where the diagonals AC and



and AC was 5 metres while AC was 40 metres and  $\angle ACB = 60^{\circ}$ A horizontal bridge joins points X and Y. Cables were used to support the bridge as shown in the diagram. The distance between the cables AB

- i) Find the exact height of A above the bridge
- ii) Use the cosine rule to find the length of the cable AB


7 a)



The diagram shows the graph of  $y = \frac{1}{1}$ 


- The curve cuts the y-axis at A. Find the co-ordinates of A.
- ii) Show that the gradient of the normal at A is 1
- iii) Find the equation of the normal at A.
- The normal cuts the curve again at B. Find the co-ordinates of B. iv)
- The number of cubes in the side of each square layer forms an Arithmetic Progression. The first 3 terms are 80, 76, 72...... A tower is made by adding layers of cubes. Each layer is a square b) 10
- i) Find the eighteenth term of the sequence.
- ii) Find the sum of the first eighteen terms.
- If 685,580 cubes are available, is it possible to build this eighteen layer tower?  $\Xi$

2



- <u>a</u> Find the value of x, giving reasons. In this diagram PTLRS. Copy the diagram onto your page.
- w ত is rotated about the x axis. Find the volume when the curve  $y = e^{3x}$  between x = 0 and x = 2
- w C spray it delivers 15 mL and the user continues to spray. When used it only delivers 90% of its previous spray. On its first A bottle of cleaner fluid has a spray nozzle which is defective.
- i) On the fifth spray how many mL's does it deliver?
- ii) What is the total limiting amount the spray can deliver.
- Given  $\log 6 = x$  and  $\log 2 = y$ . Find in terms of x and y
- i) log 3
- ii) log 24
- iii) Evaluate  $\frac{2x}{y}$  to 2 decimal places.

5 a)



The table shows the values of f(x) for 4 values of x for the graph y = f(x)


x	0	1	2	3
f(x)	1	15	5	2

Use the Trapezoidal Rule and the 4 function values to evaluate 

$$\int_{0}^{3} f(x) dx$$

From the diagram, decide if this approximation is an over-estimate or an under-estimate of the true value of the integral. Give a brief reason.  $\Xi$ 


7 b)



This is a glaziers plan for a window.DC is an arc of a circle centre M with radius 120 cm. ABCD is a rectangle where AM=MB=60 cm

- i) Find the size of ∠DMA
- ii) Find the size of *LDMC*
- iii) Find the length of arc DC.
- iv) Find the total area of glass needed for the window.

6 a)



In the diagram ZS.LXY and XTLYZ. Redraw the diagram on your paper.


- i) Explain why  $\angle YXT = \angle SZY$
- ii) Prove  $\Delta TXY$  is similar to  $\Delta TZM$
- iii) If XT = 8 cm, YT = 5 cm and SZ bisects XT. Find the length of TZ.
- 0 9 The volume of water in a tank with capacity 400 litres is controlled by two pumps. When pump A operates the volume of water in the tank is given by  $V = 400 - 4t - \frac{1}{5}$  (t is in minutes)
- i) Find the volume after 20 minutes.
- Ξ) Find the rate of flow of water after 20 minutes. Explain your answer.

At this time (20 minutes) pump A stops and pump B begins. Pump B fills the tank at a rate given by  $\frac{dV}{dt} = \frac{t}{10} + 5$ 

iii) How much water will be in the tank 10 minutes after pump B starts.

- The population of insects is given by  $P = 6000e^{0.3t}$  where t is in days <u>a</u> 3
- i) Find the population when t=5.
- ii) How long does it take for the population to reach 50000.

**O** 



The diagram shows parts of the curves y=sinx and y=cos2x

- i) Show that the curves intersect at  $x = \frac{\pi}{6}$
- ii) Calculate the shaded area. Leave the answer in exact form.
- is  $C = 0.04v^2 + \frac{17500}{1000} + 275$  dollars, where v is the average speed A bus company has established that the cost of a 1000 km journey in kilometres per hour. At what speed should the buses travel to achieve a minimum cost. ত

4

N <u>a</u> The weight w, in kilograms, of a new baby in the first few weeks is given in the following table.

weight (w)	Time(t)
0.6	1st week
0.8	2nd
1.05	3rd
1.35	4th
1.8	5th
2.5	6th

Use the information to find the signs of

- $\frac{dw}{dt}$
- ii)  $\frac{d^2w}{dt^2}$
- 10 b) For the curve  $y = x^3 3x^2$
- Find the co-ordinates of any stationary points and determine their nature.
- ii) Find the co-ordinates of any points of inflexion.
- III) points of inflexion and intercepts. Draw a neat sketch of the curve indicating stationary points,

Use your graph or otherwise to state the values of x for which the curve is

- iv) increasing
- v) concave down
- <u>¥</u>. The curve and the line y = -kx + 3k intersect. For what values of k will there be only one point of intersection.

3 a) If 
$$\int_{1}^{p+1} 3x^2 dx = 19$$
 Find the values of p

9

- cross a sandbar and his boat requires water at least 2 metres deep On a particular day high tide occurred at midnight and the depth (D) of water over the sandbar on the ensuing day is given by the the ocean tides. To reach the open sea from his mooring he must A fisherman moors his boat in a river which is heavily effected by formula  $D = 2\cos\frac{4\pi}{25}t + 1$  where t is the number of hours after midnight. **P**
- i) What is the depth of the water at high tide
- ii) What is the depth of the water at low tide. Explain your answer.
- iii) What time elapses between high tides.
- iv) Use the above information to draw the graph of  $D = 2\cos\frac{4\pi}{25}t + 1 \text{ for } 0 \le t \le 24.$
- Draw another line on your graph and using the two graphs indicate the times of the day (on your graph) when he would not be able to cross the sandbar in his boat **\( \)**

### STANDARD INTEGRALS

$$x^n dx$$

$$= \frac{1}{n+1} x^{n+1}, \ \ n \neq -1; \ \ x \neq 0, \ \text{if } n < 0$$

$$\int_{x}^{1} dx$$

$$=\ln x, x>0$$

$$e^{ax} dx$$

$$=\frac{1}{a}e^{ax}, a\neq 0$$

$$= \frac{1}{a} \sin ax, \quad a \neq$$

$$a = \frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax \, dx$$

$$= \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx$$

sec ax tan ax dx

 $= \frac{1}{a} \sec ax, \quad a \neq 0$ 

$$= \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx$$

$$=\sin^{-1}\frac{x}{a}$$
,  $a > 0$ ,  $-a < x <$ 

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx$$

$$=\ln\left(x+\sqrt{x^2-a^2}\right), \quad x>a>0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx$$


$$= \ln\left(x + \sqrt{x^2 + a^2}\right)$$

NOTE: 
$$\ln x = \log_e x$$
,  $x > 0$ 



一( 1) (1) (0,4)	
of diagonal	" 5 - lab
Vii) diagonals meeta much por	= 6. lub - 1
VI 711 (13 6)	
11 100 5	
v) Area = 1×15×10	,
	(ii) d x 1/3 = 1/3 x - 2/3
,,  ō	
132422	ii) d xtoxx = tanx:1+ x. me'x
iv   d=   3x1 +2x0 - 131	
3x+2y-13=0	111日(ストー6) 4 = 4.2x. (ストー6)
29-4 = -32+9	
: y-2 =-3/(z-3)	
1 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	j) i) (0,-2)
iii ) mBC = 8-2	27-5/3
	13 * \ 5
ii) * 3C= [6].	t) 4-3x 19
-1 + - 3	A * 20
A	2 215
, D	× 515 - 3J5
	1 JA = 5.15-145
	30
Question 3	9.
And the same of th	± 30x-92+3
4×0. 4×8	ō
	1 2 1 32-1
: 22-84 >0	
Ò	.) 0.96355
3-	) 3
c);) A = 2 - 4.2.4	(22-5)(22+5)
	Fucstim!


Question S $T$ $Z$	$b_1 V_{\pm} = 32.$ $b_1 V_{\pm} = \left[ \frac{2}{4} \left( e^{3x} \right)^2 dx \right].$ $= \left[ \frac{1}{4} \left( \frac{e^{6x}}{6} \right)^2 \right].$ $= \left[ \frac{1}{4} \left( \frac{e^{6x}}{6} \right)^2 \right].$	$= (e^{12} - 1) \frac{\pi}{6} \text{ or equivalent}.$ $c) i) T_{S} = 15 \times (0.9)^{4}$ $ii) limiting som = \frac{15}{1-0.9}$ $ii) limiting som = \frac{15}{1-0.9}$	d) i) log 3 = log 6 - log 2  ii) log 24 = log 6 + log 4  iii) log 24 = log 6 + 2 log 2  = log 6 + 2 log 2  = 2 log 6  = 2 log 6  = log 32  = log 4  = 5.17 (2010)
3) i) sin $60 = AD$ $\frac{40}{40}$ $\frac{2}{10}$ $\frac{40}{10}$ $\frac{1}{10}$	11) x=0 y= -1.  11) y= -1.(x-1)-2  24(0,-1)  24(0,-1)	14	



 $\int_{0}^{\infty} f(x) dx = \frac{3-0}{6} + 43$ 

6

Underestimate. The first trapezium's area is much less than the actual area.



III NAMO = 60

iii) Arc = 120 x II (125.7)

tan 60 = AD AD= 60 tan 60


Areu: 60 × 60 tan 60 + 1 × 120 × 11

= 3600 d 3 + 2400 H

3.775 cm²

Question 7

\$247 = 1×47 ... 47- 0b = \$24 7 47- 0b = 1×47



Since trangles are simila

x = 6.4cm

67 is \$= 20 V= 400-4×20-202	ii) The Area = ( Sinx - Coolx clx
ii dv = -4 - 2 to dt = -4 - 2 to	$\frac{\int_{\pi/\epsilon}}{\pi/\epsilon} = \left[ -\cos x - \ell, \sin 2x \right]^{\pi/\epsilon}$
) · "	= [evox + 1/2 sn 2x ] #7/2
Lewins .	
11. alv = t +5	
1	c) C= 0.04v2+17500+275
t=0 c=240	dc = 0.08v - 17500
:. V = £2 + S & +240	/ \ <b>∀</b> _
4=10 CO + Sx10 + 240	٠٠ ٥٠٥٤١ - ١٦٥٥٥ - ١
σ	0.087) = 4780.0
3 ter. 8	60.
a) i)   = 6000 e	08 09 05 1
ii) 50000 = 6000 c 0.3 t 50 = c 0.3 t	av I of the
η.	speed is 60 km/hr.
shop	Questiona
b) i) x = 17/6 sum 17/2 = 0.5	al an
	1) du >0
	ii) dry >0

