

# Cell Physiology lecture #5

Andrew Moorhouse;  
a.moorhouse@unsw.edu.au; 9385 2575

## Learning Aims – lecture 4

- To understand the forces that dictate how an ion moves across the cell membrane, including being able to describe electrochemical equilibrium and to predict how an ion moves across the membrane given the membrane potential and equilibrium potential.
- To know the Nernst equation and be able to use it (if given the values of the constants) to calculate equilibrium potentials
- To be able to describe how the resting membrane potential is generated and how changes in membrane potential occur (in terms of ion flows)
- Define depolarization, hyperpolarization, electrochemical equilibrium

### Supporting activities


- Membrane Potentials Virtual Prac (via Moodle)
- Stanfield, Chapter 7.3, start of 7.4.

## Recap: How membrane potentials are generated

A nice set of simulations to play with @ Francisco Bezanilla's home page :

<http://nerve.bsd.uchicago.edu/>

(use explorer not chrome & needs to allow JAVA to run)



## Recap: How membrane potentials are generated

When ions move across membranes they change the membrane potential. To move across a membrane their must be 1) a force and 2) the membrane must be permeable. So the relevant physiology is about:

### Electrochemical gradients


- Chemical force (Diffusion)
- Electrical Force (Electrostatics)

### Membrane Permeability

- What type of ion channels are open?

## The electrochemical forces for ions in physiological conditions when a cell is at rest


- The difference between the equilibrium potential ( $E_x$ ) and the membrane potential ( $V_m$ ) is the driving force for movement of ion x



Copyright © 2002, Elsevier Science (USA). All rights reserved.


## The nerve action potential....

1. Concept of Threshold
2. Features / Phases
3. Ionic Currents involved
4. Ion Channels involved
5. Molecular Physiology
6. Voltage-dependent Gating



## The nerve action potential.... Underlying ionic Currents.


- The 1<sup>st</sup> nerve action potential recording, from inside the giant axon of a squid



See also <https://www.youtube.com/watch?v=k48jXzFGMc8> for video of this early experiment  
Hodgkin & Huxley, *Nature*, 1939, *J Physiol.*, 1945

## The nerve action potential....

- The 1<sup>st</sup> nerve from inside the g



See also <https://www.youtube.com/watch?v=k48jXzFGMc8> for video of this early experiment  
Hodgkin & Huxley, *Nature*, 1939, *J Physiol.*, 1945



## The nerve action potential.... Underlying ionic Currents.




Sir Alan Hodgkin  
Nobel Prize 1963



Sir Bernard Katz  
Nobel Prize 1971

- One of the 1<sup>st</sup> Action Potential experiments....

*J. Physiology*, 1949



- 1 = control,
- 2 = low  $\text{Na}^+$  in the sea water,
- 3 = wash / recovery response

Conclusion?:



## The nerve action potential.... Underlying ionic Currents.

1. Inward  $\text{Na}^+$  current,  
•  $\text{Na}^+$  influx



**Rapid  
depolarization**

2. Outward  $\text{K}^+$  current  
•  $\text{K}^+$  efflux



**Slow  
repolarization**



## The nerve action potential.... Underlying ionic Currents and the Channels involved.

1. Inward  $\text{Na}^+$  current,

- $\text{Na}^+$  influx
- Fast, voltage gated channels



**Rapid  
depolarization**

2. Outward  $\text{K}^+$  current

- $\text{K}^+$  efflux
- Slow, voltage gated channels



**Slow  
repolarization**




# The molecular physiology of the nerve action potential

- some more important contributors.....


- 1) The giant squid
- 2) Hodgkin and Huxley
- 3) ....
- 4) ....

(Biology provides some great insights into physiology and opportunities for discoveries)




# The molecular physiology of the nerve action potential

- From the 1<sup>st</sup> K<sup>+</sup> channel gene from a fruit fly a family of different types of K<sup>+</sup> ion channels genes were discovered (in mammals). The structures and properties have been extensively. Different families have similar functional properties but with subtle differences to give diversity.



# The molecular physiology of the nerve action potential

- Voltage-dependent activation...
- All voltage-dependent channels have a voltage sensor that detects the membrane potential and acts as a molecular switch and opens (or activates) the channels at a specific voltage level or threshold



The voltage sensor is a region of the channel protein (containing charged amino acids) that can detect and respond to changes in the membrane potential. The images suggest different models on how it may move in response to membrane potential changes. This movement initiates opening of the gate when the voltage threshold is reached.









## The molecular physiology of the nerve action potential

- Voltage-dependent *inactivation*...The 3 conformational states of the voltage dependent  $\text{Na}^+$  channel




- 1. Closed state
  - Activation gate closed
  - Inactivation gate open
  - no ion movement
- 2. Activated state:
  - Activation gate open
  - inactivation gate open
  - ions pass through
- 3. Inactivated state:
  - activation gate open
  - inactivation gate closed
  - no ion movement

## The nerve action potential....summary



## The nerve action potential....

- Generated in a specialised region of high  $\text{Na}^+$  channel density called the axon hillock. This signal must then be communicated to other nerves.




- How does this signal travel...

[http://hplismagazine.com/wp-content/uploads/neuron\\_structure2.jpg](http://hplismagazine.com/wp-content/uploads/neuron_structure2.jpg)

## Propagation of Electrical Signals

- 1) Some ionic current enters the axon during the depolarization phase of the action potential. A single part of the axon (e.g., the axon hillock at the base of the axon) becomes depolarized while the axonal region ahead (or behind) is still hyperpolarized

Triggering event opens  $\text{Na}^+$  channels




© 2001 Brooks/Cole - Thomson Learning



## Propagation of Electrical Signals

- 2) The point depolarization initiates a spread of current along the length of the axon. This **passively** depolarizes adjacent sections of the axon. This passive spread of current along a fibre also called "**local current flows**"

Local current flow occurs between the active and adjacent inactive areas




Spread of depolarization  
(c)



## Propagation of Electrical Signals

- 3) Some of this local current leaks out across the cell membrane (e.g., via "leak" channels) and hence the depolarization gets smaller and smaller as one moves along the axon. If too much current leaks out, the action potential would fail to propagate



\* Numbers refer to the local potential in mV at various points along the membrane.

© 2001 Brooks/Cole - Thomson Learning




## Propagation

- 4) The adjacent region is depolarized by the passive spread of current. Voltage-gated channels in the adjacent region are activated if the membrane potential reaches the voltage threshold, and the action potential ( $\text{Na}^+$  influx etc) occurs once again. The action potential is regenerated along the entire length of the axon. This is **active propagation**.

- If the action potential is not regenerated the depolarization will passively decay and die out.

- The region behind the propagating action potential is refractory due to  $\text{Na}^+$  channel inactivation




(d) Propagation continues



## Propagation of Electrical Signals


Some central and peripheral axons are **myelinated**. The wrappings of insulation decrease current leak out of the axon and thereby enable the passive voltage to spread further. This means the action potential propagation increases markedly, as the potential jumps ahead to the next Node of Ranvier. The need for ionic fluxes along the entire axon is reduced, so also increases efficiency.



(a)

## Propagation of Electrical Signals

The jumping nature of the spread of depolarisation is called **saltatory conduction**. Voltage-dependent  $\text{Na}^+$  channels and transmembrane ionic influx occurs only at the Nodes of Ranvier, located between the myelin sheaths.



*The amount of myelin and hence the conduction velocity can be regulated and may be fine-tuned according to cell function.*

## Learning Aims – lecture 5

- To understand the physiological mechanisms of the Action Potential, including:
  - Being able to draw the nerve AP with correctly identified x and y axis, and with the major phases of the AP identified
  - To appreciate the voltage threshold concept and to describe briefly what happens here to elicit the action potential (ie voltage-dependent gating).
  - To describe the ionic currents that mediate the depolarisation and repolarisation phases of the AP
  - To be able to describe the ion channels that mediate the above ionic currents, and to link the different conformation states of these channels (i.e. their gates) to the different phases of the AP. To define what is meant by voltage-dependent gating, activation and inactivation (in terms of these ion channels)
- To describe what is meant by passive and active propagation and describe how these two processes contribute to AP propagation along an axon
- To appreciate what myelin is and how it affects AP propagation velocity


Supporting activities:

- Membrane Potentials Virtual Prac (via Moodle)
- Cell Physiology tutorial #2. Action Potentials
- Stanfield, Chapter 7.4

## Next Concept (final one)

### Communication of Signals from cell to cell

A nerve cell communicates with other nerve cells and with peripheral organs, and does this largely by converting the action potential into a chemical messengers at synapses



Copyright © 2005 Pearson Education, Inc., publishing as Benjamin Cummings.