

Physiology 1A

Haemodynamics

Tim Murphy
Wallace Wurth 316
tim.murphy@unsw.edu.au

1

Haemodynamics

Physical laws governing blood flow and blood pressure.

How is the circulatory system organised for effective delivery of blood flow to tissues and organs?

Inter-relationships between:

- velocity of blood flow
- blood pressure
- dimensions of the components (vessels) in the systemic circulation (resistance).

3

From this lecture you should:

Know Poiseuille's Law and what it describes.

Understand the relationship between blood flow velocity and cross-sectional area in different sections of the vascular system.

Understand the concept of resistance to blood flow and the structure and arrangement of blood vessels to counter it.

Appreciate the concepts of laminar and turbulent blood flow.

Reference: Stanfield 6th Ed. Chapter 14 pp. 394 - 404.



2

For blood to flow through the vessels, we need a driving force - pressure.

Flow is opposed by the resistance provided by the blood vessels.

$$\text{Flow} = \frac{\text{Pressure gradient}}{\text{Resistance}} \quad Q = \frac{\Delta P}{R} \text{ (cf. Ohm's Law)}$$


4




5

Flow is directly proportional to the pressure gradient (inflow pressure – outflow pressure)

$$Q \propto \Delta P (P_i - P_o)$$



Flow is inversely proportional to the length of the conducting vessel: $Q \propto 1 / l$



7

Pressure and Flow


Poiseuille's Law describes laminar flow of newtonian (homogenous) fluids through cylindrical tubes in terms of:

- Flow
- Pressure
- Tube dimensions
- Viscosity of fluid


6

Flow varies directly as the fourth power of the radius of the conducting vessel

$$Q \propto r^4$$



Flow is inversely proportional to the viscosity of the fluid: $Q \propto 1 / \eta$



8

Put together, Poiseuille's Law:

$$Q = \frac{\pi (\Delta P) r^4}{8\eta l} \quad \frac{\pi}{8} = \text{constant}$$

Radius of conducting vessel is most influential determinant.


9

Blood flows faster in narrower vessels.

True?

Total cross-sectional area (cm²)

Aorta	2.5
Arteries	20
Arterioles	40
Capillaries	2500
Venules	250
Veins	80
Venae cavae	8



Stanfield Fig. 14.15

11

Blood Flow Velocity


Velocity = distance/time (cm/s)

Flow = volume/time (cm³/s)

Volume depends upon cross-sectional area of blood vessel. Therefore:

Velocity = Flow / Area $v = Q / A$

Flow is constant in a 'rigid' system




10

Resistance to Flow

Resistance (R) to blood flow is the ratio of the pressure gradient to flow (c.f. Ohm's Law, $R = V / I$).

$$R = \frac{\Delta P}{Q} = \frac{8\eta l}{\pi r^4} \quad (\text{Stanfield; Poiseuille's Law})$$

Note effect of radius



Stanfield Fig. 14.9

12


Resistance to flow is greatest in arterioles:

- low cross-sectional area relative to radius
- constriction of arterioles

Resistance in series and parallel

Separate elements of circulation (artery, arteriole, capillary etc.) are arranged in series

Similar elements are arranged in parallel




Resistance in series is sum of individual resistances:

$$P_i \rightarrow R_1 \rightarrow R_2 \rightarrow R_3 \rightarrow P_o \quad R = \Delta P / Q$$

$$\Delta P_1 \quad \Delta P_2 \quad \Delta P_3 \quad R_t = R_1 + R_2 + R_3$$

13

Reciprocal of resistance in parallel is sum of reciprocal of individual resistances:



$$R = \Delta P / Q: \quad \frac{1}{R_t} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} \quad \frac{1}{R} \text{ is conductance}$$

If R_1 , R_2 and R_3 were equal then $1 / R_t = 3 / R_1$: $R_t = R_1 / 3$

For parallel arrangement, total resistance < individual resistance of any one segment:

$$R_t < R_1, R_2 \text{ or } R_3$$

15

14

Total Peripheral Resistance (TPR)

$$TPR = \frac{\text{arterio-venous pressure drop } (P_a - P_v)}{\text{flow}}$$

$$= \frac{\text{Blood Pressure}}{\text{Cardiac Output}} \quad (BP = CO \times TPR)$$

$$= \frac{100 \text{ mmHg} - 0 \text{ mmHg}}{5000 \text{ ml/min}}$$

$$= 0.02 \text{ mmHg/ml/min} = 0.02 \text{ PRU}$$

Individual tissues/organs may differ (e.g. kidney)

16


Effects of pressure and vascular resistance on tissue blood flow

Blood vessels are not rigid tubes

Aorta, large arteries:
elastic, fibrous conductance, compliance

Compliance is a measure of vessel 'elasticity'.

$$\text{Compliance} = \frac{\Delta V}{\Delta P}$$




17


Venules:

Thin wall, little muscle. Low resistance
capacitance (reservoir)

Veins: Thin wall, v. low resistance
capacitance, conductance




Distribution of blood by volume



19

Small arteries,
arterioles:
Muscular, resistance,
alter diameter, control
flow




Capillaries: thin
endothelial cells -
exchange

Copyright © 2009 Pearson Education, Inc., publishing as Benjamin Cummings

18

Laminar and Turbulent Flow

Due to friction with the vessel wall, flow of blood
(and other fluid) at a constant velocity in a
cylinder is laminar (series of layers).




Increased velocity enhances the effect



Increased velocity increases shear stress or viscous drag on walls

20

Laminar flow



Hodder Arnold / An Introduction to Cardiovascular Physiology © 2010 J. Rodney LeVick

21

Turbulent flow is more likely if:

- vessel is large diameter
- flow is faster
- blood viscosity is low (anemia)

Turbulent flow can be predicted by Reynold's number:


$$N_R = \frac{\text{density} \times \text{diameter} \times \text{velocity}}{\text{viscosity}} = \frac{\rho D v}{\eta} \quad \begin{array}{l} <2000, \text{ laminar} \\ >3000, \text{ turbulent} \end{array}$$

Turbulence may cause noise:

- e.g. anemia (low viscosity, high velocity)
- heart murmurs

23


Irregular fluid motions within vessel causes turbulent flow



Greater pressure is required to force turbulent flow through a tube (Poiseuille)

$$\begin{array}{l} \text{Laminar } Q \propto \Delta P \\ \text{Turbulent } Q \propto \sqrt{\Delta P} \end{array}$$

22



24