

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

Logic Summer School
Australian National University
4–16 December, 2023
Canberra, ACT, Australia



THE UNIVERSITY of EDINBURGH
informatics

lfcs

Laboratory for Foundations
of Computer Science



supported by:



THE ROYAL
SOCIETY

The
Alan Turing
Institute

Facebook Research NCSC

Plan:

- 1) Type-driven probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Wed)
- 3) Quasi Borel spaces (Wed) Simple type structure (Thu)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

please ask questions!

snibble



Course
web
page

Discrete model

$\text{type} : \text{Obs} \quad W := [0, \infty] \quad Bx := (\text{Thur})$

$DX := (\text{Fri})$

$Px := \{ \mu \in DX \mid C_{\mu}[X] = 1 \} \quad (\text{Thu})$

$C_{\mu}[E] := (\text{Fri}) \quad \delta_n := (\text{Fri})$

$\phi \mu_k := (\text{Fri})$

Def: Quasi-Borel space $X = (\mathcal{L}X, \mathcal{R}_X)$


$\mathcal{R}_X \subseteq \mathcal{L}X^{\mathbb{R}}$ Closed under:


Set
"carrier"

Set of
functions $\alpha: \mathbb{R} \rightarrow \mathcal{L}X$
"random elements"

- Constant S : 

- recombination



- precomposition: 

Examples

recombination of constants

$$- \mathbb{R} = (\mathbb{R}_J, \text{Meas}(\mathbb{R}, \mathbb{R}))$$

qbs underlying \mathbb{R}

$$- X \in \text{Set}, \quad \ulcorner X \urcorner^{\text{qbs}} := (X, \sigma\text{-simple}(\mathbb{R}, X))$$

$\lambda r. \left\{ \begin{array}{l} \vdots \\ r \in A_n: x_n \\ \vdots \end{array} \right.$

discrete qbs on X

$$- \quad \ulcorner X \urcorner_{\text{qbs}} := (X, X^{\mathbb{R}_J})$$

all functions

Indiscrete qbs on X

Validate gibbs axioms for: $\mathcal{W} := ([0, \infty], \text{Meas}(\mathbb{R}, \mathcal{W}))$

• Constants:

$E : \mathcal{B}_{\mathcal{W}}, x : \mathcal{W} \vdash$

$$(\lambda r : \mathbb{R}. x)^{-1}[E] = \begin{cases} x \in E : & \mathbb{R} \\ x \notin E : & \emptyset \end{cases} \in \mathcal{B}_{\mathbb{R}} \quad \checkmark$$

Validate gbs axioms for: $\mathcal{W} := ([0, \infty], \text{Meas}(\mathbb{R}, \mathcal{W}))$

- Precomposition:

$$\alpha: \text{Meas}(\mathbb{R}, \mathcal{W}), \quad \varphi: \text{Meas}(\mathbb{R}, \mathbb{R}) \vdash$$

$$\mathbb{R} \xrightarrow{\varphi} \mathbb{R} \xrightarrow{\alpha} \mathcal{W} \in \text{Meas}(\mathbb{R}, \mathcal{W})$$

\uparrow
 Meas is a cat.

Explicitly:

$$(\alpha \circ \varphi)^{-1}[E] \in \mathcal{B}\mathbb{R} \xleftarrow{\varphi^{-1}} \alpha^{-1}[E] \in \mathcal{B}\mathbb{R} \xleftarrow{\alpha^{-1}} E \in \mathcal{B}\mathcal{W} \quad \checkmark$$

Validate gbs axioms for: $W := ([0, \infty], \text{Meas}(\mathbb{R}, W))$

• RL Combination

$$I \text{ ctbl}, \alpha_-: \text{Meas}(\mathbb{R}, W)^I, E_-: B_{\mathbb{R}}^I, R = \bigcup_{i \in I} E_i, F: B_W^+$$

$$\left(\lambda r. \left\{ \begin{array}{c} \vdots \\ r \in E_i : \alpha_i r \\ \vdots \end{array} \right\} \right)^{-1}[F]$$

$$\beta := \bigcup_{i \in I} \alpha_i^{-1}[F] \cap E_i \in B_{\mathbb{R}}$$

In fact:


$$r \in \text{LHS} \Leftrightarrow \beta r \in F \Leftrightarrow \exists i \in I. r \in E_i \wedge \alpha_i r \in F \Leftrightarrow r \in \text{RHS}$$

✓

σ -simple function

$\alpha: \mathbb{R} \rightarrow X$ s.t. $\text{Im } \alpha := \alpha[\mathbb{R}]$ is ctbl 1

$\forall x \in \text{Im } \alpha. \alpha^{-1}[x] \in \mathcal{B}_{\mathbb{R}}$



Validate gbs axioms for: $\Gamma_X^{\text{Obs}} := (X, \sigma\text{-simple}(X))$

- Constants

$$\text{Im}(\lambda r. x) = \{x\} \text{ ctbl } \checkmark$$

NB: f σ -simple:
 $\text{Im } f \text{ ctbl } \wedge$
 $f^{-1}[x] \in \mathcal{B}_{\mathbb{R}}$

$$g: X \vdash (\lambda r. x)^{-1}[y] = \begin{cases} x=y: \mathbb{R} \\ x \neq y: \emptyset \end{cases} \in \mathcal{B}_{\mathbb{R}} \checkmark$$

Validate gbs axioms for: $\Gamma^{Obs}_X := (X, \sigma\text{-simple}(X))$

• PreComposition:

$\alpha : \sigma\text{-simple}(X), \varphi : \text{Meas}(\mathbb{R}, \mathbb{R}) \vdash$

$\text{Im}(\alpha \circ \varphi) \subseteq \text{Im} \alpha \text{ ctbl} \quad \checkmark$

NB: $f \sigma\text{-simple} :$
 $\text{Im } f \text{ ctbl} \wedge$
 $f^{-1}[x] \in \mathcal{B}_{\mathbb{R}}$

$x : X \vdash$

$(\alpha \circ \varphi)^{-1}[x] = \varphi^{-1}[\alpha^{-1}(x)] \in \mathcal{B}_{\mathbb{R}} \quad \checkmark$

$\alpha^{-1}(x) \in \mathcal{B}_{\mathbb{R}}$

$\varphi : \mathbb{R} \rightarrow \mathbb{R} \text{ measurable}$

Validate gbs axioms for: $\Gamma_X^{Obs} := (X, \sigma\text{-simple}(X))$

• recombination:

$$\alpha_i : (\sigma\text{-simple}(X))^I, E_i : \mathcal{B}_R, R = \biguplus_{i \in I} E_i \vdash$$

NB: $f \sigma\text{-simple}$:
 $\text{Im } f \text{ ctbl } \wedge$
 $f^{-1}[x] \in \mathcal{B}_R$

$$\text{Im}[E_i \cdot \alpha_i]_{i \in I} \subseteq \bigcup_{i \in I} \text{Im } \alpha_i \quad \text{ctbl} \quad \checkmark$$

$x : X \vdash$

$$[E_i \cdot \alpha_i]_{i \in I}^{-1}(x) = \bigcup_{i \in I} \alpha_i^{-1}[x] \cap E_i \in \mathcal{B}_R \quad \checkmark$$

Prop: $X : \text{Set}, A : \text{Obs} \vdash$

$$\bullet \quad \forall f : X \rightarrow_{\perp} A. \quad f : \ulcorner X^{\text{Obs}} \urcorner \rightarrow A$$

$$\bullet \quad \forall f : \ulcorner A \urcorner \rightarrow X. \quad f : A \rightarrow_{\perp} \ulcorner X \urcorner_{\text{Obs}}$$

Prop: $X : \text{Set}, A : \text{Qbs} \vdash$

$$\bullet \forall f: X \rightarrow \perp A, \tilde{f}: \ulcorner X^{\text{Qbs}} \urcorner \rightarrow A$$

Pf: $\alpha: R_{r_X^{\text{Qbs}}} \vdash \alpha \text{ } \sigma\text{-simple} \Rightarrow$

$$\alpha = [\alpha^{-1}[x].\lambda r. x]_{x \in \text{Im } \alpha} \Rightarrow$$

$$(f \circ \alpha) = [\alpha^{-1}[x].\lambda r. f x]_{x \in \text{Im } \alpha}$$

Borel



constat $\in B_A$



recombination
 $\in \mathcal{R}_A$




Prop: $X : \text{Set}, A : \text{Qbs} \vdash$

- $\forall f : X \rightarrow_L A, \tilde{f} : \overset{\text{Qbs}}{X} \rightarrow A$
- $\forall f : \overset{\text{Qbs}}{A} \rightarrow X, \tilde{f} : A \rightarrow \overset{X}{\underset{\text{Qbs}}{L}}$

Prf: $\alpha : R_A \vdash (f \circ \alpha : R \rightarrow X) \in R_{\overset{X}{\underset{\text{Qbs}}{L}}} \text{ always. } \checkmark$



Useful adjunctions:



$$\mathcal{L}_{\text{Qbs}}^V := (\mathcal{L}V, \text{Meas}(R, V))$$

$$(V \in \text{Meas})$$

$$\Gamma_X^{\text{Meas}} := \left\{ A \subseteq \mathcal{L}X \mid \forall \alpha \in R_X. \alpha^{-1}[A] \in \mathcal{B}_R \right\}$$

- limits (products, subspaces)
and colimits (coproducts, quotients)
as in Set

- Slogan: every measurable space is carried by a qbs

Example

Product $(X \times Y, \pi_1, \pi_2)$:

- $L_{X \times Y} = L_{X \times Y}$ *necessarity!*

- $R_{X \times Y} = \{ \lambda r. (\alpha r, \beta r) \mid \alpha \in R_X, \beta \in R_Y \}$

*correlated
random
elements*

rest of structure as in Set.

Function Spaces

Straightforward!

$$- \text{ } \llbracket Y^X \rrbracket := \text{Obs}(X, Y)$$

$$- R_{Y^X} := \text{uncurry}[\text{Obs}(\mathbb{R} \times X, Y)]$$


$$= \{ \alpha: \mathbb{R} \rightarrow \llbracket Y^X \rrbracket \mid \lambda(r, x). \alpha r x: \mathbb{R} \times X \rightarrow Y \}$$

$$- \text{eval}: Y^X \times X \rightarrow Y$$
$$\text{eval}(f, x) := fx$$

Meas vs Obs

By generalities:

σ -algebra
on $\text{Meas}(R, R)$



$\vdash^{\text{Meas}}_{\mathbb{R}} \neg$

$\mathbb{R} \times \mathbb{R}$


\longrightarrow

$\vdash^{\text{Meas}}_{\mathbb{R}} \neg$

$\mathbb{R} \times \mathbb{R}$

~~\dashv~~

$\vdash \mathbb{R} = \mathbb{R}$



$\vdash^{\text{Meas}}_{\text{Eval}}$

$\left(\vdash^{\text{So}}_{\mathbb{R}} \mathbb{R} \times \mathbb{R} \neq \vdash \mathbb{R} \times \mathbb{R} \right)$

No factorisation
by
Aumann's
Theorem.

Simple Type Structure

"Simple" because:

- Simply-typed λ -calculus
- types are simple: $A, B : \text{Type} \vdash B^A : \text{Type}$
 - no polymorphism
 - no term dependency
- Contexts for terms: $\Gamma \vdash t : A$
are simple: $\Gamma = x_1 : A_1, \dots, x_n : A_n$
i.e. $\text{List}(\text{Type})$

Simple Type Structure

"Simple" because:

- interpretation is simple:

$$\llbracket x_1:A_1, \dots, x_n:A_n \rrbracket := \prod_{i=1}^n A_i$$

$$\llbracket \Gamma \vdash t:A \rrbracket : \llbracket \Gamma \rrbracket \longrightarrow A$$

in Qbs

Simple Type Structure Curry-Howard-Lambek

$$\frac{\Gamma \vdash t : A \quad \Gamma \vdash s : B}{\Gamma \vdash \langle t, s \rangle : A \times B} \rightsquigarrow \llbracket \Gamma \rrbracket \xrightarrow{\lambda r. \langle tr, sr \rangle} A \times B$$

is measurable

$$\frac{\Gamma \vdash t : A \times B \quad \Gamma, x:A, y:B \vdash s : C}{\Gamma \vdash \text{let } (x, y) = t \text{ in } s : C}$$

$$\Gamma \vdash \text{let } (x, y) = t \text{ in } s : C \rightsquigarrow$$

measurability
by
type!

$$\lambda r. \text{let } (a, b) = tr \text{ in } s \vdash [x \mapsto a, y \mapsto b]$$

$$\llbracket \Gamma \rrbracket \xrightarrow{\quad} C$$

is measurable. etc.

Random element Space

$$R_X := X^{\mathbb{R}} \quad \text{since} \quad \llbracket X^{\mathbb{R}} \rrbracket = R_X \text{ as sets.}$$

Why?

$$(\subseteq) \quad \alpha \in \llbracket X \rrbracket^{\mathbb{R}} \Rightarrow \alpha: \mathbb{R} \rightarrow X \text{ in Obs.}$$

$$\text{id}_{\mathbb{R}}: \mathbb{R} \rightarrow \mathbb{R} \text{ measurable} \Rightarrow \text{id} \in R_{\mathbb{R}}$$

$$\Rightarrow \alpha = \alpha \circ \text{id} \in R_X$$

$$(\supseteq) \quad \alpha \in R_X \Rightarrow \forall \psi \in R_{\mathbb{R}} = \text{Meas}(\mathbb{R}, \mathbb{R}). \quad \alpha \circ \psi \in R_X \Rightarrow \alpha: \mathbb{R} \rightarrow X \Rightarrow \alpha \in \llbracket X \rrbracket^{\mathbb{R}}$$

Pre composition
↙

Subspaces

For $X \in \text{Obs}$, $A \subseteq X$ Set:

$$R_A := \{ \alpha: \mathbb{R} \rightarrow A \mid \alpha \in R_X \}$$

Then $A = (A, R_A)$ is the **subspace** qbs

We write $A \hookrightarrow X$

Borel subspaces ensemble

The σ -algebra $B_X := \{ A \subseteq X \mid \forall \alpha \in R_X. \alpha^{-1}[A] \in B_R \}$


internalises as $B_X = 2^X$, the qbs of Borel subsets.

$\left(\begin{array}{l} B \\ \downarrow (B_R) \end{array} \right)$ are the Borel-on-Borel sets from descriptive set theory.
cf. [Sabou et al.'21]

Standard Borel Spaces

Def: A qbs S is **standard Borel** when

$$S \cong A \text{ for some } A \in \mathcal{B}_{\mathbb{R}}$$



Slogan: Qbs **conservative extension** of Sbs

Example $C_0 := \{f: \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ continuous}\} \hookrightarrow \mathbb{R}^{\mathbb{R}}$

C_0 is sbs. (Well-known!)

Proof:

$C_0' \in B_{\mathbb{R}^{\mathbb{Q}}}$ ^{sbs!}

$C_0' := \left\{ g \in \mathbb{R}^{\mathbb{Q}} \mid \begin{array}{l} \forall a, b \in \mathbb{Q}, \varepsilon \in \mathbb{Q}^+ \\ \exists \delta \in \mathbb{Q}^+ \forall p, q \in \mathbb{Q}^+ \cap [a, b] \\ |p - q| < \delta \Rightarrow |g(p) - g(q)| < \varepsilon \end{array} \right\}$

Then $C_0 \cong C_0' \in B_{\mathbb{R}^{\mathbb{Q}}}$:

$C_0 \rightarrow C_0'$

$\varphi \mapsto \varphi|_{\mathbb{Q}}$

$C_0' \rightarrow C_0$

$\psi \mapsto \lambda r. \lim_{n \rightarrow \infty} g(\text{approx } r \text{ by } (\frac{1}{m})_{m \in \mathbb{N}})_n$

on closed intervals
(= compact intervals)
Continuity \Uparrow uniform continuity
Borel measurable by type checks

Example (ctd)

C_0 is sbs, and $\text{eval}: C_0 \times \mathbb{R} \rightarrow \mathbb{R}$

is measurable.

Avoids:

- constructing complete separable metrics
- proving that evaluation is measurable
w.r.t. metric σ -algebra.

Non-examples ~ [Sabok et al. '21]

$$- \{ A \in \mathcal{B}_{\mathbb{R}} \mid A \neq \emptyset \} \hookrightarrow \mathcal{B}_{\mathbb{R}}$$

$$- \{ (A_1, A_2) \in \mathcal{B}_{\mathbb{R}}^2 \mid A_1 \subseteq A_2 \} \hookrightarrow \mathcal{B}_{\mathbb{R}}^2$$

$$- \{ A \in \mathcal{B}_{\mathbb{R}} \mid A \text{ open} \} \hookrightarrow \mathcal{B}_{\mathbb{R}}$$

Partiality cf. [Vàkàr et al. '19]

A Borel embedding $e: X \hookrightarrow Y$

- injective function $e: [X] \rightarrow [Y]$
- its image is Borel: $e[X] \in \mathcal{B}_Y$
- e is **Strong**: $\alpha \in R_X \iff e \circ \alpha \in R_Y$

Examples

- $\mathbb{1} \hookrightarrow \mathbb{2}$
- S is sbs $\iff \exists S \hookrightarrow \mathbb{R}$

Def: A Partial map $f: X \rightarrow Y$ is a morphism

$$f: X \rightarrow Y \sqcup \{\perp\}$$

Its domain of definition $\text{Dom } f := \{x \mid fx \neq \perp\}$

Partial hom-sets are ordered:



for $f, g: X \rightarrow Y$ $f \leq g$ when $\forall x. fx \neq \perp \Rightarrow gx = fx.$

[Cockett-Lack'06]

A model of restriction categories / axiomatic domain theory

[Fiorè-Plotkin'94]

Borel embeddings
are the admissible monos

Plan:

- 1) Type-driven probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Wed)
- 3) Quasi Borel spaces (Wed) simple type structure (Thu) ✓
- 4) Dependent type structure & standard Borel spaces (Thu) ✓
- 5) Integration & random variables (Fri)

please ask questions!

snibble



Course
web
page

Dependent Type Structure

Types can contain terms : a type referring to a term

$$X : \text{Type}, E : B_X \vdash \{x \in X \mid x \in E\} : \text{Type}$$

a type, just like
STLC

a term!

Dependent Type Structure

Types can contain terms :

$$X : \text{Type}, E : B_X \vdash \{x \in X \mid x \in E\} : \text{Type}$$

a type referring
to a term

a type, just like
STLC

a term!

Content formation:

$$\frac{\Gamma \vdash A : \text{Type}}{\Gamma, x : A \vdash}$$

Dependent Type Structure

types denote spaces-in-Content

$$\llbracket \Gamma \vdash A \rrbracket$$

$\downarrow \text{dep}$

$$\llbracket \Gamma \rrbracket$$

Dependent types denote spaces-in-Content

$\Gamma \vdash$ ← Context

$\llbracket \Gamma \vdash A \rrbracket$ ← Space in Context

$\Gamma \vdash A$
↑
Type in Context

↓ dep

$\llbracket \Gamma \rrbracket$

← Context Space

assigns
environment

E.g.:

A

↓

$\mathbb{1}$

Simple types

$\llbracket E : B_A \vdash \{x \in A \mid x \in E\} \rrbracket$

$\{ (E, a) \in B_A^{x_A} \mid a \in E \}$

↓ π_1

B_A

decoder

Content extension

$$\frac{\Gamma \vdash A}{\Gamma, a:A \vdash}$$

$$\llbracket \Gamma \vdash A \rrbracket$$

dep ↓

$$\llbracket \Gamma \rrbracket \quad \llbracket \Gamma, a:A \rrbracket := \llbracket \Gamma \vdash A \rrbracket$$

Substitution

$$\Gamma \vdash \sigma : \Delta$$

$$\llbracket \sigma \rrbracket : \llbracket \Gamma \rrbracket \longrightarrow \llbracket \Delta \rrbracket$$

E.g. *Weakening*

$$\Gamma, a:A \vdash \text{wkn} : \Gamma$$

$$\llbracket \Gamma, a:A \rrbracket := \llbracket \Gamma \vdash A \rrbracket \xrightarrow[\text{dep}]{\text{wkn}} \llbracket \Gamma \rrbracket$$

Action of Substitution on types

$$\begin{array}{ccc}
 \llbracket \Gamma \vdash A[\sigma] \rrbracket & & \\
 \downarrow \text{dep} := \pi_1 & \xrightarrow{\pi_2} & \llbracket \Delta \vdash A \rrbracket \\
 \{ (r, a) \in \llbracket \Gamma \rrbracket \times \llbracket A \rrbracket \mid \text{dep } a = r \} & & \Delta \\
 \downarrow & \xrightarrow{\sigma} & \downarrow \text{dep} \\
 \llbracket \Gamma \rrbracket & & \Delta
 \end{array}$$

E.g.

$$\begin{array}{ccc}
 \llbracket x:A \vdash B[x] \rrbracket := A \times B & \xrightarrow{\pi_2} & B \\
 \downarrow \text{dep} := \pi_1 & & \downarrow \\
 x:A & \xrightarrow{\langle \rangle} & \mathbb{1}
 \end{array}$$

simple type

Terms : sections

$$\llbracket \Gamma \rrbracket \xrightarrow{\llbracket \Gamma \vdash M : A \rrbracket} \llbracket \Gamma \vdash A \rrbracket$$

$$=$$

$$\llbracket \Gamma \rrbracket \xrightarrow{\text{dep}} \llbracket \Gamma \vdash A \rrbracket$$

e.g.

$$\mathbb{R} \xrightarrow{\llbracket x : \mathbb{R} \vdash [x, \infty) : \mathcal{B}_{\mathbb{R}}[wkn] \rrbracket} \mathbb{R} \times \mathcal{B}_{\mathbb{R}}$$

$$=$$

$$\mathbb{R} \xrightarrow{\pi_1} \mathbb{R}$$

E.g. Variables:

$$\llbracket \Gamma, a : A \vdash a : A \rrbracket$$

$$\llbracket \Gamma, a : A \rrbracket \xrightarrow{\langle \text{id}, \text{dep}_{\Gamma \vdash A} \rangle} \llbracket \Gamma, a : A \vdash A[wkn] \rrbracket$$

$$=$$


$$\llbracket \Gamma, a : A \rrbracket \xrightarrow{\text{dep}} \llbracket \Gamma, a : A \rrbracket$$

Exercise:
action of substitution
 $M[\sigma]$

Dependent Pairs

$$\frac{\Gamma, a:A \vdash B}{\Gamma \vdash \frac{\llcorner \llcorner}{a \in A} B}$$

$$\llbracket \frac{\llcorner \llcorner}{a \in A} A \rrbracket := \llbracket \Gamma, a:A \vdash B \rrbracket$$



Dependent products

$$\frac{\Gamma, a:A \vdash B}{\Gamma \vdash \prod_{a \in A} B}$$

$$\Gamma \vdash \prod_{a \in A} B$$

$$\llbracket \prod_{a \in A} B \rrbracket :=$$

$$\left\{ (r_0, f : \{ a \in \llbracket A \rrbracket \mid \text{dep } a = r_0 \} \rightarrow \llbracket \prod_{a:A} B \rrbracket) \mid \forall a \in \llbracket \prod_{a:A} B \rrbracket. \text{dep } a = r_0 \Rightarrow \text{dep } (f a) = a \right\}$$

Exercise: Find the random elements.

aha: $(a:A) \rightarrow B$

Discrete model

type : Obs $w := [0, \infty]$ $\mathcal{B}X \cong \mathcal{B}^X$

$DX := (\text{Fri})$

$PX := \{ \mu \in DX \mid C_{\mu}[X] = 1 \}$

$C_{\mu}[E] := (\text{Fri})$ $\delta_n := (\text{Fri})$

$\phi \mu k := (\text{Fri})$

Plan:

- 1) Type-driven probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Wed)
- 3) Quasi Borel spaces (Wed) Simple type structure (Thu) ✓
- 4) Dependent type structure & standard Borel spaces (Thu) ✓
- 5) Integration & random variables (Fri) ✓

please ask questions!

snibble



Course
web
page