

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

Logic Summer School
Australian National University
4–16 December, 2023
Canberra, ACT, Australia



THE UNIVERSITY of EDINBURGH
informatics

lfcs

Laboratory for Foundations
of Computer Science



supported by:



THE ROYAL
SOCIETY

The
Alan Turing
Institute

Facebook Research NCSC

Plan:

- 1) Type-driven probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Wed)
- 3) Quasi Borel spaces (Web) simple type structure (Thu)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)


please ask questions!

smile



Course
web
page

discrete model measure only histograms:



Want:

- lengths
- areas
- volumes.

Continuous *Caveat:*

Thm: No $\lambda: \mathcal{P}R \rightarrow [0, \infty]$:

$$\lambda(a, b) = b - a$$

(generalises length)

$$\lambda(r + A) = \lambda A$$

(translation invariant)

$$\lambda\left(\bigcup_{n=0}^{\infty} A_n\right) = \sum_{n=0}^{\infty} \lambda A_n$$

σ -additive

Thm: no $\lambda: \mathcal{P}\mathbb{R} \rightarrow [0, \infty]$:

$$\lambda(a, b) = b - a$$

(generalises length)

$$\lambda(r + A) = \lambda A$$


(translation invariant)

$$\lambda\left(\bigcup_{n=0}^{\infty} A_n\right) = \sum_{n=0}^{\infty} \lambda A_n$$

σ -additive

Direct proof in standard analysis courses. Idea behind typical proof is:

Thm: no $\lambda: \mathcal{P}S^1 \rightarrow [0, \infty]$
s.t.



$$r: \mathbb{R} \vdash \text{rotate}_r, [\theta] := [\theta + \tau r]$$

a) satisfy measure axioms for $BS' := \mathcal{P}S'$

b) invariant under rotations: $E: BS' \vdash$

$$\lambda \text{ rotate}[E] = \lambda E$$

c) $\lambda S' = \tau (= 2\pi)$

reduce $(S', \lambda^{S'})$ to (R, λ^R) via restriction & push forward

$$\lambda^R|_{\mathcal{P}[0,1]} := \lambda E \subseteq [0,1]. \lambda E : \mathcal{P}[0,1] \rightarrow W$$


$$\lambda^{S'} := \lambda E \subseteq S'. \lambda^R_{\mathcal{P}[0,1]}(\llbracket - \rrbracket^{-1}[E]) : \mathcal{P}S' \xrightarrow{\llbracket - \rrbracket^{-1}} \mathcal{P}[0,1] \xrightarrow{\lambda^R_{\mathcal{P}[0,1]}} W$$

noting

rotations in $S' \iff$ translations in R

Since $\nexists \lambda^{S'}$, we have $\nexists \lambda^R$ either.

Thm: no $\lambda: \mathcal{P}S' \rightarrow [0, \infty]$
s.t.



a) satisfy measure axioms for $BS' := \mathcal{P}S'$

b) invariant under rotations: $E: BS' \vdash$

$$\lambda \text{ rotate}_r[E] = \lambda E$$

c) $\lambda S' = \tau$ ($:= 2\pi$)

Proof: $a + b \Rightarrow \neg c$:

1) Using axiom of choice (AoC):

$$S' = \bigoplus_{i=0}^{\infty} E_i$$

$$E_i = \text{rotate}_{r_i}[E_0]$$

$$2) \lambda S' = \sum_{i=0}^{\infty} \lambda E_i = \sum_i \lambda \text{ rotate}_{r_i} 0 = \sum_{i=0}^{\infty} \lambda 0 = \begin{cases} \lambda 0 = 0 : 0 \\ \lambda 0 > 0 : \infty \end{cases} \neq \tau$$

Constructing E_i :

$$x, y: S' \vdash x \sim y := \exists q \in Q. \text{rotate}_q x = y \quad : \text{Prop}$$

$$\equiv \exists q \in [0, 1) \cap Q. \text{rotate}_q x = y$$

\sim -equivalence classes:

$$x: S' \vdash [x]_\sim := \{ y \in S' \mid x \sim y \} \quad : \mathcal{P}S'$$

$$C := \{ [x]_\sim \in \mathcal{P}S' \mid x \in S' \}$$

$$\forall e \in C, e \neq \emptyset, \text{ so by } AC: \exists \xi: C \longrightarrow S'. \xi_e \in e.$$

NB: ξ injective

Take $C_0 := \{z_e \in S' \mid e \in C\} \in \mathcal{P} S'$

Note: $x \sim y, x, y \in C_0 \vdash x = y$.

$q: \mathbb{Q} \vdash C_q := \underset{q}{\text{rotate}}[C_0] \in \mathcal{P} S'$

Let $(r_i)_{i=0}^{\infty}$ enumerate $\mathbb{Q} \cap [0, 1)$ s.t. $r_0 = 0$

Take $E_i := C_{r_i}$

By fiat: $E_i = C_{r_i} = \text{rotate}_{r_i}[C_0] = \text{rotate}_{r_i}[E_0]$

RTP: $S' = \bigoplus_{i=0}^{\infty} E_i$

NB: $x, y: S' \vdash$
 $x \sim y: \text{Prop}$
 $C = \sim\text{-equiv.}$
 $z: C \rightarrow S'$
 $e: C \vdash z_e \in e$

$$E_i \cap E_j = \emptyset, \quad i \neq j:$$

$$x \in E_1 \cap E_2 \Rightarrow \exists y_i \in \mathcal{C}. x = \text{rotate}_{r_i} y_i$$

$$\Rightarrow y_1 \sim x \sim y_2 \Rightarrow y_1 = y_2 =: y$$

$$\Rightarrow \text{rotate}_{r_2 - r_1} y = y, \quad |r_2 - r_1| < 1$$

$$\Rightarrow r_1 = r_2$$

$$\underline{S'} = \bigcup_{i=0}^{\infty} \underline{E_i} : \quad \text{letting } e := \xi_{[x]_n} : \rho S'$$

$$\xi_e, x \in e \Rightarrow \xi_e \sim x$$

$$\Rightarrow \exists q \in (\mathbb{Q} \cap [0, 1)) . \text{rotate}_q \xi_e = x .$$

$$\text{As } \xi_e \in C_0 : x \in C_q . \text{ Find } i \text{ s.t. } r_i = q$$

$$\text{and } x \in C_{r_i} = E_i .$$



Takeaway: Taking $B/R := \mathcal{D}R$

Excludes measures such as:

length, area, volume

Workaround: only measure well-behaved subsets

Df: The Borel subsets $\mathcal{B}_{\mathbb{R}} \subseteq \mathcal{P}\mathbb{R}$:

- open intervals $(a, b) \in \mathcal{B}_{\mathbb{R}}$

closure under σ -algebra operations:

$$\frac{}{\emptyset \in \mathcal{B}_{\mathbb{R}}}$$

↖
empty set

$$\frac{A \in \mathcal{B}_{\mathbb{R}}}{A^c := \mathbb{R} \setminus A \in \mathcal{B}}$$

↖
complements

$$\frac{\vec{A} \in \mathcal{B}_{\mathbb{R}}^{\mathbb{N}}}{\bigcup_{n=0}^{\infty} A_n \in \mathcal{B}_{\mathbb{R}}}$$

↖
countable unions

Examples

discrete Countable: $\{r\} = \bigcap_{\varepsilon \in \mathbb{Q}^+} (r - \varepsilon, r + \varepsilon) \in \mathcal{B}_{\mathbb{R}}$

I countable $\Rightarrow I = \bigcup_{r \in I} \{r\} \in \mathcal{B}_{\mathbb{R}}$

closed intervals: $[a, b] = (a, b) \cup \{a, b\}$

Non-examples?

More complicated: analytic, Lebesgue

Df: Measurable space $V = (V, B_V)$

Set
(carrier)

Family of
Subsets
 $B_V \subseteq P(V)$

closed under σ -algebra operations:

$$\frac{}{\emptyset \in B_V}$$

↖
empty set

$$\frac{A \in B_V}{A^c := V \setminus A \in B_V}$$

↖
complements

$$\frac{\vec{A} \in B_V^{\mathbb{N}}}{\bigcup_{n=0}^{\infty} A_n \in B_V}$$

↖
countable unions

Idea: Structure all spaces after the worst-case scenario

Examples

- Discrete spaces $\mathcal{X}^{\text{meas}} = (X, P_X)$
- Euclidean spaces \mathbb{R}^n — replace intervals with
boxes $\prod_{i=1}^n (a_i, b_i)$
 \mathbb{R}^N similarly
 $\{C \cap A \mid C \in \mathcal{B}_V\}$
/
- Sub spaces: $A \in \mathcal{P}_{\mathcal{V}_1}$ $A := (A, [B_V] \cap A)$
- Products: $A \times B := (\mathcal{L}A_1 \times \mathcal{L}B_1, \sigma([B_A] \times [B_B]))$

Def: Borel measurable functions $f: V_1 \rightarrow V_2$

- functions $f: V_1 \rightarrow V_2$
- inverse image preserves measurability:

$$f^{-1}[A] \in \mathcal{B}_{V_1} \iff A \in \mathcal{B}_{V_2}$$

Examples

- $(+), (\cdot) : \mathbb{R}^2 \rightarrow \mathbb{R}$
- $| \cdot |, \sin : \mathbb{R} \rightarrow \mathbb{R}$
- any continuous function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$
- any function $f: X \rightarrow V$

Category Meas

Objects : Measurable spaces

Morphisms : Measurable functions

Identities:

$$\text{id} : V \rightarrow V$$

Composition:

$$f : V_2 \rightarrow V_3 \quad g : V_1 \rightarrow V_2$$

$$f \circ g : V_1 \rightarrow V_3$$

Meas Category

Products, Coproducts / disjoint union, Subspaces
Categorical limits, colimits, but:

Thm [Aumann '61] No σ -algebras $B_{\mathbb{R}}, B_{\mathbb{R}^{\mathbb{R}}}$ for measurable


membership predicate $\leftarrow (\exists) : (B_{\mathbb{R}}, B_{B_{\mathbb{R}}}) \times \mathbb{R} \longrightarrow \text{Bool}$
 $(U, r) \longmapsto [r \in U]$

eval : $(\text{Meas}(\mathbb{R}, \mathbb{R}), B_{\mathbb{R}^{\mathbb{R}}}) \times \mathbb{R} \rightarrow \mathbb{R}$
 $(f, r) \mapsto f(r)$

Questions! skip proof?

Proof (Sketch):

Borel hierarchy:



Stabilises at $\Delta_{\omega_1}^0 = \mathcal{B}(\Sigma_0^0) = \Delta_{\omega_1+1}^0$

$$\text{rank } A := \min \{ \alpha < \omega_1 \mid A \in \Delta_\alpha^0 \}$$

then
for $B_{B_R} = P(B_R)$

$$(\exists) : (B_R, B_{B_R}) \times \mathbb{R} \rightarrow \mathbb{R}$$

$$(U, r) \mapsto [r \in U]$$

$$B_{V \times U} = B([B_V] \times [B_U])$$

If measurable:

$$\alpha := \text{rank}((\exists)^{-1}[\text{true}]) < \omega,$$

Take $A \in B_R$, $\text{rank } A > \alpha$

But:

$$\alpha < \text{rank } A = \text{rank}(A, \rightarrow)^{-1}[(\exists)^{-1}[\text{true}]] \leq \text{rank}((\exists)^{-1}[\text{true}]) \leq \alpha$$

~~##~~

More details in Ex. B

Sequential Higher-order structure:

I Countable : $V^I = \prod_{i \in I} V$

\Rightarrow Some higher-order structure in Meas:

Cauchy $\in B_{[-\infty, \infty]^N}$

$$\text{Cauchy} = \bigcap_{\epsilon \in \mathbb{Q}^+} \bigcup_{k \in \mathbb{N}} \bigcap_{\substack{m, n \in \mathbb{N} \\ m, n \geq k}} \{ \vec{y} \in [-\infty, \infty]^N \mid |y_m - y_n| < \epsilon \}$$

$\limsup : [-\infty, \infty]^N \rightarrow [-\infty, \infty]$ $\lim : \text{Cauchy} \rightarrow \mathbb{R}$

Compose higher-order building blocks:

lim is measurable!
↗

$$\text{VanishingSeq}(\mathbb{R}) := \left\{ \vec{r} \in \mathbb{R}^{\mathbb{N}} \mid \lim_{n \rightarrow \infty} r_n = 0 \right\} \in \mathcal{B}_{\mathbb{R}^{\mathbb{N}}}$$

$$\text{approx}_\Delta : \text{VanishingSeq}(\mathbb{R}^+) \times \mathbb{R} \longrightarrow \mathbb{Q}^{\mathbb{N}}$$

$$\text{s.t.}; \quad |(\text{approx}_{\Delta} \vec{r})_n - r| < \Delta_n$$

Slogan: Measurable by Type !

Not all operations of interest fit:

$$\limsup : ([-\infty, \infty]^{\mathbb{R}})^{\mathbb{N}} \longrightarrow [-\infty, \infty]^{\mathbb{R}}$$

$$\limsup := \lambda \vec{f}. \lambda x. \limsup_{n \rightarrow \infty} f_n x$$

Intrinsically higher-order !

Want

Slogan: measurability by type!

But

For higher-order building blocks

defer measurability proofs until

we resume 1st order fragment \Rightarrow non compositional

Plan:

- 1) Type-driven probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Wed)
- 3) Quasi Borel spaces (Web) simple type structure (Thu)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

please ask questions!

smile



Course
web
page

Plan

Def: $V \in \text{Meas}$ is **Standard Borel** when

$$V \cong A \quad \text{for some } A \in \mathcal{B}_{\mathbb{R}}$$


the "good part" of Meas — the subcategory

$$\text{Sbs} \hookrightarrow \text{Meas}$$

Conservative extensions:

Concrete spaces
we "observe"

Standard Borel spaces



Sbs includes

- Discrete \mathbb{I} , \mathbb{I} countable
- Countable products of Sbs:

$$\mathbb{R}^n, \mathbb{R}^{\mathbb{N}}, \mathbb{Z}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}$$

- ~ Borel subspaces of Sbs:

$$\mathbb{I} := [0, 1]$$

$$\mathbb{R}^+ := (0, \infty) \quad \mathbb{R}_{\geq 0} := [0, \infty]$$

- Countable coproducts of Sbs:


$$\mathbb{N} := [0, \infty]$$

$$\overline{\mathbb{R}} := [-\infty, \infty]$$

Cone idea


Measure Theory

sample space Ω Obs Theory




random element

$\downarrow \alpha$



Primitive notions:

measurable subset



Derived

notions:

random

elements

$\alpha: \Omega \rightarrow \text{Space}$

measure

measurable subsets

Def: Quasi-Borel space $X = (\mathcal{L}X, \mathcal{R}_X)$

$\mathcal{R}_X \subseteq \mathcal{L}X^{\mathbb{R}}$ Closed under:

Set
"carrier"


Set of
functions $\alpha: \mathbb{R} \rightarrow \mathcal{L}X$
"random elements"

- Constant S:

$$\frac{x \in \mathcal{L}X}{(\lambda r. x) \in \mathcal{R}_X}$$

- Precomposition:

- recombination



Def: Quasi-Borel space

$$X = (\mathcal{X}, R_X)$$

Set
"carrier"


Set of
functions $\alpha: \mathbb{R} \rightarrow \mathcal{X}$
"random elements"

$R_X \subseteq \mathcal{X}^{\mathbb{R}}$ Closed under:

- precomposition:

$$\alpha \in R_X \quad \varphi: \mathbb{R} \rightarrow \mathbb{R} \text{ in Sbs}$$

$$\varphi \circ \alpha: \mathbb{R} \xrightarrow{\varphi} \mathbb{R} \xrightarrow{\alpha} \mathcal{X} \in R_X$$



Def: Quasi-Boel space

$$X = (\mathcal{X}, \mathcal{R}_X)$$

Set
"carrier"



Set of
functions $\alpha: \mathbb{R} \rightarrow \mathcal{X}$
"random elements"

$$\mathcal{R}_X \subseteq \mathcal{L}(\mathbb{R}, \mathcal{X}) \quad \text{Closed under:}$$

- re combination

$$\vec{\alpha} \in \mathcal{R}_X^{\mathbb{N}} \quad \mathbb{R} = \bigcup_{n=0}^{\infty} A_n \quad \text{EBR}$$

$$\lambda r. \left\{ \begin{array}{c} \vdots \\ r \in A_n: \alpha_n r \\ \vdots \end{array} \right.$$



Def: Quasi-Borel space $X = (\mathcal{L}X, \mathcal{R}_X)$


$\mathcal{R}_X \subseteq \mathcal{L}X^{\mathbb{R}}$ Closed under:


Set
"carrier"

Set of
functions $\alpha: \mathbb{R} \rightarrow \mathcal{L}X$
"random elements"

- Constant S: 

- recombination



- precomposition: 

Examples

recombination of constants

$$- \mathbb{R} = (\mathbb{R}_J, \text{Meas}(\mathbb{R}, \mathbb{R}))$$

qbs underlying \mathbb{R}

$$- X \in \text{Set}, \quad \ulcorner X \urcorner^{\text{qbs}} := (X, \sigma\text{-simple}(\mathbb{R}, X))$$

$\lambda r. \begin{cases} \vdots \\ r \in A_n: x_n \\ \vdots \end{cases}$

discrete qbs on X

$$- \quad \ulcorner X \urcorner_{\text{qbs}} := (X, X^{\mathbb{R}_J})$$

all functions

Indiscrete qbs on X

Qbs morphism $f: X \rightarrow Y$

- function $f: X \rightarrow Y$

$$\frac{\alpha \downarrow_{X_1} \in R_X}{\text{---}}$$

$$\begin{array}{c} R \\ \alpha \downarrow \\ X_1 \\ f \downarrow \\ Y_1 \end{array} \in R_Y$$

Example

- Constant functions

are qbs
morphisms

- σ -simple functions
are qbs morphisms

Category Qbs

\Leftarrow

- identity, composition

Discrete model

$\text{type} : \text{Obs} \quad W := [0, \infty] \quad B_X := (\text{Thur})$

$D_X := (\text{Fri})$

$P_X := \{ \mu \in D_X \mid C_{\mu}[X] = 1 \} \quad (\text{Thu})$

$C_{\mu}[E] := (\text{Fri}) \quad \delta_n := (\text{Fri})$

$\phi \mu_k := (\text{Fri})$

Plan:

- 1) Type-driven probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Wed)
- 3) Quasi Borel spaces (Web) simple type structure (Thu)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

please ask questions!

snibble



Course
web
page