

A domain theory for statistical probabilistic programming

Matthijs Vákár, Ohad Kammar, and Sam Staton



Logic and Semantics Seminar
Department of Computer Science and Technology
The Computer Laboratory
22 February 2019



Statistical probabilistic programming


$\llbracket - \rrbracket$: programs \rightarrow unnormalised distributions

- ▶ Bayesian inference: compiler computes normalisation
- ▶ Continuous types: $\mathbb{R}, [0, \infty]$
- ▶ Probabilistic effects:

normally
distributed
sample

sample(μ, σ) : \mathbb{R}

$\llbracket \text{sample}(0, 2) \rrbracket$



scale
distribution
by r

$r : [0, \infty]$
score(r) : 1

conditioning/fitting
to observed data
with likelihood

prior

```
let  $x = \text{sample}(0, 2)$   
in score( $\text{normalPdf}(1.1 \mid x, \frac{1}{4})$ );  
    score( $\text{normalPdf}(1.9 \mid 2x, \frac{1}{4})$ );  
    score( $\text{normalPdf}(2.7 \mid 3x, \frac{1}{4})$ );  
 $x$ 
```

posterior

Statistical probabilistic programming

► Commutativity/exchangability/Fubini

Exact Bayesian inference
using disintegration
[Shan-Ramsey'17]

$$\left[\begin{array}{l} \text{let } x = K \text{ in} \\ \text{let } y = L \text{ in} \\ f(x, y) \end{array} \right] = \left[\begin{array}{l} \text{let } y = L \text{ in} \\ \text{let } x = K \text{ in} \\ f(x, y) \end{array} \right] \quad \int \llbracket K \rrbracket (dx) \int \llbracket L \rrbracket (dy) f(x, y) = \int \llbracket L \rrbracket (dy) \int \llbracket K \rrbracket (dx) f(x, y)$$

probability
distributions



σ -finite
distributions



arbitrary
distributions



s-finite
distributions



not closed under
push-forward


full definability
[Staton'17]

Statistical probabilistic programming


Express continuous distributions using:

- Higher-order functions:

(e.g. generative random function models)



piecewise(random-constant)




piecewise(random-linear)

- Term recursion:

```
rw(x, σ) = λ().    // thunk  
  let y = sample(x, σ)  
  in (x, rw(y, σ))
```

(e.g. Gaussian random walk)



- Type recursion (à la FPC)

(e.g. dynamic types, IRs)

$$Dynamic = \mu\alpha. \{ \text{Val}(\mathbb{R}) \mid \text{Fun}(\alpha \rightarrow \alpha) \}$$

Application: modular Bayesian inference

Resample-Move Sequential Monte Carlo

[Ścibior et al.'18a+b]


resamples k particles n moves t recursion

```
rmsmc k n t =  
  marginal . finish . compose k (  
    advance . hoistS (  
      compose t mhStep . hoistT resample  
    )  
  ) . hoistST (spawn n >>)
```

inference representation

inference transformer


higher order



inference transformation (invariant preserving)

recursive types

ProbProg: Important Language Features

Church  WebPPL Venture	sample	\mathbb{R}	score	higher order	term rec	type rec	Fubini (commute)
sets + probability	✓	✗	✗	✓	✗	✗	✓
meas space + subprobability	✓	✓	✗	✗	1 st	✗	✓
CPO + subprobability	✓	✓	✗	✓	✓	✓	?
cont domain + subprobability [Jones-Plotkin'89]	✓	✓	✗	✗	1 st	✗	✓
⋮ [Jung-Tix'98]	⋮	⋮	⋮	⋮	⋮	⋮	⋮
meas + s-finite distributions [Staton'17]	✓	✓	✓	✗	1 st	✗	✓
qbs + s-finite distributions [Heunen et al'17, Ścibior et al'18]	✓	✓	✓	✓	1 st	✗	✓
coh/meas cone + probability [Ehrhard-Pagani-Tasson'18, Ehrhard-Tasson'15-'19]	✓	✓ ✗	✗	✓	✓	? ✓	? ✓
ω qbs + s-finite distributions [This work]	✓	✓	✓	✓	✓	✓	✓



Summary

Contribution

- ▶ $\omega\mathbf{Qbs}$: a category of pre-domain quasi-Borel spaces
- ▶ M : commutative probabilistic powerdomain over $\omega\mathbf{Qbs}$
- ▶ Axiomatic treatment of measure and domain theory in $\omega\mathbf{Qbs}$
- ▶ Adequacy: $(\omega\mathbf{Qbs}, M)$ adequately interprets:
 - ▶ Statistical FPC
 - ▶ Untyped Statistical λ -calculus

This talk

- ▶ $\omega\mathbf{Qbs}$
- ▶ A probabilistic powerdomain
- ▶ Axiomatic treatment
- ▶ Characterising $\omega\mathbf{Qbs}$

Iso-recursive types: FPC

type variable contexts

$$\Delta = \{\alpha_1, \dots, \alpha_n\}$$

[Fiore-Plotkin'94]

$$\frac{\Delta, \alpha \vdash_k \tau : \text{type}}{\Delta \vdash_k \mu\alpha.\tau : \text{type}}$$

$$\begin{aligned} \text{Lam} = \mu\alpha. \{ & \text{Bool}\{\text{True} \mid \text{False}\} \\ & \mid \text{App}(\alpha * \alpha) \\ & \mid \text{Abs}(\alpha \rightarrow \alpha) \} \end{aligned}$$

type recursion

$$\frac{\Gamma \vdash t : \sigma[\alpha \mapsto \tau]}{\Gamma \vdash \tau.\text{roll}(t) : \tau} \quad \frac{\Gamma \vdash t : \tau \quad \Gamma, x : \sigma[\alpha \mapsto \tau] \vdash s : \rho}{\Gamma \vdash \text{match } t \text{ with roll } x \Rightarrow s : \rho}$$

Iso-recursive types: FPC

[Fiore-Plotkin'94]

type variable contexts

$$\Delta = \{\alpha_1, \dots, \alpha_n\}$$

$$\frac{\Delta, \alpha \vdash_k \tau : \text{type}}{\Delta \vdash_k \mu\alpha.\tau : \text{type}}$$

ω Cpo-enriched
category of
domains


$$\llbracket \Delta \vdash_k \tau : \text{type} \rrbracket : (\mathcal{C}^{\text{op}})^n \times \mathcal{C}^n \rightarrow \mathcal{C}$$

$$\llbracket \Delta \vdash_k \mu\alpha.\tau : \text{type} \rrbracket = \text{minimal invariants}$$

[Freyd'91,92, Pitts'96]

locally continuous
functor

Challenge

- ▶ probabilistic powerdomain
 - ▶ commutativity/Fubini
 - ▶ domain theory
 - ▶ higher-order functions
- continuous domains [Jones-Plotkin'89]
- open problem [Jung-Tix'98]
- 

traditional approach:

domain \mapsto Scott-open sets \mapsto Borel sets \mapsto distributions/valuations

our approach: ^{as in} [Ehrhard-Pagani-Tasson'18]



(domain, quasi-Borel space) \mapsto distributions

separate
but compatible



Rudimentary measure theory

Borel sets

- ▶ $[a, b]$ Borel
- ▶ A Borel $\implies A^c$ Borel
- ▶ $(A_n)_{n \in \mathbb{N}}$ Borel $\implies \bigcup_{n \in \mathbb{N}} A_n$ Borel

Measurable functions $f: \mathbb{R} \rightarrow \mathbb{R}$

$$f^{-1}[A] \text{ Borel} \iff A \text{ Borel}$$

Measures $\mu: \text{Borel} \rightarrow [0, \infty]$

- ▶ monotone:
 $A \subseteq B \implies \mu(A) \leq \mu(B)$
- ▶ Scott-continuous:
 $A_0 \subseteq A_1 \subseteq \dots \implies \mu(\bigcup_n A_n) = \bigvee_n \mu(A_n)$
- ▶ strict ($\mu \emptyset = 0$) and additive ($\mu(A \uplus B) = \mu A + \mu B$)

Rudimentary measure theory

Borel sets

- ▶ $[a, b]$ Borel
- ▶ A Borel $\implies A^c$ Borel
- ▶ $(A_n)_{n \in \mathbb{N}}$ Borel \implies
 $\bigcup_{n \in \mathbb{N}} A_n$ Borel

Measurable functions $f: \mathbb{R} \rightarrow \mathbb{R}$

$$f^{-1}[A] \text{ Borel} \iff A \text{ Borel}$$

Measures $\mu: \text{Borel} \rightarrow [0, \infty]$

- ▶ monotone:
 $A \subseteq B \implies \mu(A) \leq \mu(B)$
- ▶ Scott-continuous:
 $A_0 \subseteq A_1 \subseteq \dots \implies \mu(\bigcup_n A_n) = \bigvee_n \mu(A_n)$
- ▶ strict ($\mu \emptyset = 0$) and additive ($\mu(A \uplus B) = \mu A + \mu B$)

1 dimensional

Example (Lebesgue measures)

$$\begin{aligned} \lambda[a, b] &= b - a \text{ on } \mathbb{R} \\ (\lambda \otimes \lambda)([a, b] \times [c, d]) &= \\ &= (b - a)(d - c) \text{ on } \mathbb{R}^2 \end{aligned}$$

2 dimensional

Example (Push-forward measure)

$$f_*\mu(A) := \mu(f^{-1}[A])$$

Borel set

measure

$f: \mathbb{R} \rightarrow \mathbb{R}$

Quasi-Borel pre-domains


ω -qbs:

set partial order on \underline{X} subset of functions $\mathbb{R} \rightarrow \underline{X}$
"random elements"

$$X = (\underline{X}, \leq_X, M_X)$$

Quasi-Borel pre-domains


ω -qbs:




$$\blacksquare x_0 \leq x_1 \leq x_2 \leq \dots \quad \implies \quad \exists \bigvee_n x_n$$

Quasi-Borel pre-domains

ω -qbs:




- $\lambda_{_}.x \in M_X$



Quasi-Borel pre-domains

ω -qbs:



- $\lambda_.x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$

$\mathbb{R} \xrightarrow{\varphi}_{\text{Borel}} \mathbb{R}$



$\models \varphi$




$\models \alpha$



Quasi-Borel pre-domains

ω -qbs:




- $\lambda_.x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$
- $(\alpha_n \in M_X)_{n \in \mathbb{N}} \implies [r \in S_n. \alpha(r)] \in M_X$

$$\mathbb{R} \xrightarrow{\varphi} \text{Borel } \mathbb{R}$$



$$[S_n. \alpha_n]$$




Borel measurable
countable partition

$$\mathbb{R} = \biguplus_{n \in \mathbb{N}} S_n$$


Quasi-Borel pre-domains

ω -qbs:




- $\lambda _ . x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$
- $(\alpha_n \in M_X)_{n \in \mathbb{N}} \implies [r \in S_n . \alpha(r)] \in M_X$

s.t.:




Quasi-Borel pre-domains

ω -qbs:



- $\lambda _ . x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$
- $(\alpha_n \in M_X)_{n \in \mathbb{N}} \implies [r \in S_n . \alpha(r)] \in M_X$

s.t.:



Morphisms $f : X \rightarrow Y$: Scott continuous qbs maps

monotone and
 $f \bigvee_n x_n = \bigvee_n f x_n$

$\forall \alpha \in M_X .$
 $f \circ \alpha \in M_Y$

Quasi-Borel pre-domains

Example

$S = (\underline{S}, \Sigma_S)$ measurable space

$$(\underline{S}, =, \{\alpha : \mathbb{R} \rightarrow \underline{S} \mid \alpha \text{ Borel measurable}\})$$

so $\mathbb{R} \in \omega\mathbf{Qbs}$



Reminder

$$\mathbf{wqbs}: X = (\underline{X}, \leq_X, M_X)$$

- $\lambda _ . x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$
- $(\alpha_n \in M_X)_{n \in \mathbb{N}} \implies [r \in S_n . \alpha(r)] \in M_X$

s.t.:

$$(\alpha_n) \in M_X^\omega \implies \bigvee_n \alpha_n \in M_X$$


Quasi-Borel pre-domains

Example

$P = (\underline{P}, \leq_P)$ ω -cpo

lubs of
step functions

$$\left(\underline{P}, \leq_P, \left\{ \bigvee_k [_ \in S_n^k \cdot a_n^k] \mid \forall k. \mathbb{R} = \biguplus_n S_n^k \right\} \right)$$



so $\mathbb{L} = ([0, \infty], \leq, \{\alpha : \mathbb{R} \rightarrow [0, \infty] \mid \alpha \text{ Borel measurable}\}) \in \omega\mathbf{Qbs}$

Reminder

wqbs: $X = (\underline{X}, \leq_X, M_X)$


- $\lambda _ . x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$
- $(\alpha_n \in M_X)_{n \in \mathbb{N}} \implies [r \in S_n \cdot \alpha(r)] \in M_X$

s.t.:

$$(\alpha_n) \in M_X^\omega \implies \bigvee_n \alpha_n \in M_X$$

Example

X ω -qbs



$$X_{\perp} := \left(\{\perp\} + \underline{X}, \perp \leq \underline{X}, \left\{ [S.\perp, S^{\mathbb{C}}.\alpha] \mid \alpha \in M_X, S \text{ Borel} \right\} \right)$$

Reminder

wqbs: $X = (\underline{X}, \leq_X, M_X)$

- $\lambda_{\perp}.x \in M_X$
- $\alpha \in M_X \implies \alpha \circ \varphi \in M_X$
- $(\alpha_n \in M_X)_{n \in \mathbb{N}} \implies [r \in S_n.\alpha(r)] \in M_X$

s.t.:

$$(\alpha_n) \in M_X^{\omega} \implies \bigvee_n \alpha_n \in M_X$$

Quasi-Borel pre-domains

Products

$$\underline{X_1} \times \underline{X_2} = \underline{X_1} \times \underline{X_2} \qquad x \leq y \iff \forall i. x_i \leq y_i$$

$$M_{X_1 \times X_2} = \{(\alpha_1, \alpha_2) : \mathbb{R} \rightarrow \underline{X_1} \times \underline{X_2} \mid \forall i. \alpha_i \in M_{X_i}\}$$



correlated
random elements

Quasi-Borel pre-domains

Products

$$\underline{X_1} \times \underline{X_2} = \underline{X_1} \times \underline{X_2} \quad x \leq y \iff \forall i. x_i \leq y_i$$

$$M_{X_1 \times X_2} = \{(\alpha_1, \alpha_2) : \mathbb{R} \rightarrow \underline{X_1} \times \underline{X_2} \mid \forall i. \alpha_i \in M_{X_i}\}$$

Theorem

$\omega\mathbf{Qbs} \rightarrow \omega\mathbf{Cpo} \times \mathbf{Qbs}$ *creates limits*

correlated
random elements




Quasi-Borel pre-domains

Products

$$\underline{X_1} \times \underline{X_2} = \underline{X_1} \times \underline{X_2} \quad x \leq y \iff \forall i. x_i \leq y_i$$

$$M_{\underline{X_1} \times \underline{X_2}} = \{(\alpha_1, \alpha_2) : \mathbb{R} \rightarrow \underline{X_1} \times \underline{X_2} \mid \forall i. \alpha_i \in M_{X_i}\}$$



correlated
random elements

Exponentials

$$\begin{aligned} \underline{Y^X} &= \{f : \underline{X} \rightarrow \underline{Y} \mid f \text{ Scott continuous qbs morphism}\} \\ &= \mathbf{Qbs}(X, Y) \end{aligned}$$

$$\underline{f} \leq \underline{g} \iff \forall x \in \underline{X}. f(x) \leq g(x)$$

$$\underline{M_{Y^X}} = \left\{ \alpha : \mathbb{R} \rightarrow \underline{Y^X} \mid \begin{array}{l} \text{uncurry } \alpha : \mathbb{R} \times X \rightarrow Y \\ \text{Scott continuous qbs morphism} \end{array} \right\}$$

so $\underline{Y^{\mathbb{R}}} = M_Y$

Fundamentals of measure theory

s-finite measures

► μ_n **bounded**:

$$\mu_n(\mathbb{R}) < \infty$$

► μ **s-finite**:

$$\mu = \sum_n \mu_n, \mu_n \text{ bounded}$$

Randomisation Theorem

Every s-finite measure is a push-forward of Lebesgue:

$$\mu \text{ s-finite} \implies \mu = f_*\lambda \text{ for some } f : \mathbb{R} \rightarrow \mathbb{R}_\perp$$

Transfer principle

$$\tau_*\lambda = \lambda \otimes \lambda \text{ for some measurable } \tau : \mathbb{R} \rightarrow (\mathbb{R} \times \mathbb{R})_\perp$$

Randomisation monad structure

▶ $(X_{\perp})^{\mathbb{R}}$

▶ $\text{return}_X(x) : r \in [0, 1] \mapsto x$

▶ $(\alpha \gg= f) : \mathbb{R} \xrightarrow{\tau} \mathbb{R} \times \mathbb{R} \xrightarrow{\mathbb{R} \times \alpha} \mathbb{R} \times X \xrightarrow{\mathbb{R} \times f} \mathbb{R} \times (Y_{\perp})^{\mathbb{R}} \xrightarrow{\text{eval}} Y$


$\mathbb{R} \rightarrow X_{\perp}$ $X \rightarrow (Y_{\perp})^{\mathbb{R}}$

▶ sample from randomisation of normal distribution

▶ $\text{score}(r) : r' \in [0, |r|] \mapsto ()$

monad laws fail
(associativity)

Lebesgue integration




A probabilistic powerdomain

$$\begin{array}{ccc} (X_{\perp})^{\mathbb{R}} & \xrightarrow{\int} & \mathbb{L}^X \\ & \searrow \quad \swarrow & \\ & = & \\ & MX & \end{array}$$

MX : randomisable integration operators

A probabilistic powerdomain



$(\mathcal{E}, \mathcal{M}) := (\text{densely strong epi, full mono})$ factorisation system

A probabilistic powerdomain

\mathcal{E} = densely strong epis closed under:

▶ products:

$$e_1, e_2 \in \mathcal{E} \implies e_1 \times e_2 \in \mathcal{E}$$

▶ lifting:

$$e \in \mathcal{E} \implies e_{\perp} \in \mathcal{E}$$

▶ random elements:

$$e \in \mathcal{E} \implies e^{\mathbb{R}} \in \mathcal{E}$$

$\implies M$ strong monad for sampling + conditioning



[Kammar-McDermott'18]

A probabilistic powerdomain

$$(X_{\perp})^{\mathbb{R}} \begin{array}{c} \xrightarrow{\quad} \mathbb{L}^X \\ \searrow \quad \nearrow \\ \quad \quad \quad = \quad \quad \quad \\ \quad \quad \quad MX \end{array}$$

► M locally continuous \implies may appear in domain equations


► M commutative

► M models synthetic measure theory

$$M \sum_{n \in \mathbb{N}} X_n \cong \prod_{n \in \mathbb{N}} MX_n$$

[Kock'12,
Ścibior et al.'18]

\implies satisfies Fubini



Valuations


Proposition

For standard Borel X :

$$MX \cong \left\{ \mu \mid \text{Scott opens} \mid \mu \text{ is s-finite} \right\}$$

Theorem (joint with Alex Simpson)

Quasi-Borel domains freely extend continuous domains:



Axiomatic domain theory

[Fiore-Plotkin'94, Fiore'96]

Structure

- ▶ Total map category: $\omega\mathbf{Qbs}$
- ▶ Admissible monos: **Borel-open** map $m : X \rightrightarrows Y$:

$$\forall \beta \in M_Y. \quad \beta^{-1}[m[X]] \in \mathcal{B}(\mathbb{R})$$

take Borel-Scott open maps as admissible monos

- ▶ **Pos**-enrichment: pointwise order
- ▶ Pointed monad on total maps: the powerdomain

\Rightarrow model axiomatic domain theory

\Rightarrow solve recursive domain equations

Axiomatic domain theory

Structure

\mathfrak{D} total map category
 $\omega\mathbf{Qbs}$
 $f \leq g$ **Pos**-enrichment
 pointwise order
 $\mathcal{M}_{\mathfrak{D}}$ admissible monos
 Borel-Scott opens
 T monad for effects
 power-domain
 m partiality encoding
 $m : -_{\perp} \rightarrow T, \perp \mapsto 0$

Derived axioms/structure

$\mathbf{p}\mathfrak{D}$ partial map category
 $-_{\perp}$ partiality monad
 (\dashv_V) the adjunction $J \dashv L$
 is locally continuous
 (\mathbf{p}_V) $\mathbf{p}\mathfrak{D}$ is $\omega\mathbf{Cpo}$ -enriched
 (1_{\leq}) $\mathbf{p}\mathfrak{D}$ has a partial terminal

Axioms

(\dashv) every object has a partial
 map classifier $\downarrow_X : X \rightarrow X_{\perp}$
 (fup) every admissible mono is full
 and upper-closed
 (\dashv_{\leq}) $\lfloor - \rfloor$ is locally monotone
 (\bigvee) \mathfrak{D} is $\omega\mathbf{Cpo}$ -enriched
 (U) ω -colimits behave uniformly
 (1) \mathfrak{D} has a terminal object
 (\rightarrow_{\leq}) \mathfrak{D} has locally monotone
 exponentials
 $(+)$ locally continuous total
 coproducts
 $(?!)$ $0 \rightarrow 1$ is admissible
 (\times_V) \mathfrak{D} has a locally
 continuous products
 (CL) \mathfrak{D} is cocomplete
 (T_V) T is locally continuous

(\otimes) $\mathbf{p}\mathfrak{D}$ has partial products
 (\otimes_V) (\otimes) is locally continuous
 (\rightarrow_V) \mathfrak{D} has locally continuous
 exponentials
 (\Rightarrow_V) $\mathbf{p}\mathfrak{D}$ has locally continuous
 partial exponentials
 $(\mathbf{p}CL)$ $\mathbf{p}\mathfrak{D}$ is cocomplete
 $(\mathbf{p}+_V)$ $\mathbf{p}\mathfrak{D}$ has locally continu-
 ous partial coproducts
 (BC) $J : \mathfrak{D} \hookrightarrow \mathbf{p}\mathfrak{D}$ is a bilimit
 compact expansion

Characterising $\omega\mathbf{Qbs}$

$$\begin{array}{ccc}
 \mathbf{Sbs} & \xrightarrow{\text{Yoneda}} & [\mathbf{Sbs}^{\text{op}}, \mathbf{Set}]_{\text{cpp}} \\
 & \searrow \text{Yoneda} \quad \quad \nearrow & \\
 & \text{SepSh} &
 \end{array}$$

countable product preserving [Staton et al.'16]

$F : \mathbf{Sbs}^{\text{op}} \rightarrow \mathbf{Set}$ separated: $F\mathbb{R} \xrightarrow{(F(\mathbb{R} \xleftarrow{r} \mathbb{1}))} (F\mathbb{1})^{\mathbb{R}}$ injective

Thm: $\mathbf{Qbs} \simeq \text{SepSh}$

→ [Heunen et al.'17]

$$\begin{array}{ccc}
 \mathbf{Sbs} & \xrightarrow{\text{Yoneda}} & [\mathbf{Sbs}^{\text{op}}, \omega\mathbf{Cpo}]_{\text{cpp}} \\
 & \searrow \text{Yoneda} \quad \quad \nearrow & \\
 & \omega\text{SepSh} &
 \end{array}$$

$$\begin{array}{l}
 f(x) \leq f(y) \\
 \implies x \leq y
 \end{array}$$

$F : \mathbf{Sbs}^{\text{op}} \rightarrow \omega\mathbf{Cpo}$ ω -separated: $F\mathbb{R} \xrightarrow{(F(\mathbb{R} \xleftarrow{r} \mathbb{1}))} (F\mathbb{1})^{\mathbb{R}}$ full

Thm: $\omega\mathbf{Qbs} \simeq_{\omega\mathbf{Cpo}} \omega\text{SepSh}$

Characterising $\omega\mathbf{Qbs}$

Grothendieck quasi-topos \mathbf{Qbs}
strong subobject classifier:

$$\underline{\Omega} = 2$$

$$M_{\Omega} = 2^{\mathbb{R}}$$

strong monos:

$$\begin{aligned} X &\xrightarrow{f} Y \\ (f \circ)^{-1}[M_Y] &= M_X \end{aligned}$$

Internal ω -cpo P :

$$\begin{array}{c} P^2 \xrightarrow{\leq_P} \Omega \\ \text{qbs} \uparrow \quad \omega\text{-chain}(P) \xrightarrow{\bigvee} P \\ (\underline{P}, \leq_P, \bigvee) \end{array}$$

+ internal quasi-topos logic ω -cpo axioms

Theorem

$$\omega\mathbf{Qbs} \simeq \omega\mathbf{Cpo}(\mathbf{Qbs})$$

Characterising $\omega\mathbf{Qbs}$

By local presentability:

$$\omega\mathbf{Cpo} \simeq \text{Mod}(\omega\mathbf{cpo}, \mathbf{Set})$$

$$\mathbf{Qbs} \simeq \text{Mod}(\mathbf{qbs}, \mathbf{Set})$$

essentially algebraic theories



$\omega\mathbf{qbs}$:

$$\omega\mathbf{cpo} \cup \mathbf{qbs} \cup \text{compatibility axiom}$$

Theorem

$$\omega\mathbf{Qbs} \simeq \text{Mod}(\omega\mathbf{qbs}, \mathbf{Set})$$

so $\omega\mathbf{Qbs}$ locally presentable, hence cocomplete

Summary

Contribution

- ▶ $\omega\mathbf{Qbs}$: a category of pre-domain quasi-Borel spaces
- ▶ M : commutative probabilistic powerdomain over $\omega\mathbf{Qbs}$
- ▶ Axiomatic treatment of measure and domain theory in $\omega\mathbf{Qbs}$
- ▶ Adequacy: $(\omega\mathbf{Qbs}, M)$ adequately interprets:
 - ▶ Statistical FPC
 - ▶ Untyped Statistical λ -calculus

[Fiore-Plotkin'94, Fiore'96]


This talk

- ▶ $\omega\mathbf{Qbs}$
- ▶ A probabilistic powerdomain
- ▶ Axiomatic treatment
- ▶ Characterising $\omega\mathbf{Qbs}$

Also in the paper

- ▶ Axiomatic domain theory
- ▶ Operational semantics
à la [Borgström et al.'16]

ProbProg: Important Language Features

Church  WebPPL Venture	sample	\mathbb{R}	score	higher order	term rec	type rec	Fubini (commute)
sets + probability	✓	✗	✗	✓	✗	✗	✓
meas space + subprobability	✓	✓	✗	✗	1 st	✗	✓
CPO + subprobability	✓	✓	✗	✓	✓	✓	?
cont domain + subprobability [Jones-Plotkin'89]	✓	✓	✗	✗	1 st	✗	✓
⋮ [Jung-Tix'98]	⋮	⋮	⋮	⋮	⋮	⋮	⋮
meas + s-finite distributions [Staton'17]	✓	✓	✓	✗	1 st	✗	✓
qbs + s-finite distributions [Heunen et al'17, Ścibior et al'18]	✓	✓	✓	✓	1 st	✗	✓
coh/meas cone + probability [Ehrhard-Pagani-Tasson'18, Ehrhard-Tasson'15-'19]	✓	✓ ✗	✗	✓	✓	? ✓	? ✓
ω qbs + s-finite distributions [This work]	✓	✓	✓	✓	✓	✓	✓



Sorts/arities

elem

ineq

Operations

lower : ineq \rightarrow elem

upper : ineq \rightarrow elem

refl : elem \rightarrow ineq

irrel : ineq \times ineq \rightarrow ineq

Def(irrel(e_1, e_2)):

lower (e_1) = lower (e_2)

upper(e_1) = upper(e_2)

antisym : ineq \times ineq \rightarrow elem

Def(antisym(e, e^{op})):

lower (e) = upper(e^{op})

upper(e) = lower (e^{op})

trans : ineq \times ineq \rightarrow ineq

Def(trans(e_1, e_2)):

upper(e_1) = lower(e_2)

Axioms

$e_1 = \text{irrel}(e_1, e_2) = e_2$

lower(refl(x)) = x = upper(refl(x))

lower(e_1) = antisym(e_1, e_2) = lower(e_2)

lower(trans(e_1, e_2)) = lower(e_1)

upper(trans(e_1, e_2)) = upper(e_2)

Presenting $\omega\mathbf{Cpo}$

Add to **pos**:

Operations

$$\bigvee : \prod_{n \in \mathbb{N}} \text{ineq} \rightarrow \text{elem}$$

$$\text{ub}_k : \prod_{n \in \mathbb{N}} \text{ineq} \rightarrow \text{ineq}$$

$$\text{lst} : \text{elem} \times \prod_{n \in \mathbb{N}} \text{ineq} \times \prod_{n \in \mathbb{N}} \text{ineq} \rightarrow \text{ineq}$$

Axioms

$$\text{lower}(\text{ub}_k(e_n)) = \text{lower}(e_k)$$

$$\text{upper}(\text{ub}_k(e_n)) = \bigvee(e_n)$$

$$\text{lower}(\text{lst}(x, (e_n), (b_n))) = \bigvee(e_n) \quad \text{upper}(\text{lst}(x, (e_n), (b_n)_n)) = x$$

$$\begin{aligned} \text{Def}(\bigvee_{n \in \mathbb{N}} e_n): \\ \text{upper}(e_n) &= \text{lower}(e_{n+1}) \\ \text{for each } n &\in \mathbb{N} \end{aligned}$$

$$\begin{aligned} \text{Def}(\text{ub}_k(e_n)_{n \in \mathbb{N}}): \\ \text{upper}(e_n) &= \text{lower}(e_{n+1}) \\ \text{for each } n &\in \mathbb{N} \end{aligned}$$

$$\begin{aligned} \text{Def}(\text{lst}(x, (e_n), (b_n))): \\ \text{upper}(e_n) &= \text{lower}(e_{n+1}) \\ \text{upper}(b_n) &= x \\ \text{lower}(e_n) &= \text{lower}(b_n) \\ \text{for each } n &\in \mathbb{N} \end{aligned}$$

Presenting Qbs

Sorts/arities

elem

rand

Operations

$\text{ev}_r : \text{rand} \rightarrow \text{elem}$

$\text{const} : \text{elem} \rightarrow \text{rand}$

$\text{rearrange}_\varphi : \text{rand} \rightarrow \text{rand}$

$\text{match}_{(S_i)_{i \in I}} : \prod_{i \in I} \text{rand} \rightarrow \text{rand}$

$\text{ext} : \text{rand} \times \text{rand} \rightarrow \text{rand}$

$\text{Def}(\text{ext}(\alpha, \beta)):$
 $\text{ev}_r(\alpha) = \text{ev}_r(\beta)$
for each $r \in \mathbb{R}$

Axioms

$$\alpha = \text{ext}(\alpha, \beta) = \beta$$

$$\text{ev}_r(\text{const}(x)) = x$$

$$\text{ev}_r(\text{rearrange}_\varphi \alpha) = \text{ev}_{\varphi(r)} \alpha$$

$$\text{ev}_r \left(\text{match}_{(S_i)_{i \in I}} (\alpha_i)_{i \in I} \right) = \text{ev}_r(\alpha_i)$$

Presenting ω Qbs

Sorts/arities

elem

ineq

rand

Operations

Add to ω cpo and qbs:

$\sqcup : \prod_{n \in \mathbb{N}} \text{rand} \times \prod_{n \in \mathbb{N}, r \in \mathbb{R}} \text{ineq} \rightarrow \text{rand}$

Def $\sqcup((\alpha_n)_{n \in \mathbb{N}}, (e_n^r)_{n \in \mathbb{N}, r \in \mathbb{R}})$:

$$\text{lower}(e_n^r) = \text{ev}_r(\alpha_n) \quad \text{upper}(e_n^r) = \text{ev}_r(\alpha_{n+1})$$

for each $n \in \mathbb{N}$, $r \in \mathbb{R}$

Axioms

Add:

$$\text{ev}_r \left(\sqcup \left((\alpha_n)_{n \in \mathbb{N}}, (e_n^r)_{n \in \mathbb{N}, r \in \mathbb{R}} \right) \right) = \bigvee (e_n^r)_{n \in \mathbb{N}}$$