

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

Logic Summer School
Australian National University
4–16 December, 2023
Canberra, ACT, Australia



THE UNIVERSITY OF EDINBURGH

informatics IfCS

Laboratory for Foundations
of Computer Science



supported by:



THE ROYAL
SOCIETY



Facebook Research NCSC

Computational golden era of:

logic & type rich
computation

Statistical
computation

Computational golden era of:

logic & type rich
computation

Expressive type systems:

Haskell, OCaml, Idris

Mechanised mathematics:

Agda, Coq, Isabelle/Hol, Lean

Verification:

SMT-powered, realistic
systems

Statistical
computation

generative modelling
+

efficient inference:

Monte-Carlo simulation
or gradient-based
optimisation

"AI"

Computational golden era of:

logic & type rich
computation


Statistical
computation

Clear connection to

Foundations:

- Rault's
- John's courses
- Michael's
- Dominik's
- this course

Why foundations?



Plan:

- 1) type-driven Probability: discrete case (Mon + Tue (?))
- 2) Borel sets & measurable spaces (Tue)
- 3) Quasi Borel spaces, Simple type structure (Wed)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

Please ask questions!

Smibble



Course
Web
Page

Advertisement

Interested in the mathematical } computational
logical } foundations of Computer
Science ?

Hedging your bets on a funded PhD offer from ANU?

Check out our PhD Programs:



LFCS
PhD



Ifcs

Laboratory for Foundations
of Computer Science



Reversible
AI for
Robotics
CDT

Language of distribution & Probability

X type (=space) of values / outcomes

$\mathcal{D}X$ type of distributions / measures over X

$\mathcal{P}X \subseteq \mathcal{D}X$ sub type of probability measures (total measure)

$\mathcal{B}X$ type of measurable events - Subsets of X we wish to measure

\mathbb{W} type of weights : $[0, \infty]$

→ type judgment

$\mu : \mathcal{D}X, E : \mathcal{B}X \vdash c_e[E] : \mathbb{W}$

↳ measure μ assigns to E

Axioms for measures

Empty event : $\emptyset : \mathcal{B}X$

Its measure is $0 : \mathbb{W}$:

$$\mu : \mathcal{D}X \vdash \underset{\mu}{\text{Ce}}[\emptyset] = 0 : \mathbb{W}$$

Axioms for measures

BX is a Boolean Sub-algebra:

$$E : BX \vdash E^c : BX$$

$$E, F : BX \vdash E \cup F, E \cap F : BX$$

$$E, C : BX, \mu : DX \vdash \quad (\text{disjoint additivity})$$

$$\underset{\mu}{\text{Ce}}[E] = \underset{\mu}{\text{Ce}}[E \cap C] + \underset{\mu}{\text{Ce}}[E \cap C^c] : W$$

Axioms for measures

$\omega := (\mathbb{N}, \leq)$ (B, \subseteq) (W, \leq) posets

$$(BX, \subseteq)^\omega := \left\{ (E_n)_{n \in \mathbb{N}} \in (BX)^\mathbb{N} \mid E_0 \subseteq E_1 \subseteq E_2 \subseteq \dots \right\}$$

(BX, \subseteq) and (W, \leq) are ω -chain-closed:

$$E_- : (BX, \subseteq)^\omega \vdash \bigvee_n E_n : BX \quad \alpha_- : (W, \leq)^\omega \vdash \sup_n \alpha_n : W$$

$$E_- : (BX, \subseteq)^\omega, \mu : D_X \vdash \quad \text{(Scott Continuity)}$$

$$\text{Ce}_{\mu} \left[\bigvee_n E_n \right] = \sup_n \text{Ce}_{\mu} [E_n] : W$$

Axiom for Probability

$$\text{Cast} : \text{PX} \xleftarrow{\leq} \text{DX}$$

$$1 : \mathbb{W}$$

$$\mu : \text{PX} \vdash \text{Ce}[X] = 1 : \mathbb{W}$$

Cast μ

Avoid casting:

$$E : BX, \mu : \text{PX} \vdash \Pr_{\Gamma}[E] := \text{Ce}[E] : [0,1] \subseteq \mathbb{W}$$

Cast μ

Axioms for measures

Integration:

$$\mu : \mathbf{DX}, \varphi : \mathbb{W}^X \vdash \int_\mu \varphi : \mathbb{W} \quad (\text{Lebesgue integral})$$

Again, avoid casting:

$$\mu : \mathbf{PX}, \varphi : \mathbb{W}^X \vdash \mathbb{E}_{\mu}[\varphi] := \int_{\mu} (\text{cast } \mu) \varphi : \mathbb{W} \quad (\text{Expectation})$$

More structure & notation later (...technical...)

Have: language + axioms

Want: model

today: discrete measures

rest of course: discrete + continuous

Discrete model

type X : set

$$DX := \{ \mu : X \rightarrow \mathbb{W} \mid \mu \text{ is Countably Supported} \}$$

(next slide)

Support

Power set

$\mu : \mathbb{W}^X, S : \mathcal{P}X \vdash S \text{ supports } \mu :=$

$\forall x : X. \mu x > 0 \Rightarrow x \in S : \text{Prop}$

$\mu : \mathbb{W}^X \vdash \text{Supp } \mu := \{x \in X \mid \mu x > 0\} : \mathcal{P}X$

$\text{Supp } \mu$ is the smallest set supporting μ

Discrete model

type X : set

$$DX := \{ \mu : X \rightarrow \mathbb{W} \mid \mu \text{ is Countably Supported} \}$$

$$:= \{ \mu : X \rightarrow \mathbb{W} \mid \text{Supp } \mu \text{ is Countable} \}$$

Ex. measures

- X ctbl. Counting measure $\#_X : DX$
 $\#_X := \lambda x : X. 1$ (NB: $\text{Supp } \#_X = X \sqrt{\text{ctbl}}$)
- Dirac measure:
 $\sigma : X \vdash \delta_x := \lambda x'. \begin{cases} x = x' : 1 \\ \text{o.w.} : 0 \end{cases} : DX$
NB: $\text{Supp } \delta_x = \{x\} \sqrt{\text{ctbl}}$
- Zero measure $\underline{0} := \lambda x. 0 : DX$
NB: $\text{Supp } \underline{0} = \emptyset \sqrt{\text{ctbl}}$

Discrete model

type X : set

$DX := \{ \mu : X \rightarrow \mathbb{W} \mid \mu \text{ is Countably Supported} \}$

$$\mu : DX, E : BX \vdash C_E[\mu] := \sum_{x \in E} \mu x$$

$$:= \sum_{x \in E \cap \text{Supp } \mu} \mu x$$

Lemma: $\mu : DX, S \in \mathcal{P}_{\text{ctbl}}^X, S \text{ supports } \mu, E : BX \vdash$

$$C_E[\mu] = \sum_{x \in E \cap S} \mu x$$

Ex:

- $E : B X \vdash$ $C_e[E] = |\underset{\#_x}{E}| := \begin{cases} E \text{ has } n \text{ elements: } n \\ E \text{ infinite: } \infty \end{cases}$

- $E : B X, n : X \vdash C_e[E] = \underset{\delta_n}{\delta_{\in E}} = \begin{cases} x \in E : 1 \\ x \notin E : 0 \end{cases} =: [x \in E] : \mathbb{W}$

NB: $E : B X \vdash [- \in E] : X \rightarrow \mathbb{W}$

indicator
function

- $E : B X \vdash C_e[E] = \underset{\Omega}{\Omega}$

Validate axioms

$$\mu : \text{DX} \vdash \underset{\mu}{\text{Ce}}[\emptyset] = 0 : \mathbb{W}$$

$$E, C : \text{BX}, \mu : \text{DX} \vdash$$

$$\underset{\mu}{\text{Ce}}[E] = \underset{\mu}{\text{Ce}}[E \cap C] + \underset{\mu}{\text{Ce}}[E \cap C^c] : \mathbb{W}$$

$$E_- : (\text{BX}, \subseteq)^\omega, \mu : \text{DX} \vdash$$

$$\underset{\mu}{\text{Ce}}[\bigvee E_n] = \sup_n \underset{\mu}{\text{Ce}}[E_n] : \mathbb{W}$$

Kernels κ from Γ to X :

$$\kappa : (DX)^\Gamma$$

kernels are "open/parameterised" measures

Ex: Dirac kernel: $\delta_+ : (DX)^X$

Kock Integral

$$\mu : D\Gamma, \kappa : DX \vdash \int^\Gamma \mu \kappa : DX$$

In discrete model:

$$\int^\Gamma \mu \kappa := \lambda x : X. \sum_{n \in \Gamma} \underbrace{\mu n \cdot k(n; x)}_{:= h \vdash x}$$

(Weak) disintegration problem:

Input: $\mu: D\Gamma$ $V: DX$

Output: a kernel $k:(DX)^{\Gamma}$ s.t.

$$\oint \mu k = V$$

Call such $k \stackrel{a}{=} (\text{weak}) \text{ disintegration of } V$

w.r.t. μ .

(non-standard
terminology)

Ex disintegration:

$$\underline{n} := \{0, 1, 2, \dots, n-1\}$$

disintegrate $\#_{\geq \frac{n+1}{2}}$ w.r.t. $\#_{\geq}$

$$k: \left(D(\underline{\mathbb{Z}}^{\frac{n+1}{2}})\right)^2$$
$$k(x_j; f) := \begin{cases} f(n) = x: & 1 \\ \text{o.w.} & 0 \end{cases}$$

$$\left(\oint \#_{\geq} k\right) f = \sum_{n \in \underline{\mathbb{Z}}} \overset{1}{\#_{\geq} x} \cdot k(n; f)$$

NB: $\text{Supp}(ux)$
 $\sqrt{c+b}$

$$= k(0; f) + k(1; f) = u(f_n; f) = 1 = \#_{\geq \frac{n+1}{2}}(f)$$

Probability measures

$$P_X := \left\{ \mu : D_X \mid \underset{\mu}{\text{C}_e}[X] = 1 \right\} \hookrightarrow^{\subseteq} D_X$$

Lemma: $\delta_- : X \rightarrow D_X$ and $\oint : D\Gamma \times (D_X)^r \rightarrow D_X$

lift along the inclusion cast: $P \hookrightarrow^{\subseteq} D$:

$$\begin{array}{ccc} X & \xrightarrow{\delta_-} & P_X \\ & \dashv & \downarrow \text{cast} \\ & \xrightarrow{\delta_-} & D_X \end{array}$$

$$\begin{array}{ccc} P\Gamma \times (P_X)^r & \dashv \oint \dashrightarrow & P_X \\ \text{cast} \times (\text{cast}) \downarrow & & \downarrow \text{cast} \\ D\Gamma \times (D_X)^r & \xrightarrow{\oint} & D_X \end{array}$$

Prop (discrete Giry):

(Michèle Giry '82)

(P, δ_-, \oint) is a monad i.e.

$$m : \Gamma, n : (Dx)^\Gamma \vdash \oint \delta_n k = k \ r$$

$$\mu : D X \vdash \oint \mu(\lambda x) \delta_x = \mu : D X$$

$$\mu : D\Gamma, \kappa : (Dx)^\Gamma, t : (DY)^X \vdash$$

$$\oint \mu(\lambda x) \left(\oint (\kappa r) t \right) = \oint \left(\oint \mu \kappa \right) (\lambda x) t(x)$$

Corollary: (P, δ_-, \oint) is a monad.

Language of distribution & Probability

X type (=space) of values / outcomes

$\mathcal{D}X$ type of distributions / measures over X

$\mathcal{P}X \subseteq \mathcal{D}X$ sub type of probability measures (total measure)

$\mathcal{B}X$ type of measurable events - Subsets of X we wish to measure

\mathbb{W} type of weights : $[0, \infty]$

→ type judgment

$\mu : \mathcal{D}X, E : \mathcal{B}X \vdash c_e[E] : \mathbb{W}$

↳ measure μ assigns to E

Plan:

- 1) type-driven Probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Tue)
- 3) Quasi Borel spaces, Simple type structure (Wed)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

Please ask questions!

Smibble



Course
Web
Page

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

Logic Summer School
Australian National University
4–16 December, 2023
Canberra, ACT, Australia



THE UNIVERSITY OF EDINBURGH

informatics IfCS

Laboratory for Foundations
of Computer Science



supported by:



THE ROYAL
SOCIETY



Facebook Research NCSC

Language of distribution & Probability

Recap

X type (=space) of values / outcomes

$\mathcal{D}X$ type of distributions / measures over X

$\mathcal{P}X \subseteq \mathcal{D}X$ Sub type of probability measures (total measure)

$\mathcal{B}X$ type of measurable Events - Subsets of X we wish to measure

\mathbb{W} type of weights : $[0, \infty]$

→ type judgment

$\mu : \mathcal{D}X, E : \mathcal{B}X \vdash c_e[E] : \mathbb{W}$

↳ measure μ assigns to E

Axioms for measures/distributions

Recap

$$\mu : \mathbf{D}X \vdash \underset{\mu}{\text{Ce}}[\emptyset] = 0 : \mathbb{W}$$

$$E, C : \mathbf{B}X, \mu : \mathbf{D}X \vdash$$

$$\underset{\mu}{\text{Ce}}[E] = \underset{\mu}{\text{Ce}}[E \cap C] + \underset{\mu}{\text{Ce}}[E \cap C^c] : \mathbb{W}$$

$$E_- : (\mathbf{B}X, \subseteq)^\omega, \mu : \mathbf{D}X \vdash$$

$$\underset{\mu}{\text{Ce}}[\bigvee_n E_n] = \sup_n \underset{\mu}{\text{Ce}}[E_n] : \mathbb{W}$$

Kernels & their Koch integral

Recap

kernel from Γ to X : $k: (DX)^\Gamma$ or $k: \Gamma \rightarrow DX$

Dirac kernel: $\delta_- : X \rightarrow DX$

Koch integral: $\mu: D\Gamma$, $k: (DX)^\Gamma \vdash \oint \mu k : DX$
or $\oint \mu(dx) \kappa(x)$ (*dx binding occurs in $\kappa(x)$*)

Giry monads: $(D, \delta_-, \oint) \dashv (P, \mathcal{S}_-, \oint)$.

Discrete model

$$\text{type} : \text{set} \quad W := [0, \infty] \quad \mathcal{B}X := \mathcal{P}X$$

$$DX := \{\mu : X \rightarrow W \mid \text{Supp } \mu \text{ countable}\}$$

$$PX := \left\{ \mu \in DX \mid \underset{\mu}{\text{Ce}}[X] = 1 \right\}$$

$$\underset{\mu}{\text{Ce}}[E] := \sum_{x \in E} \mu_x \quad \delta_x := \lambda x'. \begin{cases} x = x' : 0 \\ x \neq x' : 1 \end{cases}$$

$$\phi \mu k := \lambda x. \sum_{m \in \Gamma} \mu^m \cdot k(m; x)$$

Ex distributions

Counting measure (λ_{ctbl}): $\#_X := \lambda_X \cdot 1$

Dirac measure δ_x (prev slide)

Zero measure $\underline{\varrho} := \lambda_X \cdot 0$

Plan:

- 1) type-driven Probability: discrete case (Mon + Tue)
- 2) Borel sets & measurable spaces (Tue)
- 3) Quasi Borel spaces, Simple type structure (Wed)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

Please ask questions!

Smibble



Course
Web
Page

Product measures

$$\mu: D X, \nu: D Y \vdash \mu \otimes \nu := \int \mu(dx) \int \nu(dy) \delta_{(x,y)} : D(X \times Y)$$

(\otimes lifts along $P \hookrightarrow D$)

$$= \lambda(x,y). \mu x \cdot \nu y$$

discrete model

$$E_{\#} : \#_{X \times Y} = \#_X \otimes \#_Y$$

Indeed:

$$(\# \otimes \#)(x,y) = \#x \cdot \#y = 1 \cdot 1 = 1 = \#(x,y)$$

build measures
compositionally

$$\text{Notation: } \lambda : D(X \times Y), \kappa : (DZ)^{X \times Y} \vdash \oint \lambda(\Delta z, dy) \kappa(z, y) \\ := \oint \lambda \kappa$$

Fubini - Tonelli Thm:

Integrate in any order:

$$\mu : DX, \nu : DY, \kappa : (DZ)^{X \times Y} \vdash$$

$$\oint \mu(dx) \oint \nu(dy) \kappa(x, y) = \oint (\mu \otimes \nu)(dx, dy) \\ = \oint \nu(dy) \oint \mu(dx) \kappa(x, y)$$

Pushing a measure forward

$$\mu: D_{\Omega}, d: X^{\Omega} \vdash \mu_f := \phi \mu(d\omega) \delta_{\alpha\omega} : DX$$

$$= \lambda x. \sum_{\omega \in \Omega} \mu \omega$$

$$\alpha\omega = x$$

$\alpha: X^{\Omega}$: random element

(w.r.t. μ)

$\mu_{\alpha}: DX$: the law of α

Ex: We can represent configurations of 2 dice using $\underline{6} \times \underline{6}$

Letting $(+): \underline{6}^2 \rightarrow \mathbb{N}^2 \xrightarrow{(+)} \mathbb{N}$

we have that the law of $(+)$:

$$(\#_{\underline{6}} \otimes \#_{\underline{6}})_{(+)} : \mathbb{D}/\mathbb{N}$$

is the number of rolls whose sum is given

build measures
compositionally

Scaling a measure

$$(\cdot) : \mathbb{W} \times D_X \longrightarrow D_X$$

$$a \cdot \mu := \lambda x. a \cdot \mu x$$

$$\boxed{NB: \text{Supp}(a \cdot \mu) = \begin{cases} a=0: \emptyset \\ a \neq 0: \text{Supp } \mu \end{cases}}$$

\checkmark_{c+61}

$(\cdot) : \mathbb{W} \times D_X \rightarrow D_X$ is an action of monoid $(\mathbb{W}, (\cdot), 1)$ on D_X :


$$\mu : D_X \vdash$$

$$1 \cdot \mu = \mu$$

$$a, b : \mathbb{W}, \mu : D_X \vdash$$

$$a \cdot (b \cdot \mu) = (a \cdot b) \cdot \mu$$

Normalisation



$\mu : D X, C_C[X] \neq 0, \infty +$

$$\|\mu\| := \left(\frac{1}{C_C[X]} \right) \cdot \mu : P X$$

Ex:

$$\emptyset \neq A \subseteq_{fin} X : U_{A \subseteq X} := \|\#_A\|_{A \subseteq X} : P X$$

$$1 \xrightarrow{\#_A} D A \xrightarrow{(-)_{A \subseteq X}} D X \xrightarrow{\|\cdot\|} P X$$

I.e.

$$U_{A \subseteq X} := \lambda n. \begin{cases} n \in A : \frac{1}{|A|} \\ n \notin A : 0 \end{cases}$$

so

$$\bigcup_{n \in A} = \delta_n$$

Standard vocabulary

Joint distributions:

$$\mu : D(X_1 \times X_2)$$

Marginal distribution:

$$X_1 \xleftarrow{\pi_1} X_1 \times X_2 \xrightarrow{\pi_2} X_2$$

law of projection

$$\mu_{\pi_i} : D X_i$$

Marginalisation: $\mu_{\pi_i} = \oint \mu(dx, dy) S_x$

integrate out y

Exercise: $\mu : P X, V : D x \vdash (\mu \otimes V)_{\pi_2} = V$

independence

Pairing R.E.S:


$$\alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash$$

$$\langle \alpha, \beta \rangle := \lambda w. \langle \alpha w, \beta w \rangle : (X \times Y)^{\Omega}$$

$$\lambda : D\Omega, \alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash \alpha \perp \beta := \lambda_{\langle \alpha, \beta \rangle} = \lambda_{\alpha} \oplus \lambda_{\beta}$$

: Prop

α, β independent w.r.t. λ



Ex^(Durrett) represent Outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : \bigcup_{c \in C} \bigcup_{c \in C} \bigcup_{c \in C} : P_\Omega$$

$$\pi_i : \Omega \rightarrow C \quad \text{Outcome of } i^{\text{th}} \text{ toss}$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{? (=)} \mathbb{B}$$

where : $(?) : C^2 \rightarrow \mathbb{B} := \{\text{True}, \text{False}\}$

$$?_{x=y} := \begin{cases} x=y : \text{True} \\ x \neq y : \text{False} \end{cases}$$

Ex ^(Durrett) represent Outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : U_C \otimes U_C \otimes U_C : P_{\Omega}$$

$$\pi_i : \Omega \rightarrow C \quad \text{Outcome of } i^{\text{th}} \text{ toss}$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{?} B$$

marginalisation

$$\lambda_{\text{Same}_{12}}^T = (U_C \otimes U_C)^T \stackrel{?}{=} \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$
$$\begin{matrix} U_C(T) \cdot U_C(T) \\ \downarrow \\ \frac{1}{4} \\ \uparrow \\ U_C(H) \cdot U_C(H) \end{matrix}$$

$$\text{so } \lambda_{\text{Same}_{12}}^F = \frac{1}{2} \text{ too}$$

Ex ^(Durrett) represent Outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : \bigcup_{C^3} \otimes \bigcup_{C^3} \otimes \bigcup_{C^3} : P_\Omega$$

$$\pi_i : \Omega \rightarrow C \quad \text{Outcome of } i^{\text{th}} \text{ toss}$$

$$\underline{i \neq j} : \lambda_{\text{Same}_{ij}} = V_B$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{?} B$$

$$\lambda : \begin{matrix} (T, T) \mapsto \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} \\ \langle \text{Same}_{12}, \text{Same}_{23} \rangle \end{matrix} \hookrightarrow \lambda(H, H, H) \quad \hookrightarrow \lambda(T, T, T)$$

$$(T, F) \mapsto \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$$

$$\hookrightarrow \lambda(H, H, T) \quad \hookrightarrow \lambda(T, T, H)$$

Ex^(Durrett) represent Outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : U_C \otimes U_C \otimes U_C : P_{\Omega}$$

$$\pi_i : \Omega \rightarrow C \quad \text{Outcome of } i^{\text{th}} \text{ toss}$$

$$\underline{i \neq j} \quad \lambda_{\text{Same}_{ij}} = V_{IB}$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{?} IB$$

$$\lambda_{\langle \text{Same}_{12}, \text{Same}_{23} \rangle} = V_{IB \times IB} = V_{IB} \otimes V_{IB} = \lambda_{\text{Same}_{12}} \otimes \lambda_{\text{Same}_{13}}$$

$$\text{So } \text{Same}_{12} \perp \lambda \text{ Same}_{13}$$

independence

Pairing R.E.S:

$$\alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash$$

$$\langle \alpha, \beta \rangle := \lambda w. \langle \alpha w, \beta w \rangle : (X + Y)^{\Omega}$$

$$\lambda : D\Omega, \alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash \alpha \perp_{\lambda} \beta := \lambda_{\langle \alpha, \beta \rangle} = \lambda_{\alpha} \otimes \lambda_{\beta} : \text{Prop}$$

α, β independent w.r.t. λ

I-ary version:

$$\lambda : D\Omega, \alpha_i : \prod_{i \in I} X_i^{\Omega} \vdash \perp_{\lambda, i \in I}^{\alpha_i} :=$$

α_i independent
w.r.t. λ

$$\forall J \subseteq_{\text{fin}} I. \quad \lambda_{\langle \alpha_j \rangle_{j \in J}} = \bigotimes_{j \in J} \lambda_{\alpha_j} : \text{Prop}$$

Ex ^(Durrett) represent outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega = C \times C \times C \quad \lambda : \bigcup_{C^3} \otimes \bigcup_{C^3} \otimes \bigcup_{C^3} : P_\Omega$$

$$\pi_i : \Omega \rightarrow C \quad \text{Outcome of } i^{\text{th}} \text{ toss}$$

$$\underline{i \neq j} : \lambda_{\text{Same}_{ij}} = V_{\mathbb{B}}$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{?} \mathbb{B}$$

$$\begin{matrix} i \neq j \\ * \\ n \end{matrix} : \text{Same}_{ij} \perp \text{Same}_{jk}$$

$$\frac{1}{\lambda} \left\{ \text{Same}_{12}, \text{Same}_{23}, \text{Same}_{13} \right\}$$

$$\text{Intuition: Same}_{13} = \text{IFF} (\text{Same}_{12}, \text{Same}_{23})$$

Calc:

$$\begin{aligned} \lambda_{\langle \text{Same}_{12}, \text{Same}_{23}, \text{Same}_{13} \rangle} (T, T, T) &= \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} \neq \frac{1}{2^3} = \lambda_{\text{Same}_{12}} \otimes \lambda_{\text{Same}_{23}} \otimes \lambda_{\text{Same}_{13}} \\ &\hookrightarrow \lambda(H, H, H) \quad \hookrightarrow \lambda(T, T, T) \end{aligned}$$

Vocabulary

(Discrete) Measure Space $(X, \mu : D_X)$

measure preserving $f : (X, \mu) \rightarrow (Y, \nu)$

function $f : X \rightarrow Y$ s.t. $\mu_f = \nu$

$\mu : D_X$, $f : X \rightarrow Y \vdash \mu$ invariant under $f :=$

$f : (X, \mu) \rightarrow (Y, \nu)$

Ex:


$\mu : D_X, \nu : D_Y \vdash$

Swap : $(X \times Y, \mu \otimes \nu) \longrightarrow (Y \times X, \nu \otimes \mu)$ so


$\mu : D_X \vdash \mu \otimes \mu$ invariant under Swap

Weights as measures

Observation:



Proof:



□

NB: unit type \rightarrow $\mathbf{1} := \{\langle\rangle\}$ unit value

Integration

$$\mu: D_X, \varphi: W^X \vdash \int^\mu \varphi : W$$
$$:= \sum_{x \in X} \mu_x \cdot \varphi_x$$

(Lebesgue integral)

Can derive it:

$$D_X \times W^X \xrightarrow{D_X \times (\cong o-)} D_X \times (D_1)^X$$
$$\downarrow \int \qquad \qquad \qquad \vdash$$
$$W \leftarrow \cong \qquad \qquad \qquad \downarrow \varphi$$
$$D_1$$

Additivity:

$$\text{I ctsl, } \mu_-(DX)^I \vdash \sum_{i \in I} \mu_i : DX$$

$$:= \lambda x. \sum_{i \in I} \mu_i x$$

NB:
 $\text{supp} \sum_i \mu_i \subseteq$
 $\bigcup_i \text{supp } \mu_i$
 $\checkmark \text{ctsll}$

Ex: Bernoulli distribution

$$p:[0,1] \vdash B(p) := p \cdot \delta_{\text{True}} + (1-p) \cdot \delta_{\text{False}} : P/B$$

$$\text{i.e. } \beta_p : \begin{aligned} \text{True} &\mapsto p \\ \text{False} &\mapsto 1-p \end{aligned}$$

Thm (affine-linearity):

ϕ is affine-linear in each argument:

$I \vdash b : I$

$$m : (\mathcal{D}\Gamma)^I, k : (\mathcal{D}x)^I \vdash \phi\left(\sum_{i \in I} a_i \cdot \mu_i\right) k = \sum_{i \in I} a_i \cdot \phi \mu_i k$$

$I \vdash b : I$, $\mu : \mathcal{D}\Gamma$, $a : W^I$, $k : \mathcal{D}x^I$

$$\int \mu(dx) \left(\sum_{i \in I} a_i \cdot k_i(x) \right) = \sum_{i \in I} a_i \cdot \phi \mu k_i$$

Prop: $\mathbb{W} \cong D1$ is a σ -semi-ring isomorphism:

$$(\mathbb{W}, \Sigma, (\cdot), 1) \cong (D1, \Sigma, (\cdot), \delta_{\leq})$$

and $(\cdot) : \mathbb{W} \times Dx \rightarrow Dx$ makes Dx into a module:

$$\left(\sum_{i \in I} a_i \right) \cdot \mu = \sum_{i \in I} (a_i \cdot \mu) \quad a \cdot \sum_{i \in I} \mu_i = \sum_{i \in I} a \cdot \mu_i$$

Corollary: \int is affine-linear in each argument.

Random variable :

NB: $\bar{\mathbb{R}} := [-\infty, \infty]$

A random element $\alpha: \bar{\mathbb{R}}^\Omega$ (wrt some $\mu: D\mathcal{L}$)

Can add, multiply r.v.'s.

To integrate r.v.'s:

$$(-)^+: \bar{\mathbb{R}}^\Omega \longrightarrow \mathbb{W}^\Omega$$

$$\alpha^+ := \lambda w. \begin{cases} \alpha \cdot w \geq 0 : \alpha w \\ 0.w : 0 \end{cases} = [\alpha \geq 0] \cdot |\alpha|$$

$$\alpha^- := \lambda w. \begin{cases} \alpha \cdot w \leq 0 : |\alpha w| \\ 0.w : 0 \end{cases} = [\alpha \leq 0] \cdot |\alpha|$$

So $\alpha = \alpha^+ - \alpha^-$

$\mu: D\Omega, \alpha: \overline{\mathbb{R}}^n, \int \mu \alpha^+ < \infty \text{ or } \int \mu \alpha^- < \infty +$

$$\int \mu \alpha := \int \mu \alpha^+ - \int \mu \alpha^- : \overline{\mathbb{R}}$$

Ex. The (discrete) Lebesgue p -space:

$$p \in [1, \infty), \mu: P\Omega \vdash L_p(\Omega, \mu) :=$$

$$\left\{ \alpha: \overline{\mathbb{R}}^n \mid \underset{\mu}{\mathbb{E}}[|\alpha|^p] < \infty \right\}$$

$L_p(\Omega, \mu)$ has a norm $\|\alpha\| := \sqrt[p]{\underset{\mu}{\mathbb{E}}[|\alpha|^p]}$ almost Banach

$L_2(\Omega, \mu)$ has an inner product $\langle \alpha, \beta \rangle := \underset{\mu}{\mathbb{E}}[\alpha \cdot \beta]$ almost Hilbert


Density

a density over X : $d : X \rightarrow W$

$$d : W^X, \mu : D_X \vdash d \odot \mu : D_X \\ := \oint \mu(dx) (dx \cdot \delta_x)$$

Warning The types of measures & densities in the discrete model are close, but still different. They coincide on countable sets, so people often confuse them. Types help us keep them separate.

Intuition:



Almost certain Properties

$$E : \mathcal{B}X, \mu : \mathcal{D}X \vdash \mu(\text{d}x) \text{-almost certainly } x \in E : \text{Prop}$$
$$:= [x \in E] \odot \mu = \mu$$
$$\text{NB: } [x \in E] = \begin{cases} x \in E : 1 \\ x \notin E : 0 \end{cases} : \mathbb{W}$$

When $\mu : \mathcal{P}X$ we say instead

$\mu(\text{d}x)$ -almost surely $x \in E$

Exercise Look up the def. of a normed space

and modify the definition so that $L_p(\Omega, \mu)$ is a normed space up-to almost sure equality.

Absolute continuity

$\mu, \nu : D^X, d : W^X \vdash d = \frac{d\mu}{d\nu} : \text{Prop}$

$$:= \mu = d \odot \nu$$

$\mu, \nu : D^X \vdash \mu \ll \nu := \mu \text{ is absolutely continuous w.r.t. } \nu : \text{Props}$

$$:= \exists d : W^X. \quad d = \frac{d\mu}{d\nu}.$$

$=: \mu \text{ has a density w.r.t. } \nu$

Lemma: $\mu, \nu : D^X,$
 $\mu \ll \nu,$
 $k : (D^Y)^X$

$$\oint V(dx) \frac{d\mu}{d\nu}(x) \cdot k_x = \oint \mu(dx) k_x$$

$$\underline{Ex}: \bigcup_{A \subseteq X} \ll (\#_A)_{\text{Cost}: A \subseteq X}$$

$$\frac{dV_{A \subseteq X}}{d(\#_A)_{\text{Cost}}} = \lambda x. \begin{cases} x \in A : & \frac{1}{|A|} \\ \text{D.W.} : & 0 \end{cases}$$

but also:

$$\frac{dV_{A \subseteq X}}{d(\#_A)_{\text{Cost}}} = \lambda x. \frac{1}{|A|}$$

Radon-Nikodym Thm: (discrete version)

$\mu, \nu : P X \vdash \mu \ll \nu$ iff $\forall x. \nu x = 0 \Rightarrow \mu x = 0$
i.e. $\text{Supp } \mu \subseteq \text{Supp } \nu$

In that case, if $d_1, d_2 = \frac{d\mu}{d\nu}$ then

$$\nu(dx)\text{-a.s. } d_1 x = d_2 x$$

Ex: for ctbl X , $\forall \mu : D X . \mu \ll \#_X$. Proof: vacuously, as $\#_X x \neq 0$.

Then $\lambda x. \mu x = \frac{d\mu}{d\#}.$


Conditional expectation

β is a conditional expectation of α wrt. μ along H

$$\mu: D\Omega, H: X^\Omega, \alpha: L_1(\Omega, \mu), \beta: L_1(X, \mu_H)$$

$$\vdash \beta = \mathbb{E}[\alpha | H = -] \quad : \text{Prop}$$

$$:= \forall \varphi: L_1(Y, \mu_H^M). \int \mu_H(d\omega) \beta(\omega) \cdot \varphi(\omega) = \int \mu(d\omega) \alpha(\omega) \cdot \varphi(H\omega)$$



Thm (Kolmogorov): (discrete version)

There is a function

$$\mathbb{E}_{\mu}[-|H=-] \in \prod_{\mu: P_{\Omega}} \prod_{H: X^{\omega}} \mathcal{L}_1(\Omega, \mu) \rightarrow \mathcal{L}_1(X, \mu_H)$$

s.t. $\mathbb{E}_{\mu}[\alpha | H=-]$ is a conditional expectation of α w.r.t. μ
along H .

Conditional Probability (discrete version):

$$\text{H}: x : \Omega, \mu : P_X \vdash \underset{\mu}{\mathbb{P}_r}[- \mid H = -] : (P_{\Omega})^X$$
$$:= \lambda x_0 : X. \lambda \omega_0 : \Omega. \underset{\omega \sim \mu}{\mathbb{E}} [\llbracket \omega_0 = w \rrbracket \mid H_w = x_0]$$

Bayes's Thm (discrete version, adapted from Williams):

Let $\lambda : P(X \times \Theta)$ joint probability distribution.

Assume $\mu : D_X$, $V : D_\Theta$ s.t. $\lambda \ll \mu \otimes V$.


with $d_{X,\Theta} = \frac{d\lambda}{d(\mu \otimes V)}$.

OBS 1: $d_X : W^X$

$$d_X := \lambda_{\Theta} \int V(d\Theta) d_{X,\Theta}(x, \theta)$$

then $d_X = \frac{d\lambda}{d\mu}$

& similarly $(d_{\Theta} : W^\Theta) := \lambda_{\Theta} \int \mu(dx) d_{X,\Theta}(x, \theta) = \frac{d\lambda_{\Theta}}{d\mu}$



Bayes's Thm (discrete version, adapted from Williams):

Let $\lambda : P(X \times \Theta)$ joint probability distribution.

Assume $\mu : D_X, V : D_\Theta$ s.t. $\lambda \ll \mu \otimes V$.

with $d_{X,H} = \frac{d\lambda}{d(\mu \otimes V)}$. $d_X = \frac{d\lambda}{d\mu}$ $d_\Theta = \frac{d\lambda_H}{dV}$

Let $d_{X|H}^{(-|\cdot)} : X \times \Theta \rightarrow W$

$$d_{X|H}^{(-|\cdot)}(x|\theta) := \begin{cases} d_H \theta \neq 0: & \\ & \\ \text{o.w.:} & \end{cases}$$

$$\frac{d_{X,H}(x,\theta)}{d_H \theta}$$


$$0$$

$$\lambda_{X|H=-} : \Theta \rightarrow P_X$$

$$\lambda_{X|H=\theta} := d_{X|H}^{(-|\theta)} \odot \mu$$

Bayes's formula:

$$P_r[-|H=-] = \lambda_{X|H=-}$$



Summary

$\mu \otimes \nu$ Product measures & Fubini-Tonelli;

μ_H Push-forward / law

$(D^X, \Sigma, (\cdot))$ module structure and affine linearity of ϕ

} Lebesgue integration

Standard vocabulary: joint dist., marginalisation, independence, invariance

density & Radon-Nikodym derivatives (heed the Warning)

almost certain properties

Conditional expectation & Probability

with Bayes's Thm.

Plan:

- 1) type-driven Probability: discrete case (Mon + Tue) ✓
- 2) Borel sets & measurable spaces (Wed) (Tue)
- 3) Quasi Borel spaces, Simple type structure (Wed)
- 4) Dependent type structure & standard Borel spaces (Thu)
- 5) Integration & random variables (Fri)

Please ask questions!

Smibble



Course
Web
Page