

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

TutorialFest
ACM-SIGPLAN 51st Symposium on
Principles of Programming Languages (POPL'24)
15 January, 2024
City of Westminster, England



THE UNIVERSITY OF EDINBURGH

informatics Ifcs

Laboratory for Foundations
of Computer Science

 BayesCentre

supported by:

 THE ROYAL
SOCIETY

The
Alan Turing
Institute

Facebook Research NCSC

Computational golden era of:

logic type rich
computation

Statistical
computation

Computational golden era of:

logic & type rich
computation

Expressive type systems:

Haskell, OCaml, Idris

Mechanised mathematics:

Agda, Coq, Isabelle/Hol, Lean

Verification:

SMT-powered, realistic
systems

Statistical
computation

generative modelling
+

efficient inference:

Monte-Carlo simulation
or gradient-based
optimisation

"AI"

Computational golden era of:

logic & type rich
computation

Statistical
computation

Clear connection to

Everything PoPL

- LAFI
- some PoPL
- this tutorial

Plan

1) Type A Setting for Probability & Statistics

2) 2 Implementations

Discrete Model



Full Model

familiar maths
to
introduce language

Same language

Why foundations?

Countably supported
measures

Discrete Probability

Continuous Probability
Lebesgue measure on
 \mathbb{R}^n

Takeaway:

use types to develop smoothly

Well-behaved probability
S-finite measures over
Standard Borel spaces

Measurable spaces
Standard, established
Poor type structure

Quasi-Borel spaces
new, experimental
Rich type structure
(this course)

Why types?

- spotlights meaningful operations

$$\int : (\text{Distribution } A) \times (A \rightarrow [0, \infty]) \rightarrow [0, \infty]$$

- documents intent

Probability Distribution A vs Density $A \rightarrow [0, \infty]$

- succinctness: easier to elaborate details
- esp. formal types: use theory without fully understanding it.

Plan:

- 1) Type-driven probability: discrete case
 - 2) Borel sets & measurable spaces
 - 3) Quasi Borel spaces
 - 4) Type structure & standard Borel spaces
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

sensible

Please ask questions!



← Course
web
page

Advertisement

Interested in the mathematical } foundations of Computer
Computational logical Science ?

Check out our PhD Programs:



THE UNIVERSITY of EDINBURGH
informatics

lfcs

Laboratory for Foundations
of Computer Science



LFCS
PhD
↗



Repetile
AI for
Robotics
CDT

Language of distribution & Probability

X type (=space) of values / outcomes

$\mathcal{D}X$ type of distributions / measures over X

$\mathcal{P}X \subseteq \mathcal{D}X$ subtype of probability measures (total measure)

$\mathcal{B}X$ type of measurable events - Subsets of X we

W type of weights : $[0, \infty]$ wish to measure

$\mu : \mathcal{D}X, E : \mathcal{B}X \vdash \underset{\mu}{\text{C}_e[E]} : W$

↳ measure μ assigns to E

→ type judgment

Axioms for measures

Empty event: $\emptyset : \mathcal{B}X$

Its measure is $0 : \mathbb{W}$:

$$\mu : \mathcal{D}X \vdash \text{Ce}[\emptyset] = 0 : \mathbb{W}$$

Axioms for measures

$\mathcal{B}X$ is a Boolean sub-algebra:

$$E : \mathcal{B}X \vdash E^c : \mathcal{B}X$$

$$E, F : \mathcal{B}X \vdash E \vee F, E \wedge F : \mathcal{B}X$$

$$E, C : \mathcal{B}X, \mu : \mathcal{D}X \vdash \quad (\text{disjoint additivity})$$

$$\underset{\mu}{\text{Ce}}[E] = \underset{\mu}{\text{Ce}}[E \cap C] + \underset{\mu}{\text{Ce}}[E \cap C^c] : \mathbb{W}$$

Axioms for measures

$\omega := (\mathbb{N}, \leq) \quad (\mathcal{B}, \subseteq) \quad (\mathbb{W}, \leq) \quad \text{posets}$

$$(\mathcal{B}X, \subseteq)^\omega := \left\{ (E_n)_{n \in \mathbb{N}} \in (\mathcal{B}X)^\mathbb{N} \mid E_0 \subseteq E_1 \subseteq E_2 \subseteq \dots \right\}$$

$(\mathcal{B}X, \subseteq)$ and (\mathbb{W}, \leq) are ω -chain-closed:

$$E_- : (\mathcal{B}X, \subseteq)^\omega \vdash \bigcup_n E_n : \mathcal{B}X \quad a_- : (\mathbb{W}, \leq)^\omega \vdash \sup_n a_n : \mathbb{W}$$

$$E_- : (\mathcal{B}X, \subseteq)^\omega, \mu : \mathcal{D}X \vdash \quad (\text{Scott Continuity})$$

$$\underset{\mu}{\text{Ce}} \left[\bigvee_n E_n \right] = \sup_n \underset{\mu}{\text{Ce}} [E_n] : \mathbb{W}$$

Axiom for Probability

Cast : $\text{px} \xleftarrow{\subseteq} \text{dx}$

$1 : \mathbb{W}$

$\mu : \text{px} \vdash \text{Ce}[x] = 1 : \mathbb{W}$

Cast μ

Avoid casting:

$E : \text{Bx}, \mu : \text{px} \vdash \Pr_{\mu}[E] := \text{Ce}[E] : [0,1] \subseteq \mathbb{W}$

Cast μ

Axioms for measures

Integration:

$$\mu: \mathcal{D}X, \varphi: \mathbb{W}^X \vdash \int \mu \varphi : \mathbb{W} \quad (\text{Lebesgue integral})$$

Again, avoid Costing:

$$\mu: \mathcal{P}X, \varphi: \mathbb{W}^X \vdash \underset{\mu}{\mathbb{E}}[\varphi] = \int (\text{Cost } \mu) \varphi : \mathbb{W} \quad (\text{Expectation})$$

More structure & notation later (...technical...)

Have: language + axioms

Want: model


Part 1 : discrete measures

Part 2: discrete + continuous

Discrete model

type X : set


$D_X :=$ histograms



Discrete model

type X : set

$D_X := \{ \mu : X \rightarrow \mathbb{W} \mid \mu \text{ is countably Supported} \}$
(next slide)



Support → Powerset

$\mu: W^X, S:\mathcal{P}X \vdash S \text{ supports } \mu :=$

$\forall x:X. \mu x > 0 \Rightarrow x \in S : \text{Prop}$

$\mu: W^X \vdash \text{Supp } \mu = \{x \in X \mid \mu x > 0\} : \mathcal{P}X$

$\text{Supp } \mu$ is the smallest set supporting μ

Discrete model

type X : set

$$DX := \{ \mu : X \rightarrow W \mid \mu \text{ is countably supported} \}$$


$$:= \{ \mu : X \rightarrow W \mid \text{Supp } \mu \text{ is countable} \}$$

Discrete model


type X : set

$D_X := \left\{ \mu: X \rightarrow W \mid \mu \text{ is countably Supported} \right\}$

$:= \left\{ \mu: X \rightarrow W \mid \text{Supp } \mu \text{ is countable} \right\}$




Ex. measures

1)  ... Counting measure $\#_X : DX$
 $a_0 \quad a_1 \quad a_2 \quad \dots (X \text{ ctbl})$ $\#_X := \lambda x : X. 1$

2) Dirac measure:

$$\sigma : X \vdash \delta_x : DX$$
$$:= \lambda x'. \begin{cases} x = x' : 1 \\ \text{o.w.} : 0 \end{cases}$$



Zero measure

$$0 := \lambda x. 0 : DX$$

Discrete model

type X : set

$$DX := \{ \mu : X \rightarrow \mathbb{W} \mid \mu \text{ is countably supported} \}$$

$$BX := \wp X$$

$$\begin{aligned} \mu : DX, E : BX &\vdash \text{C}_e[E] := \sum_{\mu} \mu \llcorner E \\ &:= \sum_{x \in E \cap \text{Supp } \mu} \mu \llcorner x \end{aligned}$$

Ex:

- Counting measure Counts Outcomes:

$$E: \mathcal{B}X \vdash C_e[E] = |E| := \begin{cases} E \text{ has } n \text{ elements: } n \\ E \text{ infinite: } \infty \end{cases}$$

- Dirac measure detects outcomes:

$$E: \mathcal{B}X, x: X \vdash C_e[E] = \delta_x = \begin{cases} x \in E: 1 \\ x \notin E: 0 \end{cases} =: [x \in E] : W$$

- Zero measure is zero:

$$E: \mathcal{B}X \vdash C_e[E] = 0$$

NB: $E: \mathcal{B}X \vdash [- \in E]: X \rightarrow W$
indicator function

Validate axioms

$$\mu : D X \vdash \underset{\mu}{C e[\emptyset]} = 0 : W$$

$$E, C : B X, \mu : D X \vdash$$

$$\underset{\mu}{C e[E]} = \underset{\mu}{C e[E \cap C]} + \underset{\mu}{C e[E \cap C^c]} : W$$

$$E_{_} : (B X, \leq)^{\omega}, \mu : D X \vdash$$

$$\underset{\mu}{C e[\bigvee_n E_n]} = \sup_n \underset{\mu}{C e[E_n]} : W$$

Kernels k from Γ to X :

$$k : (DX)^\Gamma$$

Kernels are "open/parameterised" measures

Ex: Dirac kernel: $\delta_- : (DX)^X$

Kock Integral

$$\mu : D\Gamma, u : DX \vdash \int^{\Gamma} \mu u : DX$$

In discrete model:

$$\int^{\Gamma} \mu u := \lambda x : X. \sum_{r \in \Gamma} \mu r \cdot \overbrace{k(r; x)}^{:= k(r)(x)}$$

(Weak) disintegration problem:

Input: $\mu: D\Gamma$ $V: DX$

Output: a kernel $k:(DX)^{\Gamma}$ s.t.

$$\int \mu k = V$$

Call such $k \stackrel{a}{=} (\text{weak}) \text{ disintegration of } V$

w.r.t. μ .

(Non-standard
terminology)

Ex disintegration:

$$\underline{n} := \{0, 1, 2, \dots, n-1\}$$

disintegrate $\#_{\geq \underline{n+1}}$ w.r.t. $\#_2$

Define:

$$k: \left(D(\geq \underline{n+1})\right)^2$$

$$k(x_j; f) := \begin{cases} f(n) = x: & 1 \\ \text{o.w.} & 0 \end{cases}$$

check:

$$\left(\sum_{x \in \underline{\mathbb{Z}}} k\right) f = \sum_{x \in \underline{\mathbb{Z}}} \#_2^x \cdot k(x; f)$$

$$= k(0; f) + k(1; f) = \#_2^{\underline{n+1}}(f) = 1$$

Probability measures


$$P_X := \left\{ \mu : D_X \mid \underset{\mu}{\text{Ce}}[X] = 1 \right\} \overset{\subseteq}{\hookrightarrow} D_X$$

Probability measures

$$P_X := \left\{ \mu : D_X \mid \underset{\mu}{\text{C}_e[X]} = 1 \right\} \hookrightarrow^{\subseteq} D_X$$

Lemma: $\delta_x : x \rightarrow D_X$ and $\mathfrak{f} : D^\Gamma \times (D_X)^\Gamma \rightarrow D_X$

lift along the inclusion cast: $P \hookrightarrow^{\subseteq} D$:



Prop (discrete Giry):

(Michèle Giry '82)

(D, δ_-, \oint) is a monad i.e.

$$m: \Gamma, n:(Dx)^\Gamma \vdash \oint \delta_n = k \ r$$

$$\mu: D\Gamma \vdash \oint \mu(\lambda x) \delta_x = \mu$$

$$\mu: D\Gamma, \kappa: (Dx)^\Gamma, t: (DY)^X \vdash$$

$$\oint \mu(\lambda x) \left(\oint (\kappa r) t \right) = \oint \left(\oint \mu \kappa (\lambda x) t(x) \right)$$

Corollary: (P, δ_-, \oint) is a monad.

Language of distribution & Probability

X type (=space) of values / outcomes

$\mathcal{D}X$ type of distributions / measures over X

$\mathcal{P}X \subseteq \mathcal{D}X$ subtype of probability measures (total measure)

$\mathcal{B}X$ type of measurable events - Subsets of X we

W type of weights : $[0, \infty]$ wish to measure

$\mu : \mathcal{D}X, E : \mathcal{B}X \vdash \underset{\mu}{\text{C}_e[E]} : W$

↳ measure μ assigns to E

→ type judgment

Plan:

- 1) Type-driven probability: discrete case
 - 2) Borel sets & measurable spaces
 - 3) Quasi Borel spaces
 - 4) Type structure & standard Borel spaces
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

please ask questions!
sensible



← Course
web
page

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

TutorialFest
ACM-SIGPLAN 51st Symposium on
Principles of Programming Languages (POPL'24)
15 January, 2024
City of Westminster, England



THE UNIVERSITY OF EDINBURGH

informatics Ifcs

Laboratory for Foundations
of Computer Science

 BayesCentre

supported by:



THE ROYAL
SOCIETY

The
Alan Turing
Institute

Facebook Research NCSC

Discrete model

Recap

$$\text{type: set} \quad W := [0, \infty] \quad \mathcal{B}X := \mathcal{P}X$$


$$DX := \{\mu: X \rightarrow W \mid \text{Supp } \mu \text{ countable}\}$$

$$PX := \{\mu \in DX \mid \underset{\mu}{\text{Ce}}[X] = 1\}$$

$$\underset{\mu}{\text{Ce}}[E] := \sum_{x \in E} \mu_x \quad S_x := \lambda x'. \begin{cases} x = x': 0 \\ x \neq x': 1 \end{cases}$$


$$\oint \mu k := \lambda x. \sum_{m \in \Gamma} \mu^m \cdot k(m; x)$$

Ex. measures

1)  ... Counting measure $\#_X : DX$
 $a_0 \quad a_1 \quad a_2 \quad \dots (X \text{ ctbl})$ $\#_X := \lambda x : X. 1$

2) Dirac measure:

$$\sigma : X \vdash \delta_x : DX$$
$$:= \lambda x'. \begin{cases} x = x' : 1 \\ \text{o.w.} : 0 \end{cases}$$




Zero measure

$$0 := \lambda x. 0 : DX$$


Ex distributions

Recap

Counting measure (λ_{ctg}): $\#_X := \lambda x. 1$



Dirac measure δ_x (prev slide)



Zero measure $\underline{\lambda} := \lambda x. 0$



Product measures

$$\mu: D^X, \nu: D^Y \vdash \mu \otimes \nu := \int \mu(dx) \int \nu(dy) \delta_{(x,y)}: D^{(X \times Y)}$$

(\otimes) lifts along $P \hookrightarrow D$)

$$= \lambda(x, y). \mu x \cdot \nu y$$

\uparrow
discrete
model

$$E_{\#}: \#_{X \times Y} = \#_X \otimes \#_Y$$

build measures
compositionally

Indeed:

$$(\# \otimes \#)(x, y) = \#x \cdot \#y = 1 \cdot 1 = 1 = \#(x, y)$$

Notation: $\lambda : D(X \times Y), k : (DZ)^{X \times Y} \vdash \oint \lambda(z_2, dy) k(z, y)$
 $\quad\quad\quad := \oint \lambda k$

Fubini - Tonelli Thm:

Integrate in any order:

$\mu : DX, v : DY, k : (DZ)^{X \times Y} \vdash$

$$\begin{aligned} \oint \mu(dx) \oint v(dy) k(x, y) &= \iint (\mu \otimes v)(x_2, dy) \\ &= \oint v(dy) \oint \mu(dx) k(x, y) \end{aligned}$$

Pushing a measure forward

$$\mu: D_{\Omega}, \alpha: X^n \vdash \mu_f := \int \mu(d\omega) \delta_{\alpha\omega} : DX$$

$$= \lambda x. \sum_{\omega \in \Omega} \mu \omega$$

$$\alpha\omega = x$$

$\alpha: X^{\Omega}$: random element

$\mu_\alpha: DX$: the law of α (w.r.t. μ)

Ex: represent Configurations of 2 dice using

$$\text{Die}_6 := \{1, 2, \dots, 6\} \quad \text{Die}_6^2$$

Letting $(+): \text{Die}_6^2 \rightarrow \mathbb{N}^2 \xrightarrow{(+)} \mathbb{N}$

We have that the law of $(+)$:

$$\mu = (\# \underset{\text{Die}}{\otimes} \# \underset{\text{Die}}{\circ})_{(+)} : \mathbb{D}/\mathbb{N}$$

build measures
compositionally

$\mu_S =$ Number of outcomes whose sum is S

Scaling a measure

$$(\cdot) : \mathbb{W} \times D_X \longrightarrow D_X$$

$$a \cdot \mu := \lambda x. a \cdot \mu x$$

$(\cdot) : \mathbb{W} \times D_X \rightarrow D_X$ is an action of monoid $(\mathbb{W}, \cdot, 1)$ on D_X :

$$\mu : D_X \vdash$$

$$1 \cdot \mu = \mu$$

$$a, b : \mathbb{W}, \mu : D_X \vdash$$

$$a \cdot (b \cdot \mu) = (a \cdot b) \cdot \mu$$

Normalisation

unit type \rightarrow

$$1 := \left\{ \begin{array}{l} \text{✓} \\ \text{✗} \end{array} \right\}$$

unit value

$\mu : D_X, Ce[X] \neq 0, \infty +$

$$\|\mu\| := \left(\frac{1}{Ce[X]} \right) \cdot \mu \quad : P_X$$

Ex:

$$\emptyset \neq A \subseteq_{fin} X \quad : U_{A \subseteq X} := \left\| (\#_A) \right\|_{A \subseteq X} \quad : P_X$$

$$1 \xrightarrow{\#_A} DA \xrightarrow{(-)_{A \subseteq X}} DX \xrightarrow{\|\cdot\|} P_X$$

I.e.

$$U_{A \subseteq X} := \lambda n. \begin{cases} n \in A: & \frac{1}{|A|} \\ n \notin A: & 0 \end{cases} \quad \text{so} \quad U_{\{n \in X\}} = \delta_n$$

Standard vocabulary

Joint distributions: $\mu: D(X_1 \times X_2)$

Marginal distribution: $X_1 \xleftarrow{\pi_1} X_1 \times X_2 \xrightarrow{\pi_2} X_2$

law of projection

$\mu_{\pi_i}: D X_i$

Marginalisation: $\mu_{\pi_i} = \iint \mu(dx, dy) S_x$
integrate out y

Exercise: $\mu: P X, V: D X \vdash (\mu \otimes V)_{\pi_2} = V$

independence

Pairing r.e.s:


$$\alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash$$

$$\langle \alpha, \beta \rangle := \lambda w. \langle \alpha w, \beta w \rangle : (X + Y)^{\Omega}$$

$$\lambda : D\Omega, \alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash \alpha \perp_{\lambda} \beta := \lambda_{\langle \alpha, \beta \rangle} = \lambda_{\alpha} \oplus \lambda_{\beta}$$

: prop

α, β independent w.r.t. λ



Ex ^(Durrett) represent outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : \bigcup_{C^3} \otimes \bigcup_{C^2} \otimes \bigcup_{C^1} : P_\Omega$$

$$\pi_i : \Omega \rightarrow C \quad \text{outcome of } i^{\text{th}} \text{ toss}$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{(\stackrel{?}{=})} B$$

where : $\stackrel{?}{=} : C^2 \rightarrow B := \{ \text{True}, \text{False} \}$

$$\stackrel{?}{=} := \begin{cases} x = y : \text{True} \\ x \neq y : \text{False} \end{cases}$$

E₂^(Durrett) represent outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : U_C \otimes U_C \otimes U_C : P_{\Omega}$$

$\pi_i : \Omega \rightarrow C$ outcome of i^{th} toss

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{?} B$$

marginalisation

$$\lambda_{\text{Same}_{12}}^T = (U_C \otimes U_C)^T \stackrel{?}{=} \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

\uparrow

$$\begin{matrix} U_C(T) \cdot U_C(H) \\ + \\ U_C(H) \cdot U_C(H) \end{matrix}$$

$$\text{so } \lambda_{\text{Same}_{12}}^F = \frac{1}{2} \text{ too}$$

E_r^(Durrett) represent outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega = C \times C \times C \quad \lambda : \cup_C \otimes \cup_C \otimes \cup_C : P_{\Omega}$$

$$\pi_i : \Omega \rightarrow C \quad \text{outcome of } i^{\text{th}} \text{ toss}$$

$$\underline{i \neq j} : \lambda_{\text{Same}_{ij}} = V_B$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{?} B$$

$$\lambda : \langle \text{Same}_{12}, \text{Same}_{23} \rangle \ni (T, T) \mapsto \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$$

$\hookrightarrow \lambda(H, H, H) \quad \hookrightarrow \lambda(T, T, T)$

$$(T, F) \mapsto \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$$

$\hookrightarrow \lambda(H, H, T) \quad \hookrightarrow \lambda(T, T, H)$

E₂^(Durrett) represent outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega := C \times C \times C \quad \lambda : U_C \otimes U_C \otimes U_C : P_{\Omega}$$

$$\pi_i : \Omega \rightarrow C \quad \text{outcome of } i^{\text{th}} \text{ toss}$$

$$\underline{i \neq j} : \lambda_{\text{Same}_{ij}} = U_B$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{=} B$$

$$\lambda_{\langle \text{Same}_{12}, \text{Same}_{23} \rangle} = U_{B \times B} = U_B \otimes U_B = \lambda_{\text{Same}_{12}} \otimes \lambda_{\text{Same}_{13}}$$

$$\text{So } \text{Same}_{12} \perp \lambda \text{ Same}_{13}$$

independence


Pairing r.e.s:

$$\alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash$$

$$\langle \alpha, \beta \rangle := \lambda w. \langle \alpha w, \beta w \rangle : (X + Y)$$

$$\lambda : D\Omega, \alpha : X^{\Omega}, \beta : Y^{\Omega} \vdash \alpha \perp \beta := \lambda_{\langle \alpha, \beta \rangle} = \lambda_{\alpha} \otimes \lambda_{\beta}$$

: prop
α, β independent w.r.t. λ



I-any version:

$$\lambda : D\Omega, \alpha_i : \prod_{i \in I} X_i^{\Omega} \vdash \perp_{\lambda}^{\alpha_i} :=$$

α_i-independent
w.r.t. λ

$$\forall J \subseteq_{fin} I. \quad \lambda_{\langle \alpha_j \rangle_{j \in J}} = \bigotimes_{j \in J} \lambda_{\alpha_j} : \text{Prop}$$

E₂^(Durrett) represent outcomes of 3 coin tosses:

$$C := \{T, H\} \quad \Omega = C \times C \times C \quad \lambda : \bigcup_C \otimes \bigcup_C \otimes \bigcup_C : P_{\Omega}$$

$$\pi_i : \Omega \rightarrow C \quad \text{outcome of } i^{\text{th}} \text{ toss}$$

$$\underline{i \neq j} : \lambda_{\text{Same}_{ij}} = \nu_B$$

$$\underline{i \neq j} : \text{Same}_{ij} \perp \text{Same}_{jk}$$

$$\text{Same}_{ij} : \Omega \xrightarrow{\langle \pi_i, \pi_j \rangle} C \times C \xrightarrow{(\cdot)} B$$

$$\frac{1}{\lambda} \left\{ \text{Same}_{12}, \text{Same}_{23}, \text{Same}_{13} \right\}$$

Intuition: $\text{Same}_{13} = \text{IFF} (\text{Same}_{12}, \text{Same}_{23})$

Calc:

$$\lambda_{\langle \text{Same}_{12}, \text{Same}_{23}, \text{Same}_{13} \rangle} = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} \neq \frac{1}{2^3} = \lambda_{\text{Same}_{12}} \otimes \lambda_{\text{Same}_{23}} \otimes \lambda_{\text{Same}_{13}}$$
$$\hookrightarrow \lambda(H, H, H) \quad \hookrightarrow \lambda(T, T, T)$$

Vocabulary

(Discrete) Measure space $(X, \mu : D X)$

Measure preserving $f : (X, \mu) \rightarrow (Y, \nu)$

function $f : X \rightarrow Y$ s.t. $\mu_f = \nu$

$\mu : D X$, $f : X \rightarrow Y \vdash \mu$ invariant under $f :=$

$f : (X, \mu) \rightarrow (Y, \nu)$

E.g.:

$\mu : D X$, $\nu : D Y \vdash$


Swap : $(X \times Y, \mu \otimes \nu) \longrightarrow (Y \times X, \nu \otimes \mu)$ so

$\mu : D X \vdash \mu \otimes \mu$ invariant under Swap


Weights as measures

NB: $\overset{\text{unit type}}{\sim}$ $1 := \{ \langle \rangle \}^{\text{unit value}}$

Observation:



Proof:



□

Integration

$$\mu: D_X, \varphi: W^X \vdash \int^\mu \varphi : W$$
$$:= \sum_{x \in X} \mu_x \cdot \varphi_x$$

(Lebesgue
integral)

Can derive it:

$$D_X \times W^X \xrightarrow{D_X \times (\cong o -)} D_X \times (D_1)^X$$
$$\int \downarrow := \xleftarrow{\cong} D_1 \downarrow f$$

Additivity:

$$\text{I ctsl}, \mu_i : (DX)^I \vdash \sum_{i \in I} \mu_i : DX$$
$$:= \lambda x. \sum_{i \in I} \mu_i x$$

NB:

$$\text{supp} \sum_i \mu_i \subseteq$$
$$\bigcup_i \text{supp } \mu_i$$

✓ ctsl

Ex: Bernoulli distribution

$$p : [0,1] \vdash B(p) := p \cdot \delta_{\text{True}} + (1-p) \cdot \delta_{\text{False}} : P/B$$

i.e. $B_p : \begin{cases} \text{True} \mapsto p \\ \text{False} \mapsto 1-p \end{cases}$

Thm (affine-linearity):

\int is affine-linear in each argument:

I ctbl

$$\mu: (\Omega^P)^I, k: (\Omega^X)^P \vdash \int (\sum_{i \in I} a_i \cdot \mu_i) k = \sum_{i \in I} a_i \cdot \int \mu_i k$$

I ctbl, $\mu: \Omega^P$, $a_i: W^I$, $k_i: \Omega^X^I$

$$\int \mu(dx) \left(\sum_{i \in I} a_i \cdot k_i(x) \right) = \sum_{i \in I} a_i \cdot \int \mu k_i$$

Prop: $\mathbb{W} \cong D\mathbf{1}$ is a \mathbb{G} -semi-ring isomorphism:

$$(\mathbb{W}, \Sigma, (\cdot), 1) \cong (D\mathbf{1}, \Sigma, (\cdot), \delta_{\leq})$$

and $(\cdot) : \mathbb{W} \times D\mathbf{X} \rightarrow D\mathbf{X}$ makes $D\mathbf{X}$ into a module:

$$\left(\sum_{i \in I} a_i \right) \cdot \mu = \sum_{i \in I} (a_i \cdot \mu) \quad a \cdot \sum_{i \in I} \mu_i = \sum_{i \in I} a \cdot \mu_i$$

Corollary: \int is affine-linear in each argument.

Random variable :

NB: $\bar{\mathbb{R}} := [-\infty, \infty]$

A random element $\alpha: \bar{\mathbb{R}}^{\Omega} \rightarrow \mathbb{W}^{\Omega}$ (wrt some $\mu: D\Omega \rightarrow \mathbb{R}$)

Can add, multiply r.v.'s.

To integrate r.v.'s:

$$(-)^+: \bar{\mathbb{R}}^{\Omega} \longrightarrow \mathbb{W}^{\Omega}$$

$$\alpha^+ := \lambda w \cdot \begin{cases} \alpha \cdot w \geq 0 : \alpha \cdot w \\ 0 \cdot w : 0 \end{cases} = [\alpha \geq 0] \cdot |\alpha|$$

$$\alpha^- := \lambda w \cdot \begin{cases} \alpha \cdot w \leq 0 : |\alpha \cdot w| \\ 0 \cdot w : 0 \end{cases} = [\alpha \leq 0] \cdot |\alpha|$$

$$\text{So } \alpha = \alpha^+ - \alpha^-$$

$\mu: D\Omega, \alpha: \overline{\mathbb{R}}^n, \int \mu \alpha^+ < \infty \text{ or } \int \mu \alpha^- < \infty \vdash$

$$\int \mu \alpha := \int \mu \alpha^+ - \int \mu \alpha^- : \overline{\mathbb{R}}$$

Ex. The (discrete) Lebesgue p -space:

$$p: \Sigma, \infty), \mu: P\Omega \vdash L_p(\Omega, \mu) :=$$

$$\left\{ \alpha: \overline{\mathbb{R}}^n \mid \underset{\mu}{\mathbb{E}} [|\alpha|^p] < \infty \right\}$$

$L_p(\Omega, \mu)$ has a norm $\|\alpha\| := \sqrt[p]{\underset{\mu}{\mathbb{E}} [|\alpha|^p]}$ almost Banach

$L_2(\Omega, \mu)$ has an inner product $\langle \alpha, \beta \rangle := \underset{\mu}{\mathbb{E}} [\alpha \cdot \beta]$ almost Hilbert


Density

a density over X : $d : X \rightarrow W$

$$d : W^X, \mu : D X \vdash d \odot \mu : D X \\ := \oint \mu(dx) (d_n \cdot S_n)$$

Warning The types of measures & densities in the discrete model are close, but still different. They coincide on countable sets, so people often confuse them. Types help us keep them separate.

Intuition:



Almost certain properties

$E: BX, \mu: DX \vdash \mu(\lambda x) \text{-almost certainly } x \in E : \text{Prop}$

$$:= [-\epsilon E] \odot \mu = \mu$$

$$\vdash_{\text{NB}} [\lambda x \in E] = \begin{cases} x \in E & 1 \\ x \notin E & 0 \end{cases} : \mathbb{W}$$

When $\mu: Px$ we say instead

$\mu(\lambda x) \text{-almost surely } x \in E$

Absolute continuity

d is a density of μ w.r.t. V or
 d is a Radon-Nikodym derivative w.r.t. V

$$\mu, V : D^X, d : W^X \vdash d = \frac{d\mu}{dV} \quad : \text{Prop}$$

$$:= \mu = d \odot V$$

$\mu, V : D^X \vdash \mu \ll V := \mu$ is absolutely continuous w.r.t. V : Prop

$$:= \exists d : W^X. \quad d = \frac{d\mu}{dV}.$$

$=:$ μ has a density w.r.t. V

Lemma: $\mu, V : D^X,$
 $\mu \ll V,$
 $k : (D^Y)^X$

$$\oint V(dx) \frac{d\mu}{dV}(x) \cdot k_{x^*} = \oint \mu(dx) k_{x^*}$$

$$\underline{Ex}: \bigcup_{A \subseteq X} \ll (\#_A)_{\text{Cost}: A \subseteq X}$$

$$\frac{dV_{A \subseteq X}}{d(\#)_A^{\text{Cost}}} = \lambda_x \cdot \begin{cases} x \in A: & \frac{1}{|A|} \\ \text{O.W.:} & 0 \end{cases}$$

but also:

$$\frac{dV_{A \subseteq X}}{d(\#)_A^{\text{Cost}}} = \lambda_x \cdot \frac{1}{|A|}$$

Radon-Nikodym Thm: (discrete version)

$\mu, \nu: P_X \vdash \mu \ll \nu$ iff $\forall x. V_x = 0 \Rightarrow \mu x = 0$
i.e. $\text{Supp } \mu \subseteq \text{Supp } \nu$

In that case, if $d_1, d_2 = \frac{d\mu}{d\nu}$ then

$V(dx)$ -a.s. $d_1 x = d_2 x$

Ex: for ctbl x , $\forall \mu: \Omega^x. \mu \ll \#_x$. Proof: vacuously, as $\#_x \neq 0$.

Then $\lambda x. \mu x = \frac{d\mu}{d\#}.$


Conditional expectation

β is a conditional expectation of α w.r.t. μ along H

$$\mu: \Omega \rightarrow H: X^{\Omega}, d: \mathcal{F}_1(\Omega, \mu), \beta: \mathcal{F}_1(X, \mu_H)$$

$$+ \beta = \mathbb{E}[\alpha | H = -] \quad : \text{Prop}$$

$$:= \forall \varphi: \mathcal{F}_1(Y, \mu_H^M). \int \mu_H(dx) \beta(x) \cdot \varphi(x) = \int \mu(dw) \alpha(w) \cdot \varphi(Hw)$$



Thm (Kolmogorov): (discrete version)

There is a function

$$\mathbb{E}[\cdot | \cdot] \in \prod_{\mu: P_{\Omega}} \prod_{H: X^{\Omega}} \mathcal{L}_1(\Omega, \mu) \rightarrow \mathcal{L}_1(X, \mu_H)$$

s.t. $\mathbb{E}[\alpha | H = \cdot]$ is a conditional expectation of α w.r.t. μ
along H .

Conditional Probability (discrete version):


$$\begin{aligned} H: X^{\Omega}, \mu: P_X + \Pr_{\mu}[- \mid H = -] : (P_{\Omega})^X \\ := \lambda x_0: X. \lambda w_0: \Omega. \underset{w \sim \mu}{\mathbb{E}} [\mathbb{E}_{\{w_0 = w\}} \mid H_{w_0} = x_0] \end{aligned}$$

Bayes's Thm (discrete version, adapted from Williams):

Let $\lambda : P(X \times \Theta)$ joint probability distribution.

Assume $\mu : D_X$, $\nu : D_{\Theta}$ s.t. $\lambda \ll \mu \otimes \nu$.

with $d_{X,H} = \frac{d\lambda}{d(\mu \otimes \nu)}$.



$$\text{Obs 1: } d_X : W^X \quad \text{then } d_X = \frac{d\lambda}{d\mu}$$

$$d_X := \lambda \pi_1 \int \nu(d\theta) d_{X,H}(x_\theta, \theta)$$

$$\text{And similarly } (d_{H,H} : W^\Theta) := \lambda \theta \int \mu(dx) d_{X,H}(x_\theta, \theta) = \frac{d\lambda_H}{d\nu}$$

Bayes's Thm (discrete version, adapted from Williams):

Let $\lambda : P(X \times \Theta)$ joint probability distribution.

Assume $\mu : D_X, \nu : D_{\Theta}$ s.t. $\lambda \ll \mu \otimes \nu$.

with $d_{X,H} = \frac{d\lambda}{d(\mu \otimes \nu)} \cdot d_X = \frac{d\lambda_\alpha}{d\mu} \quad d_{\Theta} = \frac{d\lambda_H}{d\nu}$

Let $d_{X|H}^{(-|\cdot)} : X \times \Theta \rightarrow W$


$$d_{X|H}^{(x|\theta)} := \begin{cases} d_H \neq 0: & \frac{d_{X,H}(x,\theta)}{d_H \theta} \\ \text{O.w.:} & 0 \end{cases}$$

$$\lambda_{X|H=-} : \Theta \rightarrow P_X$$

$$\lambda_{X|H=\theta} := d_{X|H}^{(-|\theta)} \otimes \mu$$

Bayes's formula:

$$P_r[-|H=-] = \lambda_{X|H=-}$$



Summary

$\mu \otimes \nu$ Product measures & Fubini-Tonelli

μ_H Push-forward / law

$(D^x, \Sigma, (\cdot))$ module structure and affine linearity of ϕ

Lebesgue integration

Standard vocabulary: joint dist., marginalisation, independence, invariance

density & Radon-Nikodym derivatives (heed the **Warning**)

almost certain properties

Conditional expectation & probability

with Bayes's Thm.

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces
 - 3) Quasi Borel spaces
 - 4) Type structure & standard Borel spaces
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

sensible

Please ask questions!



← Course
web
page

discrete model measure Only histograms:



Want :

- lengths
- areas
- volumes .

Continuous Caveat:

Then: No $\lambda: \mathcal{P}R \rightarrow [0, \infty]$:

$$\lambda(a, b) = b - a$$

(generalises length)

$$\lambda(r + A) = \lambda A$$

(translation invariant)

$$\lambda\left(\bigcup_{n=0}^{\infty} A_n\right) = \sum_{n=0}^{\infty} \lambda A_n$$

σ -additive

Takeaway: Taking $\mathcal{B}/R := \mathcal{P}R$

Excludes measures such as:

length, area, volume

Workaround: only measure well-behaved subsets

Df: The Borel Subsets $B_{\mathbb{R}} \subseteq \mathcal{P}(\mathbb{R})$:

- Open intervals $(a, b) \in B_{\mathbb{R}}$

Closure under σ -algebra operations:

$$\frac{\emptyset \in B_{\mathbb{R}}}{A \in B_{\mathbb{R}}} \qquad \frac{A \in B_{\mathbb{R}}}{A^c := \mathbb{R} \setminus A \in B} \qquad \frac{\vec{A} \in B_{\mathbb{R}}^N}{\bigcup_{n=0}^{\infty} A_n \in B_{\mathbb{R}}}$$

Empty set { complements countable unions

Examples

discrete Countable: $\{r\} = \bigcap_{\varepsilon \in \mathbb{Q}^+} (r-\varepsilon, r+\varepsilon) \in \mathcal{B}_{\mathbb{R}}$

I countable $\Rightarrow I = \bigcup_{r \in I} \{r\} \in \mathcal{B}_{\mathbb{R}}$

Closed intervals: $[a,b] = (a,b) \cup \{a,b\}$

Non-examples?

More complicated: analytic, lebesgue

Df: Measurable Space $V = (V, \mathcal{B}_V)$

Set \hookrightarrow
(Carrier)
Family of
Subsets
 $\mathcal{B}_V \subseteq P(V)$

Closed under σ -algebra operations:

$$\underline{\emptyset \in \mathcal{B}_V}$$

Empty set

$$\underline{A \in \mathcal{B}_V}$$

$A^c := V \setminus A \in \mathcal{B}_V$

Complements

$$\overline{\bigcup_{n=0}^{\infty} A_n \in \mathcal{B}_V}$$

Countable unions

Idea: structure all spaces after the worst-case scenario

Examples

- Discrete spaces $\overset{\text{meas}}{X} = (X, \mathcal{P}X)$
 - Euclidean spaces \mathbb{R}^n → replace intervals with
charts $\prod_{i=1}^n (a_i, b_i)$
 $\mathbb{R}^{\mathbb{N}}$ similarly
 - Sub spaces: $A \in \mathcal{P}_V$ $A := (A, [\mathcal{B}_V] \cap A)$
 - Products: $A \times B := (\mathcal{L}A \times \mathcal{L}B, \sigma([\mathcal{B}_A] \times [\mathcal{B}_B]))$
- $\{C \cap A \mid C \in \mathcal{B}_V\}$

Def: Borel measurable functions $f: V_1 \rightarrow V_2$

- functions $f: V_1 \rightarrow V_2$
- inverse image preserves measurability:

$$f^{-1}[A] \in \mathcal{B}_{V_1} \iff A \in \mathcal{B}_{V_2}$$

Examples

- $(+), (\cdot): \mathbb{R}^2 \rightarrow \mathbb{R}$
- $| - |, \sin: \mathbb{R} \rightarrow \mathbb{R}$
- any continuous function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$
- any function $f: X^n \rightarrow Y$

Category Meas

Objects: Measurable spaces

Morphisms: Measurable functions

Identities:

$$id : V \rightarrow V$$

Composition:

$$\begin{array}{ccc} f : V_2 \rightarrow V_3 & & g : V_1 \rightarrow V_2 \\ \curvearrowright & & \\ f \circ g : V_1 \rightarrow V_3 & & \end{array}$$

Meas Category

Products, Co-products / disjoint union, Subspaces
Categorical limits, colimits, but:

Theorem [Arrow 61] No σ -algebras B_{B_R} , B_{R^R} for measurable

membership predicate \leftarrow (\exists) : $(B_R, B_{B_R}) \times R \rightarrow \text{Bool}$
 $(U, r) \mapsto [r \in U]$

eval : $(\text{Meas}(R, R), B_{R^R}) \times R \rightarrow R$
 $(f, r) \mapsto f(r)$

Sequential Higher-order structure:

I Countable : $V^{\mathbb{I}} = \prod_{i \in \mathbb{I}} V$

\Rightarrow Some higher-order structure in Meas:

Cauchy $\in \mathcal{B}_{[-\infty, \infty]^N}$

$$\text{Cauchy} := \bigcap_{\varepsilon \in \mathbb{Q}^+} \bigcup_{k \in \mathbb{N}} \bigcap_{\substack{m, n \in \mathbb{N} \\ m, n > k}} \{ \vec{y} \in [-\infty, \infty]^N \mid |y_m - y_n| < \varepsilon \}$$

$$\lim \text{Sup} : [-\infty, \infty]^N \rightarrow [-\infty, \infty] \quad \lim : \text{Cauchy} \rightarrow \mathbb{R}$$

Compose higher-order building blocks:

lim is measurable!

$$\text{VanishingSeq}(\mathbb{R}) := \left\{ \vec{r} \in \mathbb{R}^N \mid \lim_{n \rightarrow \infty} r_n = 0 \right\} \subset \mathbb{B}_{\mathbb{R}^N}$$

$$\text{approx_}: \text{VanishingSeq}(\mathbb{R}^+) \times \mathbb{R} \rightarrow \mathbb{Q}^N$$

$$\text{s.t.: } |(\text{approx}_{\Delta} \vec{r})_n - r| < \Delta_n$$

Slogan: Measurable by Type !

Not all operations of interest fit:

$$\limsup : ([-\infty, \infty]^{\mathbb{R}})^{\mathbb{N}} \rightarrow [-\infty, \infty]^{\mathbb{R}}$$

$$\limsup := \lambda \vec{f}. \lambda x. \limsup_{n \rightarrow \infty} f_n x$$

Intrinsically
higher-order !

Want

Slogan: measurability by type!

But

For higher-order building blocks

defer measurability proofs until

we resume 1st order fragment \Rightarrow ^{non}composition

Plan

Def: $V \in \text{Meas}$ is Standard Borel when

$$V \cong A \quad \text{for some } A \in \mathcal{B}_R$$

the "good part" of Meas - the subcategory

$$\text{Sbs} \hookrightarrow \text{Meas}$$

Sbs including

- Discrete ' \mathbb{I} ', \mathbb{I} countable
- Countable products of Sbs:

$$\mathbb{R}^n, \mathbb{R}^{\mathbb{N}}, \mathbb{Z}^n, \mathbb{N}^{\mathbb{N}}$$

- Borel subspaces of Sbs:

$$\mathbb{I} := [0, 1]$$

$$\mathbb{R}^+ := (0, \infty) \quad \mathbb{R}_{\geq 0} := [0, \infty]$$

- Countable coproducts of Sbs:

$$\mathbb{W} := [0, \infty]$$


$$\overline{\mathbb{R}} := [-\infty, \infty]$$

Conservative extensions:

Concrete spaces

→ we "observe"

Standard Borel spaces



Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces
 - 4) Type structure & standard Borel spaces
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

Please ask questions!
sensible



← Course
web
page

Core idea

Measure Theory


Abs Theory

Sample space Ω




Primitive notions:

measurable subset



random element

$\downarrow \alpha$



Derived

notions:

measure

random

elements

$\alpha: \Omega \rightarrow \text{Space}$

Events

$E \in \mathcal{B}_X$


Def: Quasi-Borel space $X = (X_1, \mathcal{R}_X)$

$$\mathcal{R}_X \subseteq {}^{\text{L}(X_1)}\mathbb{R} \quad \text{closed under:}$$

Set ↗
"carrier"
Set of
functions $\alpha: \mathbb{R} \rightarrow X_1$
"random elements"

- Constants:

$$\frac{x \in X_1}{(\lambda r, x) \in \mathcal{R}_X}$$



- precomposition:

- recombination

Def: Quasi-Borel space $X = (X_s, \mathcal{R}_X)$



$$\mathcal{R}_X \subseteq {}^L X_s \quad \text{Closed under:}$$

- precomposition:

$$\alpha \in \mathcal{R}_X \quad \varphi: \mathbb{R} \rightarrow \mathbb{R} \text{ in } \mathcal{S}_{\mathbb{R}}$$

$$(\varphi \circ \alpha): \mathbb{R} \xrightarrow{\varphi} \mathbb{R} \xrightarrow{\alpha} X_s \in \mathcal{R}_X$$

Set \curvearrowleft "carrier"
Set \curvearrowright "functions $\alpha: \mathbb{R} \rightarrow X_s$ "
"random elements"



Def: Quasi-Banach space $X = (X, \mathcal{R}_X)$

$$\mathcal{R}_X \subseteq L^{\mathcal{R}_X}$$



Closed under:

- recombination

$$\vec{\alpha} \in R_X^N \quad \mathcal{R} = \bigcup_{n=0}^{\infty} A_n$$

$$\lambda r. \left\{ \begin{array}{l} r \in A_n : \alpha_n r \\ \vdots \\ r \in A_0 : \alpha_0 r \end{array} \right.$$



Set ↗
"carrier"
Set of
functions $\alpha: \mathbb{R} \rightarrow X$
"random elements"




Ref: Quasi-Banach space $X = (LX, \mathcal{R}_X)$

$$\mathcal{R}_X \subseteq LX^J \quad \text{Closed under:}$$

Set ↗
"carrier"
Set of
functions $\alpha: \mathbb{R} \rightarrow X$
"random elements"

- Constants: 
- Precomposition: 

- recombination



Examples

recombination of
constants

$$- \mathbb{R} = (\mathbb{R}, \text{Meas}(\mathbb{R}, \mathbb{R}))$$

abs underlying \mathbb{R}

$$- X \in \text{Set}, \quad \lceil X \rceil := (X, \sigma\text{-simple}(\mathbb{R}, X))$$

discrete qbs on X

$$- " \quad \lceil X \rceil_{\text{Qbs}} := (X, X^{\mathbb{L}(\mathbb{R})})$$

all functions

Indiscrete qbs on X

Qbs morphism $f: X \rightarrow Y$

- function $f: X_i \rightarrow Y_j$

- $\alpha \stackrel{R}{\downarrow} \in R_X$

$\begin{array}{c} R \\ \alpha \downarrow \\ x_i \\ f \downarrow \\ y_j \end{array} \in R_Y$

Example

- Constant functions

one qbs
morphism

- σ -single functions

are qbs morphisms

Category Qbs \Leftarrow - identity, composition

Full model

type : Obs $\mathbb{W} := [0, \infty]$ $\mathcal{B}_X :=$ 

$\mathcal{D}_X :=$ 

$P_X := \left\{ \mu \in \mathcal{D}_X \mid \underset{\mu}{\text{Ce}}[X] = 1 \right\}$

$\underset{\mu}{\text{Ce}}[E] :=$  $\delta_n :=$ 

$\phi_{\mu k} :=$ 

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

Please ask questions!
Sensible



← Course
web
page

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

TutorialFest
ACM-SIGPLAN 51st Symposium on
Principles of Programming Languages (POPL'24)
15 January, 2024
City of Westminster, England



THE UNIVERSITY OF EDINBURGH

informatics Ifcs

Laboratory for Foundations
of Computer Science

 BayesCentre

supported by:

 THE ROYAL
SOCIETY

The
Alan Turing
Institute

Facebook Research NCSC

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

Please ask questions!
Sensible



← Course
web
page


Full model

type : Obs $\mathbb{W} := [0, \infty]$ $\mathcal{B}_X :=$ 

$\mathcal{D}_X :=$ 

$\mathcal{P}_X := \left\{ \mu \in \mathcal{D}_X \mid \underset{\mu}{\text{Ce}}[X] = 1 \right\}$



$\underset{\mu}{\text{Ce}}[E] :=$  $\delta_n :=$ 

$\phi_{\mu k} :=$ 


Ref: Quasi-Banach space $X = (LX, \mathcal{R}_X)$

$$\mathcal{R}_X \subseteq LX^J \quad \text{Closed under:}$$

Set ↗
"carrier"
Set of
functions $\alpha: \mathbb{R} \rightarrow X$
"random elements"

- Constants: 
- Precomposition: 

- recombination



Examples

recombination of
constants

$$- \mathbb{R} = (\mathbb{R}, \text{Meas}(\mathbb{R}, \mathbb{R}))$$

abs underlying \mathbb{R}

$$- X \in \text{Set}, \quad \lceil X \rceil := (X, \sigma\text{-simple}(\mathbb{R}, X))$$

discrete qbs on X

$$- " \quad \lceil X \rceil_{\text{Qbs}} := (X, X^{\mathbb{L}(\mathbb{R})})$$

all functions

Indiscrete qbs on X

Validate gbs axioms for: $\mathbb{W} := ([0, \infty], \text{Meas}(R, \mathbb{W}))$

- Constants:

$$E : \mathcal{B}_{\mathbb{W}}, n : \mathbb{W} \vdash$$
$$(2_{r:R.x})^{-1}[E] = \begin{cases} x \in E : & R \\ n \notin E : & \emptyset \end{cases} \in \mathcal{B}_R$$

Validate gbs axioms for: $\mathbb{W} := ([0, \infty], \text{Meas}(R, \mathbb{W}))$

- Precomposition:

$\alpha: \text{Meas}(R, \mathbb{W}), \varphi: \text{Meas}(R, R) \vdash$

$$R \xrightarrow{\varphi} R \xrightarrow{\alpha} \mathbb{W} \quad \in \text{Meas}(R, \mathbb{W})$$

\Downarrow
Meas is a cat.

Explicitly:

$$(\alpha \circ \varphi)[E] \in \beta R \xleftarrow{\varphi^{-1}} \varphi[E] \in \beta R \xleftarrow{\alpha^{-1}} E \in \beta \mathbb{W} \quad \checkmark$$

Validate gbs axioms for: $\mathbb{W} := ([0, \infty], \text{Meas}(R, \mathbb{W}))$

- RL Combination

$I \text{ ctbl}, \alpha : \text{Meas}(R, \mathbb{W})^I, E_i : B_{\mathbb{W}}, R = \bigcup_{i \in I} E_i, F : B_{\mathbb{W}} \vdash$

$$\left(\exists r. \left\{ \begin{array}{l} : \\ r \in E_i : \alpha_i r \\ : \end{array} \right\}^{-1} [F] \right)$$

$$\beta :=$$

$$= \bigcup_{i \in I} \alpha_i^{-1}[F] \cap E_i \in B_R$$

In fact:


$$r \in \text{LHS} \Leftrightarrow \beta r \in F \Leftrightarrow \exists i \in I. r \in E_i \wedge \alpha_i r \in F \Leftrightarrow r \in \text{RHS}$$



σ-Simple function

$\alpha: \mathbb{R} \rightarrow X$ s.t. $\text{Im } \alpha := \alpha[\mathbb{R}]$ is ctbl 1

$\forall x \in \text{Im } \alpha. \alpha^{-1}[x] \in \mathcal{B}_{\mathbb{R}}$



Validate qbs axioms for: $\Gamma^{\text{qbs}} := (X, \sigma\text{-simple}(X))$

- Constants

$$\text{Im}(\lambda r. n) = \{n\} \text{ ctbl } \checkmark$$

NB: $f \sigma\text{-Simple:}$
 $\text{Im } f \text{ ctbl } 1$
 $\tilde{f}[x] \in B_R$

$$g: X \vdash (\lambda r. n)^{-1}[y] = \begin{cases} x=y: R \\ x \neq y: \emptyset \end{cases} \in B_R \checkmark$$

Validate gbs axioms for: $\overset{\text{gbs}}{X} := (X, \sigma\text{-simple}(X))$

• Precomposition:

$\alpha : \sigma\text{-simple}(X), \varphi : \text{Meas}(R, R) \vdash$

NB: $f \sigma\text{-Simple} : \text{Im } f \text{ ctbl } 1$
 $f^{-1}[x] \in B_R$

$$\text{Im}(\alpha \circ \varphi) \subseteq \text{Im } \alpha \text{ ctbl} \quad \checkmark$$

$x : X \vdash$

$$(\alpha \circ \varphi)^{-1}[x] = \varphi^{-1}[\alpha^{-1}(x)] \in B_R \quad \checkmark$$



$$\alpha^{-1}(x) \in B_R$$

$\varphi : R \rightarrow R$ measurable

Validate gbs axioms for: $\stackrel{\text{gbs}}{X} := (X, \sigma\text{-simple}(X))$

• recombination:

$$\alpha_- : (\sigma\text{-simple}(X))^I, E : \mathcal{B}_{IR}^I, R = \bigoplus_{i \in I} E_i +$$

NB: $f \sigma\text{-Simple}$:
 $\text{Im } f \subset \text{tbl } A$
 $f[x] \in \mathcal{B}_{IR}$

$$\text{Im}[E_i \cdot \alpha_i]_{i \in I} \subseteq \bigcup_{i \in I} \text{Im } \alpha_i \quad \text{tbl} \quad \checkmark$$

$x : X \vdash$

$$[E_i \cdot \alpha_i]_{i \in I}^{-1}(x) = \bigcup_{i \in I} \tilde{\alpha}_i^{-1}[x] \cap E_i \in \mathcal{B}_{IR} \quad \checkmark$$

Prop: $X: \text{Set}, A: \text{Qbs} \vdash$

• $\forall f: X \rightarrow {}_L A_1 . \quad f: {}^{\text{Qbs}}_X \rightarrow A$

• $\forall f: {}_L A_1 \rightarrow X . \quad f: A \rightarrow {}^X_{\text{Qbs}}$

Prop: $X: \text{Set}, A: \text{Qbs} \vdash$

- $\forall f: X \rightarrow {}_L A_1 . \quad f: {}^{Qbs}_X \longrightarrow A$

Prf: $\alpha: R_{r_{Qbs}}, \vdash \alpha \text{ } \sigma\text{-simple} \Rightarrow$

$$\alpha = \left[\alpha^{-1}[x].\lambda r. x \right]_{x \in \text{Im } \alpha} \Rightarrow$$

$$(f \circ \alpha) = \left[\alpha^{-1}[x].\lambda r. fx \right]_{x \in \text{Im } \alpha} \stackrel{\text{recombination}}{\Rightarrow} \in R_A$$

↑
constat $\in B_A$ ctbl

✓

Boole


Prop: $X: \text{Set}, A: \text{Qbs} \vdash$

- $\forall f: X \rightarrow {}_L A_J . f: {}^{Qbs}_X \rightarrow A$
- $\forall f: {}_L A_J \rightarrow X . f: A \rightarrow {}^X_{Qbs}$

Prf: $\alpha: R_A \vdash (f \circ \alpha: R \rightarrow X) \in R_{{}^X_{Qbs}}$ always. ✓



Useful adjunctions:



$$\mathcal{L}_{\text{Qbs}}^{\text{Meas}} := (\mathcal{L}_{\text{V}}, \text{Meas}(R, V)) \quad (V \in \text{Meas})$$

$$\mathcal{L}_{\text{X}}^{\text{Meas}} := \left\{ A \subseteq \mathcal{L}_{\text{X}} \mid \forall a \in R_x, a^{-1}[A] \in \mathcal{B}_R \right\}$$

- limits (products, subspaces)
and colimits (coproducts, quotients)
as in Set
- Slogan: every measurable space is carried by a qbs

Example

Product $(X \times Y, \pi_1, \pi_2)$:

necessarily!

$$- L_{X \times Y} = L_{X_1 \times_1 Y_1}$$

$$- R_{X \times Y} = \{ \lambda r. (\alpha r, \beta r) \mid \alpha \in R_X, \beta \in R_Y \}$$

correlated

random

elements

rest of structure as in Set.

Function Spaces

Straightforward!

$$- \mathbb{Y}^X := \text{Qbs}(X, \mathbb{Y})$$

$$- \mathbb{R}_{\mathbb{Y}^X} := \text{Uncurrying}[\text{Qbs}(\mathbb{R}^{XX}, \mathbb{Y})]$$

$$= \left\{ \alpha: \mathbb{R} \rightarrow \mathbb{Y}^X \mid \lambda(r, x). \alpha r x: \mathbb{R} \times X \rightarrow \mathbb{Y} \right\}$$

$$- \text{eval}: \mathbb{Y}^X \times X \rightarrow \mathbb{Y}$$
$$\text{eval}(f, x) := fx$$


Meas vs Qbs

By generalities:

σ -algebra
on $\text{Meas}(R, R)$

$$\Gamma^{\text{Meas}}(R) \rightarrow R \times R \rightarrow \Gamma^{\text{Meas}}(R) \times R \rightarrow \Gamma(R) = R$$

Γ^{Meas}



No factorisation
by
Aumann's
Theorem.

$$\left(\text{So } R^R \times R \neq (R^R \times R) \right)$$

Simple Type Structure

"Simple" because:

- Simply-typed λ -calculus
- types are simple: $A, B : \text{Type} \vdash B^A : \text{Type}$
 - no polymorphism
 - no term dependency
- Contexts for terms: $\Gamma \vdash t : A$
 - are simple: $\Gamma = x_1 : A_1, \dots, x_n : A_n$
 - i.e. $\text{List}(\text{Type})$

Simple Type Structure


"Simple" because:

- interpretation is simple:

$$\llbracket x_1 : A_1, \dots, x_n : A_n \rrbracket := \prod_{i=1}^n A_i$$

$$\llbracket \Gamma \vdash t : A \rrbracket : \llbracket \Gamma \rrbracket \longrightarrow A$$

in QBS



Simple Type Structure Curry-Howard-Lambek

$$\frac{\Gamma \vdash t : A \quad \Gamma \vdash s : B}{\Gamma \vdash \langle t, s \rangle : A \times B} \rightsquigarrow \boxed{\Gamma} \xrightarrow{\lambda r. \langle tr, sr \rangle} A \times B$$

is measurable

$$\frac{\Gamma \vdash t : A \times B \quad \Gamma, x:A, y:B \vdash s : C}{\Gamma \vdash \text{let } (x,y) = t \text{ in } s : C}$$

$$\Gamma \vdash \text{let } (x,y) = t \text{ in } s : C \rightsquigarrow$$

measurability
by
type!

$$\boxed{\Gamma} \xrightarrow{\lambda r. \text{let } (a,b) = tr \text{ in } sr[x \mapsto a, y \mapsto b]} C$$

is measurable . etc.

Random element Space

$R_X := X^{\mathbb{R}}$ since $\lfloor X^{\mathbb{R}} \rfloor = R_X$ as sets.

Why?

(\subseteq) $\alpha \in \lfloor X \rfloor^{\mathbb{R}} \Rightarrow \alpha: \mathbb{R} \rightarrow X$ in Qbs.

$\text{id}_{\mathbb{R}}: \mathbb{R} \rightarrow \mathbb{R}$ measurable $\Rightarrow \text{id} \in R_{\mathbb{R}}$

$\Rightarrow \alpha = \alpha \circ \text{id} \in R_X$

Pre composition

(\supseteq) $\alpha \in R_X \Rightarrow \exists \psi \in R_{\mathbb{R}} = \text{Meas}(\mathbb{R}, \mathbb{R})$. $\alpha \circ \psi \in R_X \Rightarrow \alpha: \mathbb{R} \rightarrow X$
 $\Rightarrow \alpha \in \lfloor X \rfloor^{\mathbb{R}}$

Subspaces

For $X \in \mathbb{Q}bs$, $A \subseteq X$, set:

$$R_A := \left\{ \alpha: \mathbb{R} \rightarrow A \mid \alpha \in R_X \right\}$$

Then $A = (A, R_A)$ is the *Subspace qbs*

We write $A \hookrightarrow X$

Borel Subspaces Ensemble

The σ -algebra $B_X := \left\{ A \subseteq X \mid \forall \alpha \in R_X . \alpha^*[A] \in B_R \right\}$


internalises as $B_X = 2^X$, the gbs of
Borel subsets.

$L^{(B_R)}$ are the Borel-on-Borel sets from
descriptive set theory.
(cf. [Sabou et al. '21])

Standard Borel Spaces

Def: A qbs S is Standard Borel when

$$S \cong A \text{ for some } A \in \mathcal{B}_{\mathbb{R}}$$



Slogan: Qbs Conservative extension of Sbs

Example $C_0 := \{f: \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ continuous}\} \hookrightarrow \mathbb{R}^{\mathbb{R}}$

C_0 is sbs. (Well-known!)

Proof:

$$C'_0 \in B_{\mathbb{R}^{\mathbb{Q}}}$$

sbs!

$$C'_0 := \left\{ g \in \mathbb{R}^{\mathbb{Q}} \mid \begin{array}{l} \forall a, b \in \mathbb{Q}, \varepsilon \in \mathbb{Q}^+ \\ \exists \delta \in \mathbb{Q}^+ \forall p, q \in \mathbb{Q} \cap [a, b], |p - q| < \delta \Rightarrow |g(p) - g(q)| < \varepsilon \end{array} \right\}$$

then $C_0 \cong C'_0 \in B_{\mathbb{R}^{\mathbb{Q}}}$:

$$c_0 \rightarrow c'_0$$

$$\varphi \mapsto \varphi|_{\mathbb{Q}}$$

$$c'_0 \rightarrow c_0$$

$$\varphi \mapsto \lambda r. \lim_{n \rightarrow \infty} g(\text{approx}_{\frac{1}{n}} \bigcup_{m \in \mathbb{N}} \{ \frac{1}{m} \})_n$$

on closed intervals
 (= compact intervals)

Continuity

uniform continuity

Borel measurable

by type checks

Example (ctd)

C_0 is sbs, and $\text{eval}: C_0 \times \mathbb{R} \rightarrow \mathbb{R}$
is measurable.

Avoids;

- constructing complete separable metrics
- proving that evaluation is measurable
w.r.t. metric σ -algebra.

Non-examples ~ [Sabot et al.'21]

$$- \left\{ A \in \mathcal{B}_{\mathbb{R}} \mid A \neq \emptyset \right\} \hookrightarrow \mathcal{B}_{\mathbb{R}}$$

$$- \left\{ (A_1, A_2) \in \mathcal{B}_{\mathbb{R}}^2 \mid A \subseteq B \right\} \hookrightarrow \mathcal{B}_{\mathbb{R}}^2$$

$$- \left\{ A \in \mathcal{B}_{\mathbb{R}} \mid A \text{ open} \right\} \hookrightarrow \mathcal{B}_{\mathbb{R}}$$

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces ✓
 - 5) Integration & random variables
- Lecture 1
- Lecture 2



Please ask questions!
Sensible

← Course
web
page

Dependent Type Structure

Types can contain terms :

$X:\text{Type}, E:B_X \vdash \{x \in X \mid n \in E\} : \text{Type}$

a type referring
to a term

a type, just like
STLC

a term!

Dependent Type Structure

Types can contain terms :

$$X:\text{Type}, E:B_X \vdash \{x \in |x \in E\} : \text{Type}$$

a type referring
to a term

a type, just like
STLC

Content formation:

$$\frac{\Gamma \vdash A : \text{Type}}{\Gamma, x:A \vdash}$$

Dependent Type Structure

types denote spaces-in-Content

$$\begin{array}{c} \llbracket \Gamma \vdash A \rrbracket \\ \downarrow \text{def} \\ \llbracket \Gamma \vdash \cdot \rrbracket \end{array}$$

Dependent types denote spaces-in-Content

$\Gamma \vdash$ \leftarrow Control

$[\Gamma \vdash A]$

\leftarrow Space in Content

$\Gamma \vdash A$

\leftarrow type in context

\downarrow dep

$[\Gamma]$

\leftarrow Context space

E.g.:

assigns

environment

A

\downarrow

1

simple types

$[\![E : B_A + \{x \in A \mid x \in E\}]\!]$

$\{ (E, a) \in B_A \times A \mid a \in E \}$

$\downarrow \pi_1$

B_A

decoder

Content extension

$$\frac{\Gamma \vdash A}{\Gamma, a:A \vdash}$$

$$\llbracket \Gamma \vdash A \rrbracket$$

$$dep$$

$$\llbracket \Gamma \rrbracket$$

$$\llbracket \Gamma, a:A \rrbracket := \llbracket \Gamma \vdash A \rrbracket$$

Substitution

$$\Gamma \vdash \sigma : \Delta$$


$$\llbracket \sigma \rrbracket : \llbracket \Gamma \rrbracket \rightarrow \llbracket \Delta \rrbracket$$

E.g. Weakening

$$\Gamma, a:A \vdash \text{wkn} : \Gamma$$

$$\llbracket \Gamma, a:A \rrbracket := \llbracket \Gamma \vdash A \rrbracket \xrightarrow[dep]{wkn} \llbracket \Gamma \rrbracket$$

Action of Substitution on Types



E.g.

$$\boxed{\forall x:A \vdash B[\omega_{\text{unif} }] := A \times B} \xrightarrow{\pi_2} B$$

$$\text{dep} := \pi_1, \quad \boxed{x:A} \xrightarrow{\Leftrightarrow} \mathbb{1}$$

Simple type

Terms: sections

$$\boxed{\Gamma} \xrightarrow{[\Gamma \vdash M : A]} \boxed{[\Gamma \vdash A]}$$

=

$\downarrow \text{dep}$

e.g.

$$R \xrightarrow{[\kappa : R \vdash (\kappa, \alpha) : B_R[\text{wkn}]]} R \times B_R$$

=

$\downarrow \pi_1$

R

E.g. Variables: $\boxed{[\Gamma, \alpha : A \vdash \alpha : A]}$

$$\boxed{[\Gamma, \alpha : A]} \xrightarrow{< \text{it}, \text{dep} >_{\Gamma \vdash A}} \boxed{[\Gamma, \alpha : A \vdash A[\text{wkn}]]}$$

=

$\downarrow \text{dep}$

Exercise:

action of substitution
 $M[\sigma]$

Dependent Pairs

$$\frac{\Gamma, a:A \vdash B}{\Gamma \vdash \prod_{a \in A} B}$$

$$\llbracket \prod_{a \in A} A \rrbracket := \llbracket \Gamma, a:A \vdash B \rrbracket$$

:=

$$\begin{array}{c} \downarrow \text{dep}_B \\ \llbracket \Gamma, a:A \rrbracket \\ \llbracket \Gamma \vdash A \rrbracket \\ \downarrow \\ \llbracket \Gamma \rrbracket \end{array}$$

dep_{\prod}

Dependent Products

$$\frac{\Gamma, a:A \vdash B}{\Gamma \vdash \prod_{a:A} B}$$

aha: $(a:A) \rightarrow B$

$$\Gamma \vdash \prod_{a:A} B$$

$$[\Gamma \vdash \prod_{a:A} B] :=$$

$$\left\{ (m_0, f : \{ a \in [A] \mid \text{dep } a = m_0 \} \rightarrow [\Gamma, a:A \vdash B]) \middle| \right. \\ \left. \forall a \in [\Gamma, a:A]. \text{dep } a = m_0 \Rightarrow \text{dep}(fa) = a \right\}$$

Exercise: find the random elements.

Full model

$$\text{type: Obs} \quad W := [0, \infty] \quad \mathcal{B}_X \cong \mathcal{B}^X$$

$$DX := (\text{Fri})$$

$$PX := \left\{ \mu \in DX \mid \underset{\mu}{\text{Ce}}[X] = 1 \right\}$$

$$\underset{\mu}{\text{Ce}}[E] := (\text{Fri}) \quad S_n := (\text{Fri})$$

$$\phi_{\mu k} := (\text{Fri})$$

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces ✓
 - 5) Integration & random variables
- Lecture 1
- Lecture 2



Please ask questions!
Sensible

← Course
web
page

Foundations for type-driven probabilistic modelling

Ohad Kammar
University of Edinburgh

TutorialFest
ACM-SIGPLAN 51st Symposium on
Principles of Programming Languages (POPL'24)
15 January, 2024
City of Westminster, England



THE UNIVERSITY OF EDINBURGH

informatics Ifcs

Laboratory for Foundations
of Computer Science

 BayesCentre

supported by:

 THE ROYAL
SOCIETY

The
Alan Turing
Institute

Facebook Research NCSC

Partiality cf. [Väkär et al., '19]

A Borel embedding $e: X \hookrightarrow Y$

- injective function $e: X \rightarrowtail Y$
- its image is Borel: $e[x] \in \mathcal{B}_Y$
- e is Strong: $\alpha \in R_X \Leftrightarrow e \circ \alpha \in R_Y$

Examples

- $\mathbb{1} \hookrightarrow \mathbb{2}$
- S is sbs $\Leftrightarrow \exists S \hookrightarrow \mathbb{R}$

Def: A Partial map $f: X \rightarrow Y$ is a morphism

$$f: X \rightarrow Y \amalg \{\perp\}$$

Its domain of definition

$$f: (Y \amalg \{\perp\})^X \vdash \text{Dom } f := \{x \in X \mid f_x \neq \perp\} : \text{Type}$$

Depent-type
interpretation

$$\begin{array}{ccc} \llbracket \text{Dom } f \rrbracket & \longrightarrow & \{g \in Y \mid g \in E\} \\ \downarrow \text{dep} & & \downarrow \text{dep} \\ \llbracket f: (Y \amalg \{\perp\})^X \rrbracket \llbracket \frac{E \mapsto x. f_x \neq \perp}{E: B_Y} \rrbracket & & \end{array}$$

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces ✓
 - 5) Integration & random variables
- Lecture 1
- Lecture 2

Please ask questions!
Sensible



← Course
web
page

Full model

type : Obs $\mathbb{W} := [0, \infty]$ $\mathcal{B}_X := \mathcal{B}^X$

$\mathcal{D}_X := \boxed{\quad}$

$\mathcal{P}_X := \left\{ \mu \in \mathcal{D}_X \mid \underset{\mu}{\text{Ce}}[X] = 1 \right\}$

$\underset{\mu}{\text{Ce}}[E] := \boxed{\quad}$ $\delta_n := \boxed{\quad}$

$\phi_{\mu k} := \boxed{\quad}$

Def: A measure μ over \mathbb{R} is a function

$$\mu : \mathcal{B}_{\mathbb{R}} \rightarrow \mathbb{W} := [0, \infty]$$

Satisfying the measure axioms:

$$E : \mathcal{B}^\omega \rightarrow$$

$$\mu \emptyset = 0, \quad \mu E = \mu(E \cap F) + \mu(E \cap F^c), \quad \mu(\cup_{n=1}^{\infty} E_n) = \sup_n \mu E_n$$

For measurable spaces, replace \mathbb{R} with V

We write $[GV]$ for the set of measures on V

For qbs X , take $[G^{r_{\text{meas}}} X]$

Thm (Lebesgue measure):

There is a unique measure $\lambda \in \mathcal{L}(\mathbb{R})$, s.t.:

$$\lambda(a, b) = b - a$$

The Unrestricted Giry Spaces


Equip $\llbracket GV \rrbracket$ with two gbs structures:

- X $R_{GV} := \{ \alpha: R \rightarrow GV \mid \forall A \in \mathcal{B}_V, \lambda r. \alpha(r, A): R \rightarrow \mathbb{W} \}$
- ✓ $GV \hookrightarrow \mathbb{W}^{B_X}$
- $\hookrightarrow \alpha$ is a kernel.
 - Fewer random elents
 - $R_{GV} \subseteq R_{G'V}$
 - Lebesgue integral measurable in both arguments.
(Upcoming)

Def: Simple function. $\varphi: X \rightarrow W$ when

$\exists n \in \mathbb{N}$, $\vec{A} \in \mathcal{B}_X^n$, $A_i \cap A_j = \emptyset$, $\vec{r} \in W$ s.t.
 $(i \neq j)$

$$\varphi_x = \begin{cases} \vdots & \\ x \in A_i : & r_i \\ \vdots & \\ \text{otherwise:} & 0 \end{cases}$$



Encode into a space:

$$\text{SimpleCode} := \prod_{n \in \mathbb{N}} \mathcal{B}_X^n \times \mathcal{W}^n$$

$$\text{Simple} := \{ f \in \mathcal{W}^X \mid f \text{ simple} \} \hookrightarrow \mathcal{W}^X$$

and define an interpretation:

$$[\![-]\!]: \text{SimpleCode} \longrightarrow \text{Simple}$$

$$[\![(n, \vec{A}, \vec{r})]\!] := \sum_{i=1}^n r_i \cdot [\!- \in A_i]\!]$$

↳ characteristic function
for A_i

Lemma: $f: X \rightarrow W$ is measurable → remember!
96s
morphism!

iff $f = \lim_{n \rightarrow \infty} f_n$ for some monotone sequence

$f_n \in \text{Simple}$.

Moreover, we have measurable such choice.


Simple Approx:

$\left\{ \vec{\alpha} \in \mathbb{R}^+ \mid \Delta_n \rightarrow 0 \right\} \times \left\{ \vec{\alpha}' \in W^N \mid \begin{array}{l} \vec{\alpha}' \text{ monotone} \\ a_n \rightarrow \infty \end{array} \right\} \times W \xrightarrow{X} \text{SimpleCode}$


↑
rate of convergence

↑
Range of approximation

Why?




Why?




resolution of
approximation


Why?




Why?




Why?



Why?




Why?



$$\begin{aligned} \text{Simple Approx. } \xrightarrow{\Delta_n \rightarrow 0} f &:= \sum_{i=1}^n i \cdot \Delta_n [i \cdot \Delta_n \leq f < (i+1) \cdot \Delta_n] \\ &\quad + \lfloor \frac{a_n}{\Delta_n} \rfloor \Delta_n \cdot [f \geq \lfloor \frac{a_n}{\Delta_n} \rfloor \cdot \Delta_n] \in \text{Simple} \end{aligned}$$

Integrating Simple Functions



$\int : G \times \text{Simple Code} \rightarrow W$

$$\int \mu(n, \vec{A}, \vec{r}) := \sum_{I \subseteq \{1, \dots, n\}} \left(\sum_{i \in I} r_i \right) \cdot \mu \left(\bigcap_{i \in I} A_i \setminus \bigcup_{i \notin I} A_i \right)$$

Integration

proper higher-order operation

$$\int : Gx \times W^X \longrightarrow W$$

$$\int \mu f := \sup \left\{ \int \mu \varphi \mid \varphi \in \text{Simple}, \varphi \leq f \right\}$$

measurable by type

$$= \lim_{n \rightarrow \infty} \int \mu (\text{Simple Approx}_{\vec{\Delta}, \vec{a}} f)_n$$

we also write

$$\int \mu(\delta x) t$$

$$\text{for } \int \mu(x, t)$$

for $\frac{a_n}{\Delta_n} \rightarrow 0$, e.g. $\Delta_n = \frac{1}{2^n}$ $a_n = n$.

resolution

The unrestricted Giry Strong Monad

Dirac:

$$\delta: X \rightarrow Gx$$

$$x \mapsto \lambda A. \begin{cases} x \in A : 1 \\ x \notin A : 0 \end{cases}$$

Unlike the unrestricted
Giry on Meas.

Kleisli extension/ Kock integral:

$$\oint : Gx \times Gr^x \rightarrow Gr$$

$$\oint \mu f := \lambda A. \int \mu(dx) f(x; A)$$

but: non-commutative

(Fubini fails,
just like in
Meas)

Fubini-Tonelli; fails

$$\# \in G/R \quad \# E := \begin{cases} E \text{ finite:} & |E| \\ \text{o.w.:} & \infty \end{cases}$$

$\lambda \in G/R$ Lebesgue $k: \mathbb{R} \times \mathbb{R} \rightarrow W \cong G/1$

$$\int \#(dx) \int \lambda(dx) k(x, y) = \int \# \Omega = \Omega \stackrel{\approx}{=} 0 \quad k(x, y) := [x=y]$$

$y: \mathbb{R} + \{\leftrightarrow\} \mapsto \lambda\{y\} \cdot 1 + \lambda\{y\}^c \cdot 0 = 0$ $\#$

$$\int \lambda(dx) \underbrace{\#(dx)}_{n \in \mathbb{R} + \{\leftrightarrow\}} k(x, y) = \int \lambda(dx) \delta_{\leftrightarrow} \stackrel{\approx}{=} \infty$$

$n \in \mathbb{R} + \{\leftrightarrow\} \mapsto \{x\} \cdot 1 + 0 = 1$

Randomizable measures monad

$$D \rightarrow G \quad \lambda A. \int_{\text{Dom } \alpha} \lambda(D \circ \alpha)$$

$$LDX := \left\{ \lambda \alpha \mid \alpha: \mathbb{R} \rightarrow X \right\}$$

Lebesgue measure

$$R_{Dx} := \left\{ \lambda x. \lambda_{\alpha x} \mid \alpha: \mathbb{R} \times \mathbb{R} \rightarrow X \right\}$$

$$\delta: x \rightarrow Dx \quad \oint: D^{\Gamma \times (Dx)} \rightarrow Dx \quad \text{lift along } D \rightarrow G.$$

D validates our measure axioms including Fubini-Tonelli
 $\mu \in DX, \nu \in DY \vdash$

$$\oint \mu(dx) \oint \nu(dy) \delta_{(x,y)} = \oint \nu(dy) \oint \mu(dx) \delta_{(x,y)} =: \mu \otimes \nu$$

Thm: For sbs S , $\text{PS}, D_{\leq S}, D_{<\infty S} \in \text{sbs}$

and agree with their counterparts on Meas.

$$DS_S = \{\mu \mid \mu \text{ s-finite}\}$$

see [Staton 46]

$$R_{DS} = \{K: \mathbb{R} \rightarrow G0 \mid K \text{ s-finite kernel}\}$$



Open: Is there a counterpart to D in Meas?

More modestly, is $DS \in \text{sbs}$?

(Hypothesis: **No**)

Distribution Submonads

A measure space

$$\Omega = (\Omega, \mu)$$

is a gbs Ω with
 $\mu \in D_X$.

Similarly:- finite measure space
- (Sub) Probability space.

$$P_X := \left\{ \mu \in D_X \mid \mu X = 1 \right\}$$

$$P_{\leq 1} X := \left\{ \mu \in D_X \mid \mu X \leq 1 \right\}$$

$$P_{<\infty} X := \left\{ \mu \in D_X \mid \mu X < \infty \right\}$$

$$D_X^T$$

Full model

type: Obs $W := [0, \infty]$ $\mathcal{B}X \cong \mathbb{B}^X$

$D_X := (\{\lambda_\alpha | \alpha: R \rightarrow X\}, \{\lambda_{r,-} | \alpha: R \times R \rightarrow X\})$

$P_X := \{\mu \in D_X | \text{Ce}_{\mu}[X] = 1\}$

$\text{Ce}_{\mu}[E] := \mu E$ $\delta_n := E \mapsto \begin{cases} n \in E : 1 \\ n \notin E : 0 \end{cases}$

$\oint \mu k := \lambda E. \int \mu(\lambda x) k(x; E)$

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces ✓
 - 5) Integration & random variables ✓
- Lecture 1
- Lecture 2


Please ask questions!
Sensible



← Course
web
page

Random variable: $\xi : \Omega \rightarrow \mathbb{H} \hookrightarrow \overline{\mathbb{R}}$

$\mathbb{H} :$



- \mathbb{H}^{Ω} is a space

- \mathbb{R}^{Ω} measurable vector space:

$$\alpha \xi + \zeta := \lambda \omega \cdot \alpha \cdot \xi \omega + \zeta \omega$$

- \mathbb{W}^{Ω} measurable σ -semi-module
for \mathbb{W} :

$$\sum_{n=0}^{\infty} \alpha_n \xi_m =$$

$$\lambda \omega \cdot \sum_{n=0}^{\infty} \alpha_n \cdot \xi_{m_n}$$

$$\Pr_r : P_{\Omega} \times \mathcal{B}_n \rightarrow \mathbb{W}$$

$$\Pr_\lambda A := \text{eval}(\lambda, A) = \lambda A$$

Probability Space $\mathcal{R} = (\Omega, \lambda_\Omega)$

$P : P_{\Omega} \vdash$ " P_{Ω} holds $\lambda(x)$ -almost surely "

for some $Q \subseteq \Omega$, $P \models Q$, $[-\infty Q] \cdot \lambda = \lambda$

Example $(\xi, \zeta \in \mathbb{H}^\Omega)$

$\xi = \zeta$ a.s. when $\Pr_{\omega \sim \lambda} [\xi \omega \neq \zeta \omega] = 0$

Integrating Random Variables (as discretely)

$(-)_{+}, (-)_{-} : \bar{\mathbb{R}}^{\mathbb{N}^2} \rightarrow \mathbb{W}^{\mathbb{N}^2}$ in Qbs,

$$\xi_{+} := \max(\xi, 0) \quad \xi_{-} := \max(-\xi, 0)$$

$$\text{So: } \xi = \xi_{+} - \xi_{-}$$

$$\int : P\mathcal{N} \times \mathbb{W}^{\mathbb{N}^2} \longrightarrow \mathbb{W} \quad \begin{cases} \text{respects} \\ \text{a.s. equality} \end{cases}$$

$$\int \lambda \xi := \int \lambda \xi_{+} - \int \lambda \xi_{-} \quad \begin{aligned} \xi &= \zeta \text{ (a.s.)} \\ \Rightarrow \int \lambda \xi &= \int \zeta. \end{aligned}$$

Example

$$\lambda: P\Omega \vdash \text{ASConverge}(\bar{\mathbb{R}})^{\mathbb{N}^2} : B(\bar{\mathbb{R}}^{N \times \mathbb{N}})$$
$$:= \left\{ \vec{z} \in \bar{\mathbb{R}}^{N \times \mathbb{N}} \mid \Pr_{w \sim \lambda} [\lim z_n w \neq \perp] \right\}$$

So:

$$\lim^{\text{as}}_m: \bar{\mathbb{R}}^{N \times \mathbb{N}} \longrightarrow \bar{\mathbb{R}}^{\mathbb{N}} \quad \text{Dom } \lim^{\text{as}} := \text{ASConverge}(\bar{\mathbb{R}})^{\mathbb{N}^2}$$

$$\lim^{\text{as}} \vec{z} := \lambda w. \limsup_{n \rightarrow \infty} f_n w$$

L \lim^{as} respects a.s. equality.

Thm (monotone convergence):

let $\sum \in \mathbb{W}^{N \times \omega}$ λ -a.s. monotone.

$$\xi = \lim_{n \rightarrow \infty} \xi_n \quad (\text{a.s.})$$



$$\int \lambda \xi = \lim_{n \rightarrow \infty} \int \lambda \xi_n$$

Lebesgue Space $\left(\Omega \text{ Prob. Space}, P \in [1, \infty) \right)$

$$P: [1, \infty), \lambda: P \mapsto L_{(\Omega, \lambda)}^P: \mathcal{B}(\mathbb{R}^\Omega)$$

$$:= \left\{ \xi \in \mathbb{R}^\Omega \mid \int |\xi|_P^P < \infty \right\} \hookrightarrow \mathbb{R}^\Omega$$

Ensemble $L_\Omega := \prod_{\lambda \in P_\Omega} L_{(\Omega, \lambda)}^P$

$$L \quad p \leq q \Rightarrow L_\Omega^p \supseteq L_\Omega^q$$


L^p semi norms

$$\| \cdot \| : \bigcup_{p,\lambda} L_{(2,\lambda)}^p \rightarrow \mathbb{R}_{\geq 0} \quad \| \xi \|_p := \sqrt[p]{\int \lambda |\xi|^p}$$


L^2 inner product

$$\langle \cdot, \cdot \rangle : \bigcup_{p,\lambda} L_{(2,\lambda)}^p \times L_{(2,\lambda)}^p \rightarrow \mathbb{R}$$

$$\langle \xi, \eta \rangle_p := \int \lambda \xi \eta$$



$$(\cdot)^{\rho}$$



Statistics

Expectation

$$\mathbb{E} : \bigcup_{\lambda} \mathcal{L}^1 \rightarrow \mathbb{R}$$

$$\mathbb{E}_{\lambda} \xi := \int \lambda \xi$$

Covariance and Correlation

$$\text{Cov}, \text{Corr} : \bigcup_{\lambda} \mathcal{L}^2 \rightarrow \mathbb{R}$$

$$\text{Cov}(\xi, \zeta) := \langle \xi - \mathbb{E} \xi, \zeta - \mathbb{E} \zeta \rangle$$

$$\text{Corr}(\xi, \zeta) := \frac{\langle \xi, \zeta \rangle}{\|\xi\|_2 \|\zeta\|_2} = \cos(\text{angle}(\xi, \zeta))$$

Sequential limits

$\rho: [1, \infty)$, $\lambda: P(X) \rightarrow$ Cauchy $L_{\lambda}^{\rho} := \mathcal{B}\left(\int_{(X, \Sigma)}^{\rho}\right)^N$

$$:= \left\{ \vec{\Sigma} \mid \forall \varepsilon \in \mathbb{Q}^+ \exists N \in \mathbb{N} \quad \forall m, n \geq N \quad \|\Sigma_{m+n} - \Sigma_{m+n}\|_{\rho} < \varepsilon \right\}$$

Thm: L_{λ}^{ρ} is Cauchy-complete

$\lim: \text{Cauchy } L_{\lambda}^{\rho} \rightarrow L^{\rho}$ (convergence in mean)

Why?

1. Every Cauchy sequence has an a.s. converging subseq.
2. We can find it measurable

Example

Theorem (dominated convergence)

For $\vec{z}_n, \vec{z} \in \mathbb{F}^l$ s.t. $\vec{z}_n \leq \vec{z}$ a.s.:

$$1. \lim^{as} \vec{z}_n \in \mathbb{F}^l$$

$$2. \lim^1 \vec{z}_n = \lim^{as} \vec{z}_n$$

$$3. \lim_{n \rightarrow \infty} \int \lambda \vec{z}_n = \int \lambda \lim_{n \rightarrow \infty} \vec{z}_n$$

Separability

Def: L^P separable: has countable dense subset

Fact: Separability is property of λ_2 :

TFAE:

- $\exists P \geq 1$. L^P separable
- $\forall P \geq 1$. L^P separable

Measurable separability in $I \hookrightarrow P\Omega \times [1, \infty)$

$$\vec{\beta} : \prod_{(\lambda, p) \in J} L_{(\Omega, \lambda)}^p \xrightarrow{IN} \text{S.t.}$$

$$\left\{ \vec{\beta}_n^{(p)} \mid n \in \mathbb{N} \right\} \text{ dense in } L_{(\Omega, \lambda)}^p$$

Prop. - Every SBS S measurable separable in
 $PS \times [1, \infty)$

- $I \hookrightarrow P\Omega \times \{2\}$ measurable separable

$$\Rightarrow \exists \vec{\beta} \in \prod_{\lambda \in J} L_{(\Omega, \lambda)}^2 \text{ orthonormal system } (\beta_n) \text{ dense}$$

$$\begin{aligned} \langle \beta_n, \beta_m \rangle &= 0 \\ \|\beta_n\|_2 &= 1 \end{aligned}$$

Escape

Let $S \hookrightarrow L^2$ closed Vector Subspace.

Orthogonal decomposition linear in fact.

$$\langle P, P^\perp \rangle : L^2 \rightarrow S \times S^\perp$$

When S is separable with orthonormal system β


We have a measurable version of

$$\langle P, P^\perp \rangle : L^2 \rightarrow S \times S^\perp$$

$$P\xi := \sum_{n=0}^{\infty} \langle \xi, \beta_n \rangle \beta_n \quad P^\perp := \text{Id} - P .$$

Kolmogorov's Conditional Expectation


ground truth space



(H)

Sample space

H
observation



Σ
Statistics
of interest

!

R

conditional expectation

$$\mathbb{E}[\Sigma | H = -]$$

Observed
statistic

Kolmogorov's Conditional Expectation

A Conditional expectation


of $\xi \in L^1_{\Omega}$ wrt

$H: \Omega \rightarrow \mathbb{H}$ is

$\xi \in L^1_H$ s.t. for all $A \in \mathcal{B}_H$:

$$\int_A \mu \xi = \int_{H^{-1}[A]} \lambda \xi$$

Where $\mu := \lambda_H$




Conditional expectations

1. unique a.s.

2. fundamental to modern Probability, e.g.:

a Martingale



$$\text{St. } \xi_n = \mathbb{E}[\xi_{n+1} | H_n = -]$$

Theorem (Existence)

- $\exists \mathbb{E}[-|H=-]: \mathcal{L}_{(\Omega, \lambda)}^1 \rightarrow \mathcal{L}_{(\mathbb{H}, \mu)}^1$

- When (Ω, λ) is Separable

$$\mathbb{E}[-|H=-]: \mathcal{L}_{(\Omega, \lambda)}^1 \rightarrow \mathcal{L}_{(\mathbb{H}, \mu)}^1$$

- When \mathbb{H} is \mathcal{I} -measurably separable

$$\mathbb{E}[-|H=-]: \prod_{\substack{H \in \mathbb{H} \\ \lambda \in H^*[\mathcal{I}]}} \mathcal{L}_{(\Omega, \lambda)}^1 \rightarrow \mathcal{L}_{(\mathbb{H}, \mu)}^1$$

Plan:

- 1) Type-driven probability: discrete case ✓
 - 2) Borel sets & measurable spaces ✓
 - 3) Quasi Borel spaces ✓
 - 4) Type structure & standard Borel spaces ✓
 - 5) Integration & random variables ✓
- Lecture 1
- Lecture 2

please ask questions!
sensible



← Course
web
page

Discrete model

type: set $\mathbb{W} := [0, \infty]$ $\mathcal{B}X := \mathcal{P}X$

$\mathcal{D}X := \{\mu: X \rightarrow \mathbb{W} \mid \text{Supp } \mu \text{ countable}\}$

$\mathcal{P}X := \{\mu \in \mathcal{D}X \mid \underset{\mu}{\text{Ce}}[X] = 1\}$

$\underset{\mu}{\text{Ce}}[E] := \sum_{x \in E} \mu_x \quad \delta_x := \lambda x. \begin{cases} x = x': 0 \\ x \neq x': 1 \end{cases}$

$\oint \mu k := \lambda x. \sum_{m \in \Gamma} \mu^m \cdot k(m; x)$

Full model

type: Obs $W := [0, \infty]$ $\mathcal{B}X \cong \mathbb{B}^X$

$D_X := (\{\lambda_\alpha | \alpha: R \rightarrow X\}, \{\lambda_{r,\cdot}\}_{\alpha(r,-)} | \alpha: R \times R \rightarrow X\})$

$P_X := \{\mu \in D_X | \text{Ce}_\mu[X] = 1\}$

$\text{Ce}_\mu[E] := \mu E$ $\delta_n := E \mapsto \begin{cases} n \in E : 1 \\ n \notin E : 0 \end{cases}$

$\oint \mu k := \mathbb{E} . \int \mu(d\alpha) k(\alpha; E)$

Advertisement

Interested in the mathematical } foundations of Computer
Computational logical Science ?

Check out our PhD Programs:



THE UNIVERSITY of EDINBURGH
informatics

lfcs

Laboratory for Foundations
of Computer Science



LFCS
PhD
↗



Repetile
AI for
Robotics
CDT

Enough!

Lunch.