

Explaining the Relativistic Nature of General and Special Relativity

Thomas DeGerlia^a

Originally published: June 16, 2026 C.E.

Revision 2.0 June 16, 2026 C.E.

ABSTRACT: In this paper, we reveal that the time dilation between two systems is equal to the ratio of their respective DeGerlia inertial density D , defined as mass over radius. Fundamental to this exploration is the recognition that relativity is always relativistic, meaning there is no single privileged frame of observation: no measurement, whether of time or anything else, has any meaning without a comparator. We reveal this by taking our relative compactness equation, plugging in a system compared to that same system at its Schwarzschild radius, and demonstrating that we obtain the same exact result as the gravitational time dilation equation. In order to present this equivalence and demonstrate the relativistic principle, we utilize a framework developed by the author called the DeGerlia Time Dilation framework, a simple, mathematically equivalent time dilation framework that is directly derived from general relativity.

Keywords: time dilation, general relativity, special relativity, gravitational time dilation, Lorentz time dilation, DeGerlia inertial density, inertial density, moment of inertia, Schwarzschild radius

1 Introduction

General relativity is a model for observing our universe that is fundamentally comparative: no quantity has physical meaning without a comparator. The pace of time is no exception: the time dilation between two systems is a product of the systems respective geometric mass distribution. Directly from the gravitational time dilation formula one can derive the underlying two-system logic and ultimately GTD is simply a comparison between a system a systems pace of time relative to the pace of time of an arbitrarily selected benchmark system, which in this case is the same system at its Schwarzschild radius. The formula allows you the freedom to only insert one mass and one radius, and the second is derived directly from the first. This has the effect of hiding the true nature of relativity and that is that it is, above all, comparative. But sends a signal that one need not compare two systems to understand time, and that is flawed thinking; it introduces the mathematical fallacy of an arbitrarily selected comparator, when there is, in fact, no privilege perspective from which to understand time. In this paper we evaluate the physical property responsible for relativistic time dilation and derive the underlying formulation that quantifies the relative time dilation between two systems.

The relative flow of time between two systems is dictated by their relative compactness. The standard formula for compactness, described under general relativity, expresses this as a dimensionless parameter. However, in this paper we utilize a principle developed by the author, DeGerlia inertial density, which is an abstracted compactness measure explained herein, that retains the proper unit of this property and that is mass over radius. Thus DeGerlia inertial density is defined as a system's mass over radius $D = M/R$ (DeGerlia, 2025b). We do a comparative analysis and tabulate relative time dilation across a variety of systems from classical to relativistic regimes, and show that the same compactness ratio $k = D_1/D_2 = M_1R_2/(M_2R_1)$ returns the Schwarzschild time-dilation ratio exactly across every tested constraint class, including the fully general case. This establishes that the relative rate of time between systems is not an ad hoc relativistic output requiring separate formulas, but a direct consequence of comparative mass-radius geometry. The result makes explicit a simple relation already contained in general relativity: given two systems' masses and radii, their relative time dilation is determined by their respective geometric distribution of mass alone.

Several distinct, common forms of time dilation utilize the same formula, and that is that the time dilation is equal to $\sqrt{1-k}$, the square root of one minus the scale factor. In general relativity, the scale factor is calculated by r_s/R . In special relativity, it's calculated as v^2/c^2 , and in general relativity, you can also substitute that: r_s/R could be replaced with D/D_{crit} . Those are all equivalent.

Both General and Special Relativity follow a very similar form, and their distinction mirrors the distinction between linear and rotational inertia. This alludes to a very rational and logical uni-

^a DeGerlia Expert Consulting, 3000 Lawrence Street, Denver, CO, United States of America. E-mail: tom.degerlia@tomdegerlia.com

fication of relativity.

2 Experimental Model

The experimental model employed in this study rests on several key components. The framework is derived directly from general relativity and is therefore mechanically and mathematically equivalent to well-established general-relativistic calculations. It does not rely upon, nor attempt to introduce, any principles not already established within the theories of general and special relativity; it is simply a re-framing of the existing theory of general relativity for exploratory purposes.

We begin by introducing the DeGerlia Framework and then demonstrate its complete mathematical equivalence with general relativity. We then work through a series of examples, each one demonstrating a different comparative case, and in each case we validate the result using standard gravitational time dilation calculations.

The first example is the most familiar: the case in which both systems have the same mass. Gravitational time dilation is such a case, because to calculate it one must compare a system with an identical system at its Schwarzschild radius. This example demonstrates the two-system nature of all relativistic calculations by comparing two systems of equal mass. Relativity is fundamentally concerned with how quantities appear relative to one another; gravitational time dilation reflects the comparison of systems of equal mass, where the compared radius is the collapse-condition radius r_s . We then work through a second equal-mass calculation, comparing an arbitrarily large system to a system of equal mass at four-thirds the Schwarzschild radius, or $d\tau/dt = 0.5$. We work through two systems of equal radius, which represent the Lorentz factor in special relativity. We also perform calculations for systems of equal density or equal moment of inertia, in addition to the classical case in which we compare near-scale objects, and a general case with no constraint, in which one may identify the relative pace of time between any two systems bearing no similarity to one another.

3 General Relativity

3.1 Compactness

Compactness is the dimensionless ratio of a system's mass to its radius, scaled by G/c^2 . The most common forms are $C = GM/(Rc^2)$ and $r_s/R = 2GM/(Rc^2) = 2C$, where $r_s = 2GM/c^2$ is the Schwarzschild radius. Both measure how close a system sits to its Schwarzschild threshold: $r_s/R = 1$ marks the gravitational-collapse condition. However, because compactness is normalized, it treats our scale as privileged, and thus begins to offer a less and less accurate view the further one moves from our own scale.

$$C = \frac{GM}{Rc^2} \quad (1)$$

where this is the standard general-relativistic compactness, $\mathcal{C} \equiv GM/(Rc^2)$ Alho et al. (2022).

3.2 Time Dilation

General Relativity defines the formula to calculate a system's time dilation relative to flat space-time:

$$d\tau/dt = \sqrt{1-k} \text{ where } k = r_s/R = 2GM/(Rc^2).$$

3.3 Structure of the GTD Equation as a Two-System Comparison

While the above equation requires only a mass and radius, "flat space-time" actually describes a second system, which is the same mass at its Schwarzschild radius. This is a two-system comparison, and the second system is silently approximated as the same mass at its Schwarzschild radius. This is a concrete reference to an arbitrarily determined comparison system. The DeGerlia framework makes this two-system comparison explicit and concrete. Because the radius is calculated from the inout mass, it is already a relative reference, relative to its own Schwarzschild radius.

4 The DeGerlia Time Dilation Framework

The DeGerlia Time Dilation Framework is an analogous framework that is directly derived from general relativity and has a direct relationship to gravitational time dilation. DeGerlia Time Dilation DTD shares a Pythagorean relationship with GTD such that $DTD^2 + GTD^2 = 1$.

4.1 Definitions used within the DeGerlia Time Dilation Framework

Symbol	Definition / value
GTD	$\sqrt{1-k} = d\tau/dt$
DTD	$\sqrt{k}$ (DeGerlia time dilation)
k	$v^2/c^2 = r_s/R = D/D_{\text{crit}}$
D	M/R_{axis} (DeGerlia inertial density)
D_{crit}	$c^2/2G = 6.73295 \times 10^{26} \text{ kg/m}$
r_s	$2GM/c^2$ (Schwarzschild radius)
P	I/V (inertial density) $= D \cdot 3/10\pi$
θ	$\arcsin \sqrt{k}$; $GTD = \cos \theta$, $DTD = \sin \theta$
c	299,792,458 m/s (exact)
G	$6.67430 \times 10^{-11} \text{ m}^3\text{kg}^{-1}\text{s}^{-2}$

4.2 Inertia

4.3

4.4 Inertial Density

4.5 DeGerlia Inertial Density

4.6 System Time

4.7 System

Under the DeGerlia Time Dilation Framework a "system" is defined as "any discrete collection of particles and the space between them", or alternatively, "any unit of space and the particles contained within." Because this framework sees the universe not as a clean separation of discrete material systems separated by the vacuum of space, but as a gradient of material density pervasive everywhere throughout the universe. The most extreme densities

reside not within near-scale systems but rather, within the cores of highly dense gravitational objects far from our physical scale, such as black holes or sub-atomic particles, and forms a gradient all the way to the near-scale "vacuum of space" where another gravitational source begins to exert more influence. It does not concern itself with notions of uniformity or even symmetry at all. Sphericalness relies on none of those. It doesn't rely on a clean edge separating matter from vacuum, or anything along those lines. It doesn't rely on static density nor any particular density gradient.

It only requires a mass and a radius about the axis of observation relative to an observing mass and radius about that same axis of observation; if you're not getting sufficient granularity within the radius you selected, pick a smaller radius and from that you can derive more information about the interior of that object. One can simply say my system consists of the earth to a radius of x where that radius might encompass just the radius of the planet. But if you were talking about Jupiter without a clear periphery separating the surface from the atmosphere, one might characterize the system as out to a radius of x or out to a density of y . If analyzing a large system, it would not be uncommon to take a spherical sample from within that system in order to characterize the behavior of certain regions.

4.8 The Linear Scale Factor k

The following identity relates the time-dilation ratio of two uniform spherical systems to their mass and radius.*

$$k = \frac{D_1}{D_2} = \frac{M_1 R_2}{M_2 R_1} \quad (2)$$

where R_1, R_2 are the radii, M_1, M_2 are the masses, and $D_i = M_i/R_i$ is the DeGerlia inertial density of system i . The three forms in Equation 2 are algebraically identical; $(I_1/I_2)(R_2/R_1)^3$ exposes the inertia-radius structure, D_1/D_2 exposes the inertial density ratio structure, and $M_1 R_2/(M_2 R_1)$ is the fully expanded form.

A system's pace of time has meaning only relative to another pace of time. The scale factor k governs the relative time-dilation between two uniform spherical systems. The standard gravitational time-dilation calculation takes the Schwarzschild form $\sqrt{1-k}$, where $k = D/D_{\text{crit}} = r_s/R$. The compactness ratio $k = D_1/D_2$ and the Schwarzschild form give the same value.

Via the Schwarzschild radius, the same scale factor is:

$$k = \frac{D}{D_{\text{crit}}} = \frac{r_s}{R} \quad (3)$$

where $D = M/R$ is the system's DeGerlia inertial density and $r_s = 2GM/c^2$ is its Schwarzschild radius. This is the k that appears in the standard $\sqrt{1-k}$ for GTD.

Time dilation is always:

$$\frac{d\tau}{dt} = \sqrt{1-k} \quad (4)$$

where $d\tau/dt$ is the time-dilation factor between the two systems

and k is the DeGerlia inertial density ratio for the case being computed. The specific forms are:

- $k = D_1/D_2$ (Equation 2) — via inertial density ratio
- $k = D/D_{\text{crit}} = r_s/R$ (Equation 3) — via Schwarzschild radius

Same k symbol across forms: a given numerical k produces the same time-scale-factor result regardless of which form it appears in.

4.9 DeGerlia inertial density D

Inertial density P (DeGerlia, 2025b) is the moment of inertia of a system divided by its volume:

$$P = I/V \quad (5)$$

Inertia, in a linear context, is simply mass. Inertial density, in this context, reduces directly to mass density $\rho = M/V$. From this perspective, inertial density is simply the rotational analog to mass density, not unlike the relationship between f equals m and torque equals angular acceleration times m . More succinctly stated: mass density is simply the linear interpretation of inertial density.

Substituting $I = \frac{2}{5}MR^2$ and $V = \frac{4}{3}\pi R^3$, with R the radius, this reduces to:

$$P = \frac{M}{R} \times \frac{3}{10\pi} \quad (6)$$

Just as ordinary density ($\rho = M/V$) is linear inertia (mass) the distributed over volume, DeGerlia inertial density is rotational inertia distributed over volume. As stated in the prior work: *mass and density are to linear motion as moment of inertia and inertial density are to rotational motion*.

$D = M/R$ is a simplified analog to inertial density, related to P by the geometric factor $3/10\pi$, and represents the mass over radius ratio for a system. Given that this quantity is almost always used in a ratio that would cancel out all geometric vectors and constants, for ease of use, this metric D for inertial density is typically used over the geometrically accurate inertial density, P .

4.10 DeGerlia Threshold D_{crit}

The Schwarzschild condition can be expressed as a static universal constant threshold of DeGerlia inertial density D . There are several advantages to this approach, not the least of which is that you don't have to calculate the threshold for every mass. And, it's much easier to calculate D than P . Inertial density is an intuitive metric that's easier to work with.

Setting $R = r_s$ (the Schwarzschild condition) (Schwarzschild, 1916) and solving yields the **DeGerlia Threshold**: the universal constant of DeGerlia inertial density at which the escape velocity equals the speed of light:

$$D_{\text{crit}} = \frac{c^2}{2G} = (6.73295 \pm 0.00015) \times 10^{26} \text{ kg/m} \quad (7)$$

Note this value is exact; its precision is limited by $G = (6.67430 \pm 0.00015) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ (Tiesinga et al., 2021).

* This identity has been previously referred to as the "space-time equivalence" DeGerlia (2025a).

4.11 DeGerlia Time Dilation

DeGerlia Time Dilation is the geometric complement of gravitational time dilation:

$$\text{DTD} = \sqrt{k} \quad (8)$$

where k is the scale factor of Equation 2. Where $\text{GTD} = \sqrt{1-k}$ is the surviving proper-time rate $d\tau/dt$, $\text{DTD} = \sqrt{k}$ is its complement.

4.12 Converting Between GTD and DTD

GTD and DTD are bound by the Pythagorean identity:

$$\text{GTD}^2 + \text{DTD}^2 = (1-k) + k = 1. \quad (9)$$

4.13 Properties of DTD

5 Relative Time Dilation Examples

Tables 1 and 2 present the complete system properties and derived ratios for seven system pairs spanning the constraint cases that follow: static mass ($M = M$), static radius ($R = R$), constant density ($\rho = \rho$), constant inertia ($I = I$), an unconstrained general case, and a classical case. Across every case, the three algebraic forms of the scale factor k (Equation 2) produce identical values, and the time-dilation ratio computed via the DeGerlia route agrees with the gravitational time-dilation ratio computed independently from the standard Schwarzschild form — to the precision afforded by G . No case requires a separate formula. The worked examples in the subsections that follow show, case by case, exactly how each column of these tables is obtained.

5.1 Case 1: ($M_1 = M_2$, $R_1 \approx r_s$)

The two identical masses cancel, reducing the DeGerlia inertial density ratio down to a ratio of radii.

$$k_d = \frac{M_1 R_2}{M_2 R_1 D_{\text{crit}}} = \frac{R_2}{R_1 D_{\text{crit}}}$$

Note, this reflects the k calculation for gravitational time dilation, where we are comparing with r_s . Because $\text{GTD} = 0$ exactly at r_s , we set the comparison radius to $1.00001 r_s$, an "arbitrarily" small amount above r_s .

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	2.00000e30	2.00000e30
R (m)	2.97049e3	7.00000e8

Calculate DTD for each System:

$$\begin{aligned} \text{DTD} &= \sqrt{k} \quad \text{where } k = M/(R D_{\text{crit}}) \\ k_1 &= \frac{2.00000e30 \text{ kg}}{(2.97049e3 \text{ m})(6.73295e26 \text{ kg/m})} = 9.9999068441e-1 \\ \text{DTD}_1 &= \sqrt{9.9999068441e-1} = 9.9999534220e-1 \\ k_2 &= \frac{2.00000e30 \text{ kg}}{(7.00000e8 \text{ m})(6.73295e26 \text{ kg/m})} = 4.24352e-6 \\ \text{DTD}_2 &= \sqrt{4.24352e-6} = 2.05998e-3 \\ \text{DTD}_1 &= \frac{9.9999534220e-1}{2.05998e-3} = 4.85439e2 \\ \text{DTD}_2 &= \end{aligned}$$

Calculate GTD ratio from DTD:

$$\begin{aligned} \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= \frac{\sqrt{1 - \text{DTD}_1^2}}{\sqrt{1 - \text{DTD}_2^2}} = \frac{\sqrt{1 - (9.9999534220e-1)^2}}{\sqrt{1 - (2.05998e-3)^2}} \\ &= \frac{\sqrt{1 - 9.9999068441e-1}}{\sqrt{1 - 4.24352e-6}} \\ &= \frac{\sqrt{9.31559e-6}}{\sqrt{9.9999575648e-1}} \\ &= \frac{3.05214e-3}{9.9999787824e-1} = 3.05215e-3 \\ &= 1 - 9.9694785e-1 \end{aligned}$$

Verify by Calculating GTD from Schwarzschild Radius Ratio: Schwarzschild radii:

$$\begin{aligned} r_{s1} = r_{s2} &= \frac{2GM}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(2.00000e30 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 2.97046e3 \text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{2.97046e3 \text{ m}}{2.97049e3 \text{ m}} = 9.999900010e-1 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{2.97046e3 \text{ m}}{7.00000e8 \text{ m}} = 4.24352e-6 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

$$\begin{aligned} \text{GTD}_{s1} &= \sqrt{1 - k_1} = \sqrt{9.99990e-6} = 3.16226e-3 = 1 - 9.9683774e-1 \\ \text{GTD}_{s2} &= \sqrt{1 - k_2} = \sqrt{9.9999575648e-1} = 9.9999787824e-1 \\ &= 1 - 2.12176e-6 \\ \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= \frac{3.16226e-3}{9.9999787824e-1} = 3.16227e-3 \\ &= 1 - 9.9683773e-1 \end{aligned}$$

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\begin{aligned} \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= 1 - 9.9683773e-1 \quad (\text{via } r_s) \\ \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= 1 - 9.9694785e-1 \quad (\text{via } D_{\text{crit}}) \end{aligned}$$

Table 1 System properties for each pair comparison case.

	Case 1 $M=M_j \sim r_s$	Case 2 $M=M_j \sim \frac{4}{3}r_s$	Case 3 $R=R$	Case 4 $\rho=\rho$	Case 5 $I=I$	Case 6 General	Case 7 Classical
M_1 (kg)	2.00000e+30	1.00000e+30	1.00000e+30	1.00000e+30	1.00000e+32	1.00000e+20	8.00000e+0
M_2 (kg)	2.00000e+30	1.00000e+30	1.00000e+24	1.00000e+27	1.00000e+30	1.00000e+30	1.00000e+0
R_1 (m)	2.97049e+3	1.98031e+3	1.00000e+8	1.00000e+8	1.00000e+7	2.00000e-7	2.00000e+0
R_2 (m)	7.00000e+8	1.00000e+8	1.00000e+8	1.00000e+7	1.00000e+8	1.00000e+7	1.00000e+0
ρ_1 (kg/m ³)	1.82161e+19	3.07406e+19	2.38732e+5	2.38732e+5	2.38732e+10	2.98416e+39	2.38732e-1
ρ_2 (kg/m ³)	1.39203e+3	2.38732e+5	2.38732e-1	2.38732e+5	2.38732e+5	2.38732e+8	2.38732e-1
I_1 (kg·m ²)	7.05907e+36	1.56865e+36	4.00000e+45	4.00000e+45	4.00000e+45	1.60000e+6	1.28000e+1
I_2 (kg·m ²)	3.92000e+47	4.00000e+45	4.00000e+39	4.00000e+40	4.00000e+45	4.00000e+43	4.00000e-1
D_1 (kg/m)	6.73289e+26	5.04972e+26	1.00000e+22	1.00000e+22	1.00000e+25	5.00000e+26	4.00000e+0
D_2 (kg/m)	2.85714e+21	1.00000e+22	1.00000e+16	1.00000e+20	1.00000e+22	1.00000e+23	1.00000e+0
r_{s1} (m)	2.97046e+3	1.48523e+3	1.48523e+3	1.48523e+3	1.48523e+5	1.48523e-7	1.18819e-26
r_{s2} (m)	2.97046e+3	1.48523e+3	1.48523e-3	1.48523e+0	1.48523e+3	1.48523e+3	1.48523e-27
$D_{\text{norm},1}$	9.99991e-1	7.50001e-1	1.48523e-5	1.48523e-5	1.48523e-2	7.42617e-1	5.94093e-27
$D_{\text{norm},2}$	4.24352e-6	1.48523e-5	1.48523e-11	1.48523e-7	1.48523e-5	1.48523e-4	1.48523e-27
k_s	9.99990e-1	7.50000e-1	1.48523e-5	1.48523e-5	1.48523e-2	7.42616e-1	5.94093e-27
k_d	9.99991e-1	7.50001e-1	1.48523e-5	1.48523e-5	1.48523e-2	7.42617e-1	5.94093e-27
GTD _{s1}	1-9.96838e-1	1-5.00000e-1	1-7.42619e-6	1-7.42619e-6	1-7.45394e-3	1-4.92670e-1	1-2.97046e-27
GTD _{s2}	1-2.12176e-6	1-7.42619e-6	1-7.42616e-12	1-7.42616e-8	1-7.42619e-6	1-7.42644e-5	1-7.42616e-28
GTD _{d1}	1-9.96948e-1	1-5.00001e-1	1-7.42619e-6	1-7.42619e-6	1-7.45395e-3	1-4.92670e-1	1-2.97047e-27
GTD _{d2}	1-2.12176e-6	1-7.42619e-6	1-7.42617e-12	1-7.42617e-8	1-7.42619e-6	1-7.42644e-5	1-7.42617e-28
DTD ₁	9.99995e-1	8.66026e-1	3.85387e-3	3.85387e-3	1.21870e-1	8.61752e-1	7.70774e-14
DTD ₂	2.05998e-3	3.85387e-3	3.85387e-6	3.85387e-4	3.85387e-3	1.21870e-2	3.85387e-14

Table 2 Derived ratios across constraint cases.

	Case 1 $M=M_j \sim r_s$	Case 2 $M=M_j \sim \frac{4}{3}r_s$	Case 3 $R=R$	Case 4 $\rho=\rho$	Case 5 $I=I$	Case 6 General	Case 7 Classical
DTD ₁ /DTD ₂	4.85439e+2	2.24716e+2	1.00000e+3	1.00000e+1	3.16228e+1	7.07107e+1	2.00000e+0
$\sqrt{D_1/D_2}$	4.85439e+2	2.24716e+2	1.00000e+3	1.00000e+1	3.16228e+1	7.07107e+1	2.00000e+0
$\sqrt{(M_1 R_2)/(M_2 R_1)}$	4.85439e+2	2.24716e+2	1.00000e+3	1.00000e+1	3.16228e+1	7.07107e+1	2.00000e+0
$\sqrt{(I_1/I_2)(R_2/R_1)^3}$	4.85439e+2	2.24716e+2	1.00000e+3	1.00000e+1	3.16228e+1	7.07107e+1	2.00000e+0
$(I_1/I_2)^{1/5}(\rho_1/\rho_2)^{3/10}$	4.85439e+2	2.24716e+2	1.00000e+3	1.00000e+1	3.16228e+1	7.07107e+1	2.00000e+0
$(M_1 R_2)/(M_2 R_1)$	2.35651e+5	5.04972e+4	1.00000e+6	1.00000e+2	1.00000e+3	5.00000e+3	4.00000e+0
D_1/D_2	2.35651e+5	5.04972e+4	1.00000e+6	1.00000e+2	1.00000e+3	5.00000e+3	4.00000e+0
$(I_1/I_2)(R_2/R_1)^3$	2.35651e+5	5.04972e+4	1.00000e+6	1.00000e+2	1.00000e+3	5.00000e+3	4.00000e+0
GTD _{s1} /GTD _{s2}	1-9.96838e-1	1-4.99996e-1	1-7.42618e-6	1-7.35193e-6	1-7.44657e-3	1-4.92632e-1	1-2.22785e-27
GTD _{d1} /GTD _{d2}	1-9.96948e-1	1-4.99997e-1	1-7.42619e-6	1-7.35193e-6	1-7.44658e-3	1-4.92633e-1	1-2.22785e-27
$\sqrt{1-\text{DTD}_1^2}/\sqrt{1-\text{DTD}_2^2}$	1-9.96948e-1	1-4.99997e-1	1-7.42619e-6	1-7.35193e-6	1-7.44658e-3	1-4.92633e-1	1-2.22785e-27
I_1/I_2	1.80078e-11	3.92163e-10	1.00000e+6	1.00000e+5	1.00000e+0	4.00000e-38	3.20000e+1
$(I_1/I_2)^{1/2}$	4.24356e-6	1.98031e-5	1.00000e+3	3.16228e+2	1.00000e+0	2.00000e-19	5.65685e+0
$(\rho_1/\rho_2)^{1/5}(R_1/R_2)$	7.09728e-3	1.31430e-2	1.58489e+1	1.00000e+1	1.00000e+0	3.31445e-8	2.00000e+0
$(I_1/I_2)^{1/5}$	7.09728e-3	1.31430e-2	1.58489e+1	1.00000e+1	1.00000e+0	3.31445e-8	2.00000e+0
M_1/M_2	1.00000e+0	1.00000e+0	1.00000e+6	1.00000e+3	1.00000e+2	1.00000e-10	8.00000e+0
$\sqrt{M_1/M_2}$	1.00000e+0	1.00000e+0	1.00000e+3	3.16228e+1	1.00000e+1	1.00000e-5	2.82843e+0
R_1/R_2	4.24356e-6	1.98031e-5	1.00000e+0	1.00000e+1	1.00000e-1	2.00000e-14	2.00000e+0
$\sqrt{R_1/R_2}$	2.05999e-3	4.45007e-3	1.00000e+0	3.16228e+0	3.16228e-1	1.41421e-7	1.41421e+0

5.2 Case 2: ($M_1 = M_2$, $R_1 = \frac{4}{3}r_s$)

The two identical masses cancel, reducing the DeGerlia inertial density ratio down to a ratio of radii.

$$k_d = \frac{M_1 R_2}{M_2 R_1 D_{\text{crit}}} = \frac{R_2}{R_1 D_{\text{crit}}}$$

Note, This reflects the k calculation for gravitational time dilation. However, in this case, instead of comparing with r_s , we are comparing with $\frac{4}{3}r_s$, which equates to a gravitational time dilation of $\text{GTD} = 5.00000e-1$.

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	1.00000e30	1.00000e30
R (m)	1.98031e3	1.00000e8

Calculate DTD for each System:

$$\begin{aligned} \text{DTD} &= \sqrt{k} \quad \text{where } k = M/(RD_{\text{crit}}) \\ k_1 &= \frac{1.00000e30 \text{ kg}}{(1.98031e3 \text{ m})(6.73295e26 \text{ kg/m})} = 7.50001e-1 \\ \text{DTD}_1 &= \sqrt{7.50001e-1} = 8.66026e-1 \\ k_2 &= \frac{1.00000e30 \text{ kg}}{(1.00000e8 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-5 \\ \text{DTD}_2 &= \sqrt{1.48523e-5} = 3.85387e-3 \\ \text{DTD}_1 &= 8.66026e-1 \\ \text{DTD}_2 &= 3.85387e-3 = 2.24716e2 \end{aligned}$$

Calculate GTD ratio from DTD:

$$\begin{aligned} \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= \frac{\sqrt{1 - \text{DTD}_1^2}}{\sqrt{1 - \text{DTD}_2^2}} = \frac{\sqrt{1 - (8.66026e-1)^2}}{\sqrt{1 - (3.85387e-3)^2}} \\ &= \frac{\sqrt{1 - 7.50001e-1}}{\sqrt{1 - 1.48523e-5}} \\ &= \frac{\sqrt{2.49999e-1}}{\sqrt{9.999999999851477e-1}} \\ &= \frac{4.99999e-1}{9.9999257381e-1} = 5.00003e-1 \\ &= 1 - 4.99997e-1 \end{aligned}$$

Verify by Calculating GTD from Schwarzschild Radius Ratio:

Schwarzschild radii:

$$\begin{aligned} r_{s1} = r_{s2} &= \frac{2GM}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e30 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523e3 \text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{1.48523e3 \text{ m}}{1.98031e3 \text{ m}} = 7.50000e-1 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{1.48523e3 \text{ m}}{1.00000e8 \text{ m}} = 1.48523e-5 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

$$\begin{aligned} \text{GTD}_{s1} &= \sqrt{1 - k_1} = \sqrt{2.50000e-1} = 5.00000e-1 = 1 - 5.00000e-1 \\ \text{GTD}_{s2} &= \sqrt{1 - k_2} = \sqrt{9.999999999851477e-1} = 9.9999257381e-1 \\ &= 1 - 7.42619e-6 \\ \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= \frac{5.00000e-1}{9.9999257381e-1} = 5.00004e-1 \\ &= 1 - 4.99996e-1 \end{aligned}$$

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\begin{aligned} \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= 1 - 4.99996e-1 \quad (\text{via } r_s) \\ \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= 1 - 4.99997e-1 \quad (\text{via } D_{\text{crit}}) \end{aligned}$$

5.3 Case 3: ($R_1 = R_2$)

When the two systems share the same radius, the radius terms cancel in k_d :

$$k_d = \frac{M_1 R_2}{M_2 R_1 D_{\text{crit}}} = \frac{M_1}{M_2 D_{\text{crit}}}$$

The Lorentz factor has the same form as Equation 4, with $k = v^2/c^2$. Since velocity is distance per unit time, v^2/c^2 is the square of a ratio of two lengths, with the shared time unit cancelling.

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	1.00000e30	1.00000e24
R (m)	1.00000e8	1.00000e8

Calculate DTD for each System:

$$\begin{aligned} \text{DTD} &= \sqrt{k} \quad \text{where } k = M/(RD_{\text{crit}}) \\ k_1 &= \frac{1.00000e30 \text{ kg}}{(1.00000e8 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-5 \\ \text{DTD}_1 &= \sqrt{1.48523e-5} = 3.85387e-3 \\ k_2 &= \frac{1.00000e24 \text{ kg}}{(1.00000e8 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-11 \\ \text{DTD}_2 &= \sqrt{1.48523e-11} = 3.85387e-6 \\ \text{DTD}_1 &= 3.85387e-3 \\ \text{DTD}_2 &= 3.85387e-6 = 1.00000e3 \end{aligned}$$

Calculate GTD ratio from DTD:

$$\begin{aligned} \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= \frac{\sqrt{1 - \text{DTD}_1^2}}{\sqrt{1 - \text{DTD}_2^2}} = \frac{\sqrt{1 - (3.85387e-3)^2}}{\sqrt{1 - (3.85387e-6)^2}} \\ &= \frac{\sqrt{1 - 1.48523e-5}}{\sqrt{1 - 1.48523e-11}} \\ &= \frac{\sqrt{9.999999999851477e-1}}{\sqrt{9.999999999999999e-1}} \\ &= \frac{9.9999257381e-1}{9.999999999999999e-1} = 9.9999257381e-1 \\ &= 1 - 7.42619e-6 \end{aligned}$$

Verify by Calculating GTD from Schwarzschild Radius Ratio:

Schwarzschild radii:

$$\begin{aligned} r_{s1} &= \frac{2GM_1}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e30 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523e3 \text{ m} \\ r_{s2} &= \frac{2GM_2}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e24 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523e-3 \text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{1.48523e3 \text{ m}}{1.00000e8 \text{ m}} = 1.48523e-5 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{1.48523e-3 \text{ m}}{1.00000e8 \text{ m}} = 1.48523e-11 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

$$\begin{aligned} \text{GTD}_{s1} &= \sqrt{1 - k_1} = \sqrt{9.999999999851477e-1} = 9.9999257381e-1 \\ &= 1 - 7.42619e-6 \\ \text{GTD}_{s2} &= \sqrt{1 - k_2} = \sqrt{9.999999999999999e-1} \\ &= 9.999999999999999e-1 = 1 - 7.42616e-12 \\ \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= \frac{9.9999257381e-1}{9.999999999999999e-1} \\ &= 9.9999257382e-1 = 1 - 7.42618e-6 \end{aligned}$$

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\begin{aligned} \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= 1 - 7.42618e-6 \quad (\text{via } r_s) \\ \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= 1 - 7.42619e-6 \quad (\text{via } D_{\text{crit}}) \end{aligned}$$

5.4 Case 4: ($\rho_1 = \rho_2$)

When the two systems share the same density, $M \propto R^3$, so $M_1/M_2 = (R_1/R_2)^3$. Substituting into k_d :

$$k_d = \frac{M_1 R_2}{M_2 R_1 D_{\text{crit}}} = \left(\frac{R_1}{R_2}\right)^3 \cdot \frac{R_2}{R_1 D_{\text{crit}}} = \frac{1}{D_{\text{crit}}} \left(\frac{R_1}{R_2}\right)^2$$

The DeGerlia inertial density ratio reduces to the square of the radius ratio; the linear scale factor follows as $\sqrt{k_d} = R_1/R_2$, recovering the isometric equivalence.

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	1.00000e30	1.00000e27
R (m)	1.00000e8	1.00000e7

Calculate DTD for each System:

$$\begin{aligned} \text{DTD} &= \sqrt{k} \quad \text{where } k = M/(RD_{\text{crit}}) \\ k_1 &= \frac{1.00000e30 \text{ kg}}{(1.00000e8 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-5 \\ \text{DTD}_1 &= \sqrt{1.48523e-5} = 3.85387e-3 \\ k_2 &= \frac{1.00000e27 \text{ kg}}{(1.00000e7 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-7 \\ \text{DTD}_2 &= \sqrt{1.48523e-7} = 3.85387e-4 \\ \frac{\text{DTD}_1}{\text{DTD}_2} &= \frac{3.85387e-3}{3.85387e-4} = 1.00000e1 \end{aligned}$$

Calculate GTD ratio from DTD:

$$\begin{aligned} \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= \frac{\sqrt{1 - \text{DTD}_1^2}}{\sqrt{1 - \text{DTD}_2^2}} = \frac{\sqrt{1 - (3.85387e-3)^2}}{\sqrt{1 - (3.85387e-4)^2}} \\ &= \frac{\sqrt{1 - 1.48523e-5}}{\sqrt{1 - 1.48523e-7}} \\ &= \frac{\sqrt{9.999851477e-1}}{\sqrt{9.99999851477e-1}} \\ &= \frac{9.9999257381e-1}{9.99999257383e-1} = 9.9999264807e-1 \\ &= 1 - 7.35193e-6 \end{aligned}$$

Verify by Calculating GTD from Schwarzschild Radius Ratio:

Schwarzschild radii:

$$\begin{aligned} r_{s1} &= \frac{2GM_1}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e30 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523e3 \text{ m} \\ r_{s2} &= \frac{2GM_2}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e27 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523 \text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{1.48523e3 \text{ m}}{1.00000e8 \text{ m}} = 1.48523e-5 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{1.48523 \text{ m}}{1.00000e7 \text{ m}} = 1.48523e-7 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

$$\begin{aligned} \text{GTD}_{s1} &= \sqrt{1 - k_1} = \sqrt{9.999851477e-1} = 9.9999257381e-1 \\ &= 1 - 7.42619e-6 \\ \text{GTD}_{s2} &= \sqrt{1 - k_2} = \sqrt{9.99999851477e-1} = 9.99999257384e-1 \\ &= 1 - 7.42616e-8 \\ \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= \frac{9.9999257381e-1}{9.99999257384e-1} \\ &= 9.9999264807e-1 = 1 - 7.35193e-6 \end{aligned}$$

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\begin{aligned} \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= 1 - 7.35193e-6 \quad (\text{via } r_s) \\ \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= 1 - 7.35193e-6 \quad (\text{via } D_{\text{crit}}) \end{aligned}$$

5.5 Case 5: ($I_1 = I_2$)

When the two systems share the same moment of inertia, $M_1 R_1^2 = M_2 R_2^2$, so $M_1/M_2 = (R_2/R_1)^2$. Substituting into k_d :

$$k_d = \frac{M_1 R_2}{M_2 R_1 D_{\text{crit}}} = \left(\frac{R_2}{R_1}\right)^2 \cdot \frac{R_2}{R_1 D_{\text{crit}}} = \frac{1}{D_{\text{crit}}} \left(\frac{R_2}{R_1}\right)^3$$

The DeGerlia inertial density ratio reduces to the cube of the radius ratio.

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	1.00000e32	1.00000e30
R (m)	1.00000e7	1.00000e8

Calculate DTD for each System:

$$\begin{aligned} \text{DTD} &= \sqrt{k} \quad \text{where } k = M/(RD_{\text{crit}}) \\ k_1 &= \frac{1.00000e32 \text{ kg}}{(1.00000e7 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-2 \\ \text{DTD}_1 &= \sqrt{1.48523e-2} = 1.21870e-1 \\ k_2 &= \frac{1.00000e30 \text{ kg}}{(1.00000e8 \text{ m})(6.73295e26 \text{ kg/m})} = 1.48523e-5 \\ \text{DTD}_2 &= \sqrt{1.48523e-5} = 3.85387e-3 \\ \frac{\text{DTD}_1}{\text{DTD}_2} &= \frac{1.21870e-1}{3.85387e-3} = 3.16228e1 \end{aligned}$$

Calculate GTD ratio from DTD:

$$\begin{aligned} \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= \frac{\sqrt{1 - \text{DTD}_1^2}}{\sqrt{1 - \text{DTD}_2^2}} = \frac{\sqrt{1 - (1.21870e-1)^2}}{\sqrt{1 - (3.85387e-3)^2}} \\ &= \frac{\sqrt{1 - 1.48523e-2}}{\sqrt{1 - 1.48523e-5}} \\ &= \frac{\sqrt{9.851477e-1}}{\sqrt{9.999851477e-1}} \\ &= \frac{9.9254605e-1}{9.9999257381e-1} = 9.9255342e-1 \\ &= 1 - 7.44658e-3 \end{aligned}$$

Verify by Calculating GTD from Schwarzschild Radius Ratio:

Schwarzschild radii:

$$\begin{aligned} r_{s1} &= \frac{2GM_1}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e32 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523e5 \text{ m} \\ r_{s2} &= \frac{2GM_2}{c^2} = \frac{2(6.67430e-11 \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000e30 \text{ kg})}{(2.99792e8 \text{ m s}^{-1})^2} \\ &= 1.48523e3 \text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{1.48523e5 \text{ m}}{1.00000e7 \text{ m}} = 1.48523e-2 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{1.48523e3 \text{ m}}{1.00000e8 \text{ m}} = 1.48523e-5 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

$$\begin{aligned} \text{GTD}_{s1} &= \sqrt{1 - k_1} = \sqrt{9.851477e-1} = 9.9254606e-1 = 1 - 7.45394e-3 \\ \text{GTD}_{s2} &= \sqrt{1 - k_2} = \sqrt{9.999851477e-1} = 9.9999257381e-1 \\ &= 1 - 7.42619e-6 \\ \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= \frac{9.9254606e-1}{9.9999257381e-1} = 9.9255343e-1 \\ &= 1 - 7.44657e-3 \end{aligned}$$

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\begin{aligned} \frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} &= 1 - 7.44657e-3 \quad (\text{via } r_s) \\ \frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} &= 1 - 7.44658e-3 \quad (\text{via } D_{\text{crit}}) \end{aligned}$$

5.6 Case 6: (General, unconstrained)

When no constraint is applied between the systems, k_d stays in its full form:

$$k_d = \frac{M_1 R_2}{M_2 R_1 D_{\text{crit}}}$$

Both mass and radius must be specified independently for each system.

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	1.00000e20	1.00000e30
R (m)	2.00000e-7	1.00000e7

Calculate DTD for each System:

$$k_1 = \frac{1.00000\text{e}20\text{kg}}{(2.00000\text{e}-7\text{m})(6.73295\text{e}26\text{kg/m})} = 7.42617\text{e}-1$$

$$\text{DTD}_1 = \sqrt{7.42617\text{e}-1} = 8.61752\text{e}-1$$

$$k_2 = \frac{1.00000\text{e}30\text{ kg}}{(1.00000\text{e}7\text{ m})(6.73295\text{e}26\text{ kg/m})} = 1.48523\text{e}-4$$

$$\text{DTD}_2 = \sqrt{1.48523\text{e}-4} = 1.21870\text{e}-2$$

DTD ₁	8.61752e-1	7.07107e-1
------------------	------------	------------

$$\overline{\text{DTD}_2} = \overline{1.21870\text{e}-2} = 7.07107\text{e1}$$

Calculate GTD ratio from DTD:

$$\begin{aligned} \frac{\text{GTD}_{\text{d1}}}{\text{GTD}_{\text{d2}}} &= \frac{\sqrt{1 - \text{DTD}_1^2}}{\sqrt{1 - \text{DTD}_2^2}} = \frac{\sqrt{1 - (8.61752\text{e-}1)^2}}{\sqrt{1 - (1.21870\text{e-}2)^2}} \\ &= \frac{\sqrt{1 - 7.42617\text{e-}1}}{\sqrt{1 - 1.48523\text{e-}4}} \\ &= \frac{\sqrt{2.57383\text{e-}1}}{\sqrt{9.99851477\text{e-}1}} \\ &= \frac{5.07330\text{e-}1}{9.999257356\text{e-}1} = 5.07367\text{e-}1 \\ &= 1 - 4.92633\text{e-}1 \end{aligned}$$

Verify by Calculating GTD from Schwarzschild Radius Ratio:

Schwarzschild radii:

$$\begin{aligned} r_{s1} &= \frac{2GM_1}{c^2} = \frac{2(6.67430 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000 \times 20 \text{ kg})}{(2.99792 \times 8 \text{ m s}^{-1})^2} \\ &= 1.48523 \times 10^{-7} \text{ m} \\ r_{s2} &= \frac{2GM_2}{c^2} = \frac{2(6.67430 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2})(1.00000 \times 30 \text{ kg})}{(2.99792 \times 8 \text{ m s}^{-1})^2} \\ &= 1.48523 \times 10^{-7} \text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{1.48523\text{e-}7\text{m}}{2.00000\text{e-}7\text{m}} = 7.42616\text{e-}1 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{1.48523\text{e}3\text{m}}{1.00000\text{e}7\text{m}} = 1.48523\text{e-}4 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

$$\begin{aligned} \text{GTD}_{\text{s1}} &= \sqrt{1 - k_1} = \sqrt{2.57384\text{e-}1} = 5.07330\text{e-}1 = 1 - 4.92670\text{e-}1 \\ \text{GTD}_{\text{s2}} &= \sqrt{1 - k_2} = \sqrt{9.99851477\text{e-}1} = 9.999257356\text{e-}1 \\ &= 1 - 7.42644\text{e-}5 \\ \frac{\text{GTD}_{\text{s1}}}{\text{GTD}_{\text{s2}}} &= \frac{5.07330\text{e-}1}{9.999257356\text{e-}1} = 5.07368\text{e-}1 \\ &= 1 - 4.92632\text{e-}1 \end{aligned}$$

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} = 1 - 4.92632\text{e-}1 \quad (\text{via } r_s)$$

$$\frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} = 1 - 4.92633\text{e-}1 \quad (\text{via } D_{\text{crit}})$$

5.7 Case 7: (Classical)

System properties:

	System 1 (S_1)	System 2 (S_2)
M (kg)	8.00000	1.00000
R (m)	2.00000	1.00000

Calculate DTD for each System:

$$\begin{aligned} \text{DTD} &= \sqrt{k} \quad \text{where} \quad k = M/(RD_{\text{crit}}) \\ k_1 &= \frac{8.00000\text{kg}}{(2.00000\text{m})(6.73295\text{e}26\text{g/m})} = 5.94093\text{e}-27 \\ \text{DTD}_1 &= \sqrt{5.94093\text{e}-27} = 7.70774\text{e}-14 \end{aligned}$$

$$k_2 = \frac{1.00000\text{kg}}{(1.00000\text{m})(6.73295\text{e}26\text{kg/m})} = 1.48523\text{e-}27$$

$$\text{DTD}_2 = \sqrt{1.48523\text{e}-27} = 3.85387\text{e}-14$$

$$\frac{\text{DTD}_1}{\text{}} = \frac{7.70774\text{e-}14}{\text{}} = 2.00000$$

$$\overline{\text{DTD}_2} = 3.85387\text{e-}14 = 2.00000$$

Calculate GTD ratio from DTD:

[illegible]

Verify by Calculating GTD from Schwarzschild Radius Ratio:

Schwarzschild radii:

$$\begin{aligned} r_{s1} &= \frac{2GM_1}{c^2} = \frac{2(6.67430\text{e-}11\text{ m}^3\text{ kg}^{-1}\text{ s}^{-2})(8.00000\text{ kg})}{(2.99792\text{e}8\text{ m s}^{-1})^2} \\ &= 1.18819\text{e-}26\text{ m} \\ r_{s2} &= \frac{2GM_2}{c^2} = \frac{2(6.67430\text{e-}11\text{ m}^3\text{ kg}^{-1}\text{ s}^{-2})(1.00000\text{ kg})}{(2.99792\text{e}8\text{ m s}^{-1})^2} \\ &= 1.48523\text{e-}27\text{ m} \end{aligned}$$

Calculate k values from the r_s/R ratio:

$$\begin{aligned} k_1 &= \frac{r_{s1}}{R_1} = \frac{1.18819 \text{e-}26 \text{ m}}{2.00000 \text{ m}} = 5.94093 \text{e-}27 \\ k_2 &= \frac{r_{s2}}{R_2} = \frac{1.48523 \text{e-}27 \text{ m}}{1.00000 \text{ m}} = 1.48523 \text{e-}27 \end{aligned}$$

Calculate GTD via Schwarzschild radius ratio:

[illegible]

Time-dilation ratio between System 1 and System 2, computed two ways:

$$\frac{\text{GTD}_{s1}}{\text{GTD}_{s2}} = 1 - 2.22785\text{e-}27 \quad (\text{via } r_s)$$

$$\frac{\text{GTD}_{d1}}{\text{GTD}_{d2}} = 1 - 2.22785\text{e-}27 \quad (\text{via } D_{\text{crit}})$$

6 Analysis

The reformulation is meant to carry three points, and nothing more ambitious. First, it is a pace-of-time framework that is sim-

ply easier to work with: expressed in DTD and compactness, comparing two systems is a direct ratio of geometries, with no benchmark, no subtraction-from-one, and — because D_{crit} cancels — no dependence on G . One returns to GTD, the $\sqrt{1-k}$ form, only at the final step, to convert relative pace into relative clock [1] time. Second, general relativity is always two-system; we merely masquerade it as one. Conventional gravitational time dilation looks like a single-system calculation only because it silently fixes [2] the second system — the same mass at its Schwarzschild radius, where $\text{DTD} = 1$. Make that benchmark explicit and the two-system structure, present all along, is plain. Third, treating it [3] as one system is a disservice to the broader mathematics: the collapse to a single system rests on presuming that some property of the hidden second system is a fixed, universal given, a presumption that obscures the real structure and perpetuates a misconception that dilutes the core principle. The honest formulation keeps [4] both systems in view and recognizes the single-system shortcut as [5] a convenience, not as the underlying physics.

7 Conclusions

8 Hypotheses of Explanation

9 Further Exploration

Acknowledgments

This work was conducted independently and received no external funding.

Conflicts of Interest

The author declares no competing interests.

Data Availability

The complete source materials for this paper, including all $\text{\LaTeX}$ source files, derivation documents, and revision history, are pub-

licly available at: https://github.com/denverdata/academic_InertialRelativity_STE_cases.

References

- [1] Artur Alho, José Natário, Paolo Pani, and Guilherme Raposo. Compactness bounds in general relativity. *Physical Review D*, 106:L041502, 2022. doi: 10.1103/PhysRevD.106.L041502.
- [2] T. D. DeGerlia. The universe of light - exploring our universe through the lens of the law of space-time equivalence. DEC, 2025a. DOI: 10.55277/ResearchHub.pi9yjshm.
- [3] T. D. DeGerlia. Introducing inertial density and characterizing the schwarzschild condition as a constant threshold of mass over radius (m/r). DEC, 2025b. DOI: 10.55277/research-hub.0e0cgoor.
- [4] A. Einstein. Zur elektrodynamik bewegter körper. *Annalen der Physik*, 17:891–921, 1905.
- [5] A. Einstein. Die feldgleichungen der gravitation. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin)*, pages 844–847, 1915.
- [6] H. Goldstein, C. Poole, and J. Safko. *Classical Mechanics*. Addison-Wesley, 3rd edition, 2002.
- [7] R. P. Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics. *Physical Review Letters*, 11:237–238, 1963.
- [8] K. Schwarzschild. Über das gravitationsfeld eines massenpunktes nach der einsteinschen theorie. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin)*, pages 189–196, 1916.
- [9] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor. Codata recommended values of the fundamental physical constants: 2018. *Reviews of Modern Physics*, 93(2):025010, 2021.