

EQUIVARIANT CLASSES OF ORBITS IN $\mathrm{GL}(2)$ -REPRESENTATIONS

ANAND DEOPURKAR

ABSTRACT. We compute equivariant fundamental classes of orbits in $\mathrm{GL}(2)$ -representations. As applications, we find degrees of the orbit closures corresponding to elliptic fibrations and self-maps of the projective line.

1. INTRODUCTION

If we fix a hypersurface in projective space, how complicated is the set of all hypersurfaces obtained from the fixed one by changes of coordinates? Similarly, if we fix a self-map of the projective space, how complicated is the set of all self-maps obtained from the fixed one by changes of coordinates? These questions, and many others, generalise as follows. Given a representation W of an algebraic group G , how complicated is the G -orbit of a fixed $w \in W$? One measure of complexity is the degree of the orbit closure in $\mathbf{P}W$. A more refined measure is the G -equivariant fundamental class. Our main theorem (Theorem 1.3) completely describes the equivariant fundamental classes (and hence degrees) of orbits in representations of $G = \mathrm{GL}(2)$. The case where W is irreducible was already known. The new contribution is treating reducible W ; this presents new challenges, but also has new applications. The case of $G = \mathrm{GL}(1)$, or more generally, any torus, is straightforward. We treat it in Appendix A.

The question of finding equivariant classes of orbit closures has been well studied, especially in cases where the orbits have a geometric interpretation. For the $\mathrm{GL}(2)$ representation $\mathrm{Sym}^n \mathbf{C}^2$, where the orbits represent divisors of degree n on \mathbf{P}^1 modulo changes of coordinates, the degree of the orbit closure was computed by Enriques–Fano [8] for the generic case and Aluffi–Faber in general [1]. The equivariant class was computed by Lee–Patel–Tseng [14, Appendix B]. For the $\mathrm{GL}(3)$ representation $\mathrm{Sym}^n \mathbf{C}^3$, where the orbits represent plane curves of degree n , the degree of the orbit closure was computed by Aluffi–Faber [2,3]. For the $\mathrm{GL}(4)$ representation $\mathrm{Sym}^3 \mathbf{C}^4$, where the orbits represent cubic surfaces, the equivariant class of a generic orbit closure was computed in [7]. Local analogues of equivariant orbit classes are Thom polynomials, which have been studied by Buch, Fehér, Rimányi, and Weber among others [10,11,19]. In all these cases, the equivariant class yields a counting formula—the equivariant orbit/Thom class of w gives the number of times w appears, up to isomorphism, in a given family. We expect the equivariant class to reflect the geometry of w . This is indeed the case for divisors on \mathbf{P}^1 , where the class depends on the multiplicities in the divisor, and for curves in \mathbf{P}^2 , where the class depends on the singularities and flexes of the curve.

As a direct application of the main theorem, we compute the degrees of orbit closures in two (reducible) representations of geometric significance. The first is the $\mathrm{GL}(2)$ -representation $\mathrm{Sym}^{4n}(\mathbf{C}^2) \oplus \mathrm{Sym}^{6n}(\mathbf{C}^2)$, where the orbits represent isomorphism classes of elliptic fibrations over \mathbf{P}^1 . In this case, the degree depends on the Kodaira types of the singular fibers. The second is the $\mathrm{GL}(2)$ -representation $\mathrm{Hom}(\mathbf{C}^2, \mathrm{Sym}^n \mathbf{C}^2)$, where most orbits represent isomorphism classes of

Date: May 19, 2024.

self-maps $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree n . (The space of degree n maps $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ and its quotient by changes of coordinates are important objects of study in complex dynamics, where they are usually denoted by Rat_n and \mathcal{M}_n [15, 18, 20, 21]). In this case, the degree is (surprisingly) independent of the orbit.

We first give these two applications in Section 1.1 and Section 1.2, respectively, before stating the main theorem in Section 1.3.

1.1. Elliptic fibrations. Fix a positive integer n , and let $W = \text{Sym}^{4n}(\mathbf{C}^2) \oplus \text{Sym}^{6n}(\mathbf{C}^2)$. A non-zero $(A, B) \in W$ determines an elliptic fibration over \mathbf{P}^1 defined by the Weierstrass equation

$$y^2 = x^3 + Ax + B.$$

The $\text{GL}(2)$ -orbits in W thus represent Weierstrass elliptic fibrations over \mathbf{P}^1 , up to isomorphism.

Let h be the class of the Weil divisor $\mathcal{O}(1)$ on the weighted projective space $\mathbf{P}W = (W - 0)/\mathbf{G}_m$, where \mathbf{G}_m acts by weight 2 on $\text{Sym}^{4n}(\mathbf{C}^2)$ and by weight 3 on $\text{Sym}^{6n}(\mathbf{C}^2)$. Given $(A, B) \in W$ and $u \in \mathbf{P}^1$, let $\text{ord}(A)_u$ and $\text{ord}(B)_u$ be the orders of vanishing of A and B at u . Set

$$c(u) = \min \left(\frac{1}{2} \text{ord}(A)_u, \frac{1}{3} \text{ord}(B)_u \right).$$

Theorem 1.1. *Fix a non-zero $w = (A, B) \in W = \text{Sym}^{4n}(\mathbf{C}^2) \oplus \text{Sym}^{6n}(\mathbf{C}^2)$, and let $\pi: E \rightarrow \mathbf{P}^1$ be the Weierstrass fibration defined by w . Let $D \subset \mathbf{P}^1$ be a finite set such that π is smooth on $\mathbf{P}^1 - D$. Let $\Gamma \subset \text{GL}(2)$ be the stabiliser of $w \in W$ and let $\text{Orb}([w])$ be the closure of the $\text{PGL}(2)$ -orbit of $[w] \in \mathbf{P}W$. Then*

$$|\Gamma|[\text{Orb}([w])] = 2^{4n+3} 3^{6n+1} \cdot n \cdot \left(4n^3 - \sum_{u \in D} c(u)^2 (3n - c(u)) \right) h^{10n-2}.$$

If π is a minimal Weierstrass fibration as in [16, III.3], then $c(u) < 2$ and $c(u)$ determines the Kodaira fiber type over u (see [16, IV.3.1]). See Table 1 for the Kodaira types and their contribution to the formula above. The main theorem in fact gives the equivariant class, of which the degree is a particular specialisation. See Section 7.1 for the proof.

$c(u)$	Type	Description	Contribution to the degree $c(u)^2(3n - c(u))$
0	I_N	Smooth elliptic curve, nodal rational curve, or cycle of smooth rational curves	0
1	I_N^*	\tilde{D}_{4+N} -configuration of rational curves	$3n - 1$
1/3	II	Cuspidal rational curve	$1/27 \cdot (9n - 1)$
1/2	III	Two tangent rational curves	$1/8 \cdot (6n - 1)$
2/3	IV	Three concurrent rational curves	$4/27 \cdot (9n - 2)$
4/3	IV^*	\tilde{E}_6 -configuration of rational curves	$16/27 \cdot (9n - 4)$
3/2	III^*	\tilde{E}_7 -configuration of rational curves	$27/8 \cdot (2n - 1)$
5/3	II^*	\tilde{E}_8 -configuration of rational curves	$25/27 \cdot (9n - 5)$

TABLE 1. Contributions from the singular fibers in a minimal Weierstrass fibration $y^2 = x^3 + Ax + B$ towards the degree of the orbit closure of $(A, B) \in \mathbf{P}(\text{Sym}^{4n}(\mathbf{C}^2) \oplus \text{Sym}^{6n}(\mathbf{C}^2))$.

1.2. Rational self maps. Fix a positive integer n and set $W = \mathrm{Hom}(\mathbf{C}^2, \mathrm{Sym}^n \mathbf{C}^2)$. An element $f \in W$ is equivalent to a map

$$(1) \quad \mathbf{C}^2 \otimes \mathcal{O}_{\mathbf{P}^1} \rightarrow \mathcal{O}_{\mathbf{P}^1}(n).$$

For f in a Zariski open subset, the map (1) is surjective, and hence defines a map $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree n . Conversely, every map $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree n arises from an $f \in W$, which is unique up to a scalar. Thus, most $\mathrm{GL}(2)$ -orbits in W represent maps $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree n modulo changes of coordinates.

Theorem 1.2. *Suppose $f \in \mathrm{Hom}(\mathbf{C}^2, \mathrm{Sym}^n \mathbf{C}^2)$ defines a map $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree n . Let $\bar{\Gamma} \subset \mathrm{PGL}(2)$ be the stabiliser of $[f] \in \mathbf{P}\mathrm{Hom}(\mathbf{C}^2, \mathrm{Sym}^n \mathbf{C}^2)$, and $\mathrm{Orb}([f])$ the closure of the $\mathrm{PGL}(2)$ -orbit of $[f]$. Then*

$$|\bar{\Gamma}| \cdot \deg(\mathrm{Orb}([f])) = n(n+1)(n-1).$$

Again, the main theorem gives the equivariant class, of which the degree is a particular specialisation. See Section 7.2 for the proof.

We highlight that Theorem 1.2 does not hold for all $f \in \mathrm{Hom}(\mathbf{C}^2, \mathrm{Sym}^n \mathbf{C}^2)$. For the f whose associated map (1) is not surjective— f with base-points—the stabiliser-weighted degree can be different. It is remarkable that for the f without base-points, it is constant. This is in contrast to the case of divisors on \mathbf{P}^1 , where the multiplicities in the divisor matter.

1.3. Main theorem. Fix a 2-dimensional vector space V over an algebraically closed field \mathbf{k} of characteristic 0. Fix a finite dimensional $\mathrm{GL} V$ representation

$$W = W_1 \oplus \cdots \oplus W_n, \text{ where } W_i = \mathrm{Sym}^{a_i - b_i} V \otimes \det V^{b_i}.$$

Set $d_i = a_i + b_i$, and assume that $d_i > 0$ for all i . Fix a maximal torus $T \subset \mathrm{GL} V$. We then have an isomorphism between the equivariant (rational) Chow ring $A_{\mathrm{GL} V}$ and the symmetric polynomials in $A_T = \mathbf{Q}[v_1, v_2]$.

We must now introduce some notation. Fix a non-zero $w = (w_1, \dots, w_n) \in W$, and write $w_i = f_i \otimes \delta^{b_i}$ for some $f_i \in \mathrm{Sym}^{a_i - b_i} V$ and $\delta \in \det V$. Given $u \in \mathbf{P}^1$, let r_i^u be the order of vanishing of f_i at u . Let $\Lambda^u \subset \mathbf{R}^2$ be the convex hull of the union of the shifted quadrants

$$\frac{1}{d_i}(r_i^u + b_i, b_i) + \mathbf{R}_{\geq 0}^2.$$

Let $\lambda^u(0), \dots, \lambda^u(k^u)$ be the vertices of Λ^u arranged from the bottom right to the top left. For a $p \in \mathbf{R}^2$, use p_1 and p_2 to denote the first and the second coordinates. Set

$$b = \min(b_i/d_i \mid w_i \neq 0), \quad r_{\mathrm{gen}}^u = \lambda^u(0)_1 - b, \quad r^u = \min((r_i^u + b_i)/d_i), \text{ and} \\ s^u = \begin{cases} 1 - \frac{\lambda^u(0)_1 - \lambda^u(1)_1}{\lambda^u(0)_2 - \lambda^u(1)_2}, & \text{if } k^u \geq 1, \\ 1, & \text{otherwise.} \end{cases}$$

For $j = 1, \dots, k^u$, let $\eta^u(j)$ and $\zeta^u(j)$ be the smallest integral normal vectors to the rays of Λ^u at the vertex $\lambda^u(j)$. Set $N^u(j) = \det(\eta^u(j), \zeta^u(j))$. Let $A \subset \mathbf{P}^1$ be a finite set that includes the common zero locus of $\{f_i \mid b_i/d_i = b\}$.

Given $F \in \mathbf{Q}(v_1, v_2)$, denote by F_{sym} its symmetrisation

$$F_{\mathrm{sym}} = F(v_1, v_2) + F(v_2, v_1).$$

Let $N = \dim W$ and observe that in $A_T = \mathbf{Q}[v_1, v_2]$, we have the top Chern class

$$c_N(W) = \prod_{i=1}^n \prod_{j=0}^{a_i-b_i} ((b_i + j)v_1 + (a_i - j)v_2).$$

Let $\Gamma \subset \mathrm{GL}(V)$ be the stabiliser of $w \in W$; assume that it is finite.

Theorem 1.3. *In the notation above, the $\mathrm{GL} V$ -equivariant class of the orbit closure of $w \in W$ in $A_{\mathrm{GL} V}(W) \subset \mathbf{Q}[v_1, v_2]$ is given by*

$$(2) \quad |\Gamma|[\mathrm{Orb}(w)] = c_N(W) \cdot \left(F_{\mathrm{sym}} + \sum_{u \in A} G_{\mathrm{sym}}^u + \sum_{u \in A} \sum_{j=1}^{k^u} H^u(j)_{\mathrm{sym}} \right),$$

where

$$\begin{aligned} F &= 2((1-b)v_1 + bv_2)^{-1}(v_1 - v_2)^{-3} \\ &\quad - (2b-1)((1-b)v_1 + v_2)^{-2}(v_1 - v_2)^{-2} \\ G^u &= ((1-r^u)v_1 + r^u v_2)^{-1}(v_1 - v_2)^{-3} \\ &\quad - s^u((1-b)v_1 + bv_2)^{-1}(v_1 - v_2)^{-3} \\ &\quad - r_{\mathrm{gen}}^u((1-b)v_1 + v_2)^{-2}(v_1 - v_2)^{-2}, \text{ and} \\ H^u(j) &= |N^u(j)|\eta^u(j)_1^{-1}\zeta^u(j)_1^{-1}((1-\lambda^u(j)_2)v_1 + \lambda^u(j)_2 v_2)^{-1}(v_1 - v_2)^{-3}. \end{aligned}$$

Note that in the sum of $H^u(j)_{\mathrm{sym}}$, the bottom right vertex ($j = 0$) is omitted.

Remark 1.4. It is not obvious that the expression in Theorem 1.3 is a polynomial. But it must be, as a consequence of the theorem.

1.4. Negative or mixed weights. Our main theorem applies to representations W whose direct summands have positive weights d_i . The theorem can also be used for W whose direct summands have negative weights by dualising or by twisting by a large negative n as described in Section 4.

The cases where W has summands of weight 0 or some summands of positive weights and some of negative weights are a bit strange. In these cases, a generic $w \in W$ does not contain the origin in its orbit closure. Therefore, its equivariant class of a generic orbit closure is 0, as can be seen by pulling back to the equivariant Chow ring of the origin.

1.5. Ideas in the proof. Let \mathbf{PW} be the weighted projective space $(W - 0)/\mathbf{G}_m$ for the central $\mathbf{G}_m \subset \mathrm{GL}(2)$. Given a $w \in W$, the key idea is to find a complete orbit parametrisation for $\mathrm{Orb}([w])$, namely a proper $\mathrm{PGL}(2)$ -variety X and an equivariant finite map $X \rightarrow \mathbf{PW}$ whose image is $\mathrm{Orb}([w])$. Then the class of $\mathrm{Orb}([w])$ is the push-forward of $[X]$, up to a constant factor. The push-forward also gives $\mathrm{GL}(2)$ -equivariant class of $\mathrm{Orb}(w)$ (see Proposition 3.4).

To find X , we start with $M = \mathbf{P} \mathrm{Hom}(\mathbf{k}^2, \mathbf{k}^2)$, and the rational map $M \dashrightarrow \mathbf{PW}$ given by $m \mapsto mw$. We find an explicit resolution $\widetilde{M} \rightarrow \mathbf{PW}$, which serves as our complete orbit parametrisation. We then compute the push-forward as an integral on \widetilde{M} using Atiyah–Bott localisation.

The resolution $\widetilde{M} \rightarrow M$ is a weighted blow-up. It is much more convenient to take the weighted blow-up in a stacky sense. The stacky blow-up is smooth and maps to the weighted projective stack \mathcal{PW} . We can then write the push-forward as an integral and evaluate it using localisation. The stacky blow-up is toroidal, and is completely described by the combinatorial data of the Newton polygons Λ^u .

1.6. Conventions and organisation. We work over an algebraically closed field \mathbf{k} of characteristic 0. A stack means an algebraic stack over \mathbf{k} . All schemes and stacks are of finite type over \mathbf{k} . Given a vector space/bundle V , the projectivisation $\mathbf{P}V$ refers to the space of one-dimensional sub-spaces/bundles of V , consistent with the convention in [12] and $\mathcal{O}_{\mathbf{P}V}(-1)$ denotes the universal sub-bundle. All Chow groups are with rational coefficients.

In Section 2, we recall stacky weighted blow-ups in preparation for our main construction. In Section 3, we describe how to find the equivariant class of an orbit using a complete parametrisation. Both of these sections are general (not specific to $\mathrm{GL}(2)$). In Section 4, we observe that the main theorem is invariant under a twist operation, which allows some simplification. In Section 5, we construct a complete parametrisation of a $\mathrm{GL}(2)$ -orbit using a stacky blow-up. In Section 6, we evaluate the equivariant orbit class using localisation. In Section 7, we deduce the applications to elliptic fibrations and rational self maps. In Appendix A, we explain the case of G a torus.

Acknowledgements. The project arose from conversations with Anand Patel, to whom I am deeply grateful. I thank Ming Hao Quek for sharing his expertise on stacky blow-ups. I was supported by the grant DE180101360 from the Australian Research Council.

2. RATIONAL NEWTON POLYHEDRA AND WEIGHTED BLOW-UPS

The material in this section should be well-known to experts (see, for example, [17, § 2]).

Set $M = \mathbf{Z}^n$ and $N = \mathrm{Hom}(M, \mathbf{Z})$. Let $M_{\geq 0}$ be the set of vectors with non-negative coordinates in $M \otimes \mathbf{R} = \mathbf{R}^n$, and similarly for $N_{\geq 0}$. A *rational Newton polyhedron* is a closed convex polyhedron $\Lambda \subset M$ whose recession cone is $M_{\geq 0}$ and whose vertices have rational coordinates. Such a Λ gives a fan Λ^\perp in $N \otimes \mathbf{R}$ supported on $N_{\geq 0}$, called the *normal fan* of Λ . There is an inclusion reversing bijection between the faces of Λ and the cones of Λ^\perp . To a face F of Λ , we associate the cone F^\perp of Λ^\perp defined by

$$F^\perp = \{f \in N \mid f \text{ is constant on } F \text{ and this constant is the minimum of } f \text{ on } \Lambda\}.$$

Since the recession cone of Λ is $M_{\geq 0}$, and f achieves a minimum on Λ , it must lie in $N_{\geq 0}$.

Let F be a maximal face of Λ , that is, of dimension $(n-1)$. Then F^\perp is a ray. For every F , choose a non-zero vector $\beta_F \in F^\perp$ with integer coordinates. Let r be the number of maximal faces of Λ . Then the collection $\{\beta_F\}$ gives a homomorphism $\beta: \mathbf{Z}^r \rightarrow N$ with finite cokernel. Let $\mathcal{X}_{\Lambda, \beta}$ be the toric stack defined by the data $(N, \Lambda^\perp, \beta)$ in the sense of [4]. It comes with a canonical map $\mathcal{X}_{\Lambda, \beta} \rightarrow \mathbf{A}^n$, which we call the stacky blow-up of \mathbf{A}^n defined by (Λ, β) .

Let us describe $\mathcal{X}_{\Lambda, \beta} \rightarrow \mathbf{A}^n$ in charts, following [4, Proposition 4.3]. Assume that Λ is simplicial, that is, every vertex of Λ has exactly n incident rays. Let v be a vertex of Λ . Denote the rays incident to v by R_1, \dots, R_n and the maximal faces incident to v by F_1, \dots, F_n such that R_i is the only ray not contained in F_i . Set $\beta_i = \beta_{F_i}$ and let $r_i \in R_i$ be the unique vector such that $\langle \beta_i, r_i \rangle = 1$. Then r_1, \dots, r_n is a basis of $M \otimes \mathbf{Q}$ dual to the basis β_1, \dots, β_n of $N \otimes \mathbf{Q}$. Let $M_v \supset M$ be the dual lattice of the sub-lattice of N spanned by β_1, \dots, β_n . Then M_v/M is a finite abelian group. Set $\mu_v = \mathrm{Hom}(M_v/M, \mathbf{G}_m)$. The chart of $\mathcal{X}_{\Lambda, \beta}$ defined by v is

$$(3) \quad [\mathrm{Spec} \mathbf{k}[u_1, \dots, u_n]/\mu_v],$$

with the action given as follows. A $\zeta \in \mu_v$ acts by

$$\zeta: u_i \mapsto \zeta(r_i)u_i.$$

In particular, note that $\mathcal{X}_{\Lambda, \beta}$ is a smooth Deligne–Mumford stack.

Let e_1, \dots, e_n be the standard basis vectors in M . In the chart above, the map to $\mathbf{A}^n = \text{Spec } \mathbf{k}[x_1, \dots, x_n]$ is defined by

$$(4) \quad x_i \mapsto u_1^{\langle \beta_1, e_i \rangle} \dots u_n^{\langle \beta_n, e_i \rangle}.$$

Note that $\zeta \in \mu_v$ multiplies the image of x_i by $\zeta(e)$ where

$$e = r_1 \langle \beta_1, e_i \rangle + \dots + r_n \langle \beta_n, e_i \rangle.$$

Since r_1, \dots, r_n and β_1, \dots, β_n are dual bases, we see that $e = e_i \in M$ and hence $\zeta(e) = 1$. So the map (4) is indeed μ_v -invariant. Write $r_i = (a_1, \dots, a_n)$ in standard coordinates with $a_i \in \mathbf{Q}$. Informally, it is helpful to think of u_i as $x_1^{a_1} \dots x_n^{a_n}$.

Let X_Λ be the toric variety associated to (N, Λ^\perp) . Then we have a map $\mathcal{X}_{\Lambda, \beta} \rightarrow X_\Lambda$ which is the coarse space map [4, Proposition 3.7].

Remark 2.1. Let $(N, \Lambda^\perp, \beta)$ be the stacky fan given by a rational Newton polyhedron as above. Let r be the number of rays of Λ^\perp . In [4], the stack associated to $(N, \Lambda^\perp, \beta)$ is defined as the quotient of an open subset $Z \subset \mathbf{A}^r$ by the action of a group G that acts on \mathbf{A}^r through a homomorphism $G \rightarrow \mathbf{G}_m^r$. In our case, N is a free \mathbf{Z} -module. From the construction of $G \rightarrow \mathbf{G}_m^r$ in [4, § 2], it follows that $G \rightarrow \mathbf{G}_m^r$ is injective. Let $\mathcal{X} = [Z/G]$ and $\overline{\mathcal{X}} = [Z/\mathbf{G}_m^r]$. It is easy to see that we have the pull-back diagram

$$\begin{array}{ccc} \mathcal{X} & \longrightarrow & \overline{\mathcal{X}} \\ \downarrow & & \downarrow \\ \mathbf{A}^n & \longrightarrow & [\mathbf{A}^n / \mathbf{G}_m^r]. \end{array}$$

Given Λ , we use two natural choices of β . For the first, denoted by β^{can} , we let β_F be the shortest vector with integer coordinates on the ray F^\perp . For the second, denoted by β^{res} , we let β_F be the shortest vector with integer coordinates on the ray F^\perp such that the value of β_F on F is an integer. Then we have a map

$$\mathcal{X}_{\Lambda, \beta^{\text{res}}} \rightarrow \mathcal{X}_{\Lambda, \beta^{\text{can}}},$$

which is a sequence of root stacks along the divisors defined by the rays. Precisely, it is the root stack of order $\beta_F^{\text{res}} / \beta_F^{\text{can}}$ along the divisor defined by the ray F^\perp . The map $\mathcal{X}_{\Lambda, \beta^{\text{can}}} \rightarrow X_\Lambda$ is called the *canonical desingularisation*. The map $\mathcal{X}_{\Lambda, \beta^{\text{can}}} \rightarrow \mathbf{A}^n$ is an isomorphism away from the origin. The map $\mathcal{X}_{\Lambda, \beta^{\text{res}}} \rightarrow \mathbf{A}^n$ is an isomorphism away from the union of the coordinate hyperplanes.

Let x_1, \dots, x_n be variables. For $p = (p_1, \dots, p_n) \in \mathbf{Z}_{\geq 0}^n$, we write x^p for the monomial $x_1^{p_1} \dots x_n^{p_n}$. A *weighted monomial* is a pair (x^p, d) , where d is a positive integer. Let L be a set of weighted monomials. Let Λ be the rational Newton polyhedron defined by the points $\frac{1}{d}p$ for $(p, d) \in L$, that is, the convex hull of the union of $\frac{1}{d}p + \mathbf{R}_{\geq 0}^n$ for $(p, d) \in L$. Assume that Λ is simplicial. Then we have the stacky blow-ups $\mathcal{X}_{\Lambda, \beta^{\text{res}}}$ and $\mathcal{X}_{\Lambda, \beta^{\text{can}}}$. We call $\mathcal{X}_{\Lambda, \beta^{\text{can}}}$ the *canonical weighted blow-up* in the set of weighted monomials L , and $\mathcal{X}_{\Lambda, \beta^{\text{res}}}$ the *resolving weighted blow-up*. The following two propositions justify the name.

Proposition 2.2. *In the setup above, let $v = \frac{1}{d}p$ be a vertex of Λ . Consider the chart*

$$\text{Spec } \mathbf{k}[u_1, \dots, u_n] \rightarrow \mathcal{X}_{\Lambda, \beta^{\text{res}}}$$

defined by v . The image of x^p in $\mathbf{k}[u_1, \dots, u_n]$ is the d -th power of a monomial u . Furthermore, for every $(x^q, e) \in L$, the monomial u^e divides the image of x^q .

Proof. Set $\beta_i = \beta_{F_i}^{\mathrm{res}}$. Using (4), we see that

$$x^p \mapsto \prod_i \prod_j u_j^{p_i \langle \beta_j, e_i \rangle} = \prod_j u_j^{\langle \beta_j, p \rangle}.$$

By the choice of β_j , the quantity $\langle \beta_j, p/d \rangle$ is a non-negative integer. Thus, x^p maps to the d -th power of the monomial

$$u = \prod_j u_j^{\langle \beta_j, p/d \rangle}.$$

Consider $(x^q, e) \in L$. Let r_1, \dots, r_n be the rays of Λ incident to v . Then the point q/e is in the cone defined by the vertex v and the rays spanned by r_1, \dots, r_n . That is, we can write

$$q/e = p/d + a_1 r_1 + \dots + a_n r_n$$

for some non-negative rational numbers a_i . By applying β_i to both sides, we see that $e \cdot a_i$ is a non-negative integer. Using (4) again, we get

$$x^q \mapsto u^e \prod_i u_i^{e \cdot a_i}.$$

□

Let d_1, \dots, d_m be positive integers and let $\mathcal{P}(d_1, \dots, d_m)$ be the weighted projective stack

$$\mathcal{P}(d_1, \dots, d_m) = [(\mathbf{A}^m - 0)/\mathbf{G}_m],$$

where \mathbf{G}_m acts coordinate-wise by weights d_1, \dots, d_m . Consider the rational map

$$\mathbf{A}^n \dashrightarrow \mathcal{P}(d_1, \dots, d_m)$$

defined by the monomials x^{p_1}, \dots, x^{p_m} ; that is,

$$(5) \quad (x_1, \dots, x_n) \mapsto [x^{p_1} : \dots : x^{p_m}].$$

Let Λ be the Newton polyhedron defined by the weighted monomials $(x^{p_1}, d_1), \dots, (x^{p_m}, d_m)$. Assume that Λ is simplicial.

Proposition 2.3. *The map (5) extends uniquely to a morphism*

$$\mathcal{X}_{\Lambda, \beta^{\mathrm{res}}} \rightarrow \mathcal{P}(d_1, \dots, d_m).$$

Proof. The domain is normal and the co-domain is separated. Therefore, if the map extends, it extends uniquely [9, Appendix A]. To see that it extends, we may work locally on charts. Let v be a vertex of Λ , say $v = p_i/d_i$. By Proposition 2.2, on the chart $\mathrm{Spec} \mathbf{k}[u_1, \dots, u_n]$, the pull-back of x^{p_i} is u^{d_i} for a monomial u , and the pull-back of x^{p_j} is divisible by u^{d_j} . The extension of (5) on $\mathrm{Spec} \mathbf{k}[u_1, \dots, u_n]$ is given by

$$[x^{p_1} u^{-d_1} : \dots : x^{p_{i-1}} u^{-d_{i-1}} : 1 : x^{p_{i+1}} u^{-d_{i+1}} : \dots : x^{p_m} u^{-d_m}].$$

□

We reformulate (2.3) to suit our setting.

Corollary 2.4. *Let W_1, \dots, W_m be finite dimensional \mathbf{k} -vector spaces and set $W = \bigoplus_i W_i$. Let $\mathcal{P}W$ be the weighted projective stack where W_i has weight $d_i > 0$. Let $f: \mathbf{A}^n \dashrightarrow \mathcal{P}W$ be the rational map defined by $f_i \in W_i \otimes \mathbf{A}[x_1, \dots, x_n]$ and assume that the coordinates of f_i generate the monomial ideal $\langle x^{p_i} \rangle$. Let Λ be the Newton polyhedron defined by the weighted monomials $(x^{p_1}, d_1), \dots, (x^{p_m}, d_m)$. Then the rational map f extends uniquely to a morphism*

$$\mathcal{X}_{\Lambda, \beta^{\mathrm{res}}} \rightarrow \mathcal{P}W.$$

Proof. We follow the proof of Proposition 2.3. Let v be a vertex of Λ , say $v = p_i/d_i$. Let u^{d_i} be the pull-back of x^{p_i} to the chart defined by v . Then for all j , the element $u^{-d_j} f_j \in W_j \otimes \mathbf{k}[u_1, \dots, u_m]$ has polynomial coordinates. Furthermore, for $j = i$, the coordinates generate the unit ideal. On this chart, the extension of the rational map f is given by $[u^{-d_1} f_1 : \dots : u^{-d_m} f_m]$. \square

3. CLASS OF AN ORBIT USING A COMPLETE PARAMETRISATION

Let W be a finite dimensional representation of $\mathrm{GL}(m)$. Consider the central $\mathbf{G}_m \subset \mathrm{GL}(m)$, and assume that it acts on W by positive weights. We denote by $\mathcal{P}W$ the weighted projective stack

$$\mathcal{P}W = [W - 0 / \mathbf{G}_m].$$

Fix a non-zero vector $w \in W$, and let

$$[w]: \mathrm{Spec} \mathbf{k} \rightarrow \mathcal{P}W$$

be the corresponding point of $\mathcal{P}W$. By the *stabiliser* Γ of w , we mean the fiber product

$$(6) \quad \begin{array}{ccc} \Gamma & \longrightarrow & \mathrm{PGL}(m) \\ \downarrow & & \downarrow a \mapsto a \cdot w \\ \mathrm{Spec} \mathbf{k} & \xrightarrow{[w]} & \mathcal{P}W. \end{array}$$

We have the diagram

$$\begin{array}{ccccc} \Gamma & \longrightarrow & \mathrm{GL}(m) & \longrightarrow & \mathrm{PGL}_m \\ \downarrow & & \downarrow a \mapsto a \cdot w & & \downarrow a \mapsto a \cdot w \\ \mathrm{Spec} \mathbf{k} & \longrightarrow & W & \longrightarrow & \mathcal{P}W \end{array}$$

in which the right square and the outer square are cartesian. Therefore, the left square is also cartesian. Therefore, Γ is simply the stabiliser of w in $\mathrm{GL}(m)$.

A *complete orbit parametrisation* of $[w]$ is a proper morphism

$$i: X \rightarrow \mathcal{P}W,$$

where X is a Deligne–Mumford stack together with the action of $\mathrm{PGL}(m)$ and i is a $\mathrm{PGL}(m)$ -equivariant map such that there exists an open subscheme $U \subset X$ isomorphic to $\mathrm{PGL}(m)$ as a $\mathrm{PGL}(m)$ -scheme and a point $x \in U$ whose image is $[w]$. The *orbit* of $[w]$, denoted by $\mathrm{Orb}([w])$, is the Zariski closure in $\mathcal{P}W$ of $\mathrm{PGL}(m) \cdot [w]$, with the reduced scheme structure.

Proposition 3.1. *Let $i: X \rightarrow \mathcal{P}W$ be a complete parametrisation of the orbit of w . Assume that the stabiliser $\Gamma \subset \mathrm{GL}(m)$ of w is finite. Then, we have the equality of cycles*

$$i_*[X] = |\Gamma|[\mathrm{Orb}([w])].$$

Proof. We have the fiber product

$$(7) \quad \begin{array}{ccc} \Gamma & \longrightarrow & \mathrm{PGL}(m) \\ \downarrow & & \downarrow \\ \mathrm{Spec} \mathbf{k} & \xrightarrow{[w]} & \mathrm{Orb}([w]). \end{array}$$

Consider the open inclusion $\mathrm{PGL}(m) \rightarrow X$ that sends a to $a \cdot x$. The image of this inclusion is U . The points of X in the complement of U are stabilised by a positive dimensional subgroup of $\mathrm{PGL}(m)$ and hence they map to points in $\mathrm{Orb}([w])$ that are stabilised by a positive dimensional

subgroup. In particular, they do not map to $[w]$. As a result, the fiber product (7) gives the fiber product

$$\begin{array}{ccc} \Gamma & \longrightarrow & X \\ \downarrow & & \downarrow \\ \mathrm{Spec} \mathbf{k} & \xrightarrow{[w]} & \mathrm{Orb}([w]). \end{array}$$

We see that the map $X \rightarrow \mathrm{Orb}([w])$ is generically finite of degree $|\Gamma|$. The proposition follows. \square

We now give a cohomological formula for the push-forward. We first need a lemma, adapted from [6, Proposition 2.1]. Let U be a vector space of dimension N with the action of an algebraic group G . Set $U^* = U - 0$ and let $\pi: U^* \rightarrow \mathbf{P}U$ be the projection.

Lemma 3.2. *Let Y be a Deligne–Mumford stack with a G -action and a G -equivariant map $\phi: Y \rightarrow \mathbf{P}U$. Then, in $A_G(U^*)$, we have the equality*

$$\pi^* \phi_* [Y] = \int_Y \frac{c_N(U)}{\phi^* c_1 \mathcal{O}(-1)}.$$

The integral on the right is the push-forward $A_G(Y) \rightarrow A_G$, considered as an element of $A_G(U^*)$ via the pull-back $A_G \rightarrow A_G(U^*)$.

Proof. Let Q be the cokernel of $\phi^* \mathcal{O}(-1) \rightarrow U \otimes \mathcal{O}_Y$. On $Y \times \mathbf{P}U$, let π_i for $i = 1, 2$ be the two projections. The vanishing locus of the composite map

$$\pi_2^* \mathcal{O}(-1) \rightarrow U \otimes \mathcal{O}_{Y \times \mathbf{P}U} \rightarrow \pi_1^* Q$$

is precisely the graph Z of $\phi: Y \rightarrow \mathbf{P}U$. Therefore, we have

$$(8) \quad [Z] = c_{N-1}(\pi_1^* Q \otimes \pi_2^* \mathcal{O}(1)) [Y \times \mathbf{P}U].$$

Consider the fiber square

$$\begin{array}{ccc} X \times U^* & \xrightarrow{\tilde{\pi}_2} & U^* \\ \downarrow \tilde{\pi} & & \downarrow \pi \\ X \times \mathbf{P}U & \xrightarrow{\pi_2} & \mathbf{P}U. \end{array}$$

By the push-pull formula, we have

$$(9) \quad \widetilde{\pi}_{2*} \tilde{\pi}^* [Z] = \pi^* \pi_{2*} [Z].$$

The right-hand side of (9) is $\pi^* \phi_* [X]$. Since the pull-back of $\mathcal{O}_{\mathbf{P}U}(1)$ to U^* is trivial, (8) shows that

$$\tilde{\pi}^* [Z] = c_{N-1}(\pi_1^* Q) [Y \times U^*].$$

The statement follows by applying $\widetilde{\pi}_{2*}$ to the above equation. \square

We need an analogue of Lemma 3.2 for weighted projective spaces. Let W be a vector space of dimension N with an action of a torus T . Set $W^* = W - 0$ and $\mathscr{P}W = [W^* / \mathbf{G}_m]$, where \mathbf{G}_m acts on W by positive weights and this action commutes with the action of T . Let $\pi: W^* \rightarrow \mathscr{P}W$ be the projection.

Lemma 3.3. *Let X be a Deligne–Mumford stack with a T -action and a T -equivariant map $\phi: X \rightarrow \mathscr{P}W$. Then, in $A_T(W^*)$, we have the equality*

$$\pi^* \phi_* [X] = \int_X \frac{c_N(W)}{\phi^* c_1 \mathcal{O}_{\mathscr{P}W}(-1)}.$$

We understand the right-hand side in the same sense as in Lemma 3.2.

Proof. It suffices to prove the equality in $A_{\tilde{T}}(W^*)$ where $\tilde{T} \rightarrow T$ is a finite cover by another torus. Choose a basis $\langle w_i \rangle$ of W compatible with the action of T and \mathbf{G}_m . Suppose T acts on w_i by the character $\chi_i \in \text{Hom}(T, \mathbf{G}_m)$ and \mathbf{G}_m acts on w_i by weight d_i . Let $\tilde{T} \rightarrow T$ be a finite cover by a torus such that the image of χ_i in $\text{Hom}(\tilde{T}, \mathbf{G}_m)$ is divisible by d_i . Let U be the \mathbf{k} -span of the symbols u_i . Equip U with a \tilde{T} action so that \tilde{T} acts on u_i by the character $\frac{1}{d_i}\chi_i$ and with the \mathbf{G}_m action by weight 1. Then the map

$$\mu: U \rightarrow W$$

defined by $\sum x_i u_i \mapsto \sum x_i^{d_i} w_i$ is equivariant for the \tilde{T} and \mathbf{G}_m actions and finite of degree

$$\deg \mu = \prod d_i.$$

Under the induced map $\mu: \mathbf{P}U \rightarrow \mathcal{P}W$, the pull-back of $\mathcal{O}_{\mathcal{P}W}(-1)$ is $\mathcal{O}_{\mathbf{P}U}(-1)$. Define Y by the pull-back diagram

$$\begin{array}{ccc} Y & \xrightarrow{\tilde{\phi}} & \mathbf{P}U \\ \downarrow \tilde{\mu} & & \downarrow \mu \\ X & \xrightarrow{\phi} & \mathcal{P}W. \end{array}$$

Set $U^* = U - 0$ and denote by $\tilde{\pi}: U^* \rightarrow \mathbf{P}U$ the projection. By Lemma 3.2, in $A_{\tilde{T}}(U^*)$ we have

$$(10) \quad \tilde{\pi}^* \tilde{\phi}_*[Y] = \int_Y \frac{c_N(U)}{\phi^* c_1 \mathcal{O}_{\mathbf{P}U}(-1)}.$$

Note that $c_N(U) = \prod d_i^{-1} c_N(W)$. Since $Y \rightarrow X$ is of degree $\prod d_i$, the integral on the right-hand side of (10) is equal to

$$\int_X \frac{c_N(W)}{\phi^* c_1 \mathcal{O}_{\mathcal{P}W}(-1)}.$$

Now the statement follows by applying $\mu_*: A_{\tilde{T}}(U^*) \rightarrow A_{\tilde{T}}(W^*)$ to both sides of (10). \square

Let $\text{Orb}(w) \subset W$ be the closure of the $\text{GL}(m)$ -orbit of w , with the reduced scheme structure. Let $N = \dim W$.

Proposition 3.4. *Let $i: X \rightarrow \mathcal{P}W$ be a complete parametrisation of the orbit of $w \in W$. Assume that the stabiliser $\Gamma \subset \text{GL}(m)$ of w is finite. Then, in $A_G(W) = A_G$, we have*

$$|\Gamma| \cdot [\text{Orb}(w)] = \int_X \frac{c_N(W)}{i^* c_1 \mathcal{O}_{\mathcal{P}W}(-1)}.$$

The integral on the right is the push-forward $A_G(X) \rightarrow A_G$.

Proof. Let $T \subset \text{GL}(m)$ be a maximal torus. It suffices to prove the equality in $A_T(W)$. Proposition 3.1 and Lemma 3.3 together give the equality in $A_T(W^*)$. But then the equality also holds in $A_T(W)$ since $A_T^i(U) = A_T^i(U^*)$ for $i = \text{codim Orb}(w)$. \square

4. TWIST INVARIANCE

Consider a representation W of $\mathrm{GL}(2)$ defined by

$$\rho: \mathrm{GL}(2) \rightarrow \mathrm{GL}(W).$$

For $n \in \mathbf{Z}$, we have the surjective homomorphism

$$T_n: \mathrm{GL}(2) \rightarrow \mathrm{GL}(2), \quad M \mapsto M \cdot (\det M)^n,$$

whose kernel is the diagonally embedded $\mu_{n+2} \subset \mathrm{GL}(2)$. The composite $\rho \circ T_n$ gives a new representation $\mathrm{GL}(2) \rightarrow \mathrm{GL}(W)$ which we call $W(n)$. Note that

$$(11) \quad \text{if } W \cong \mathrm{Sym}^{a-b} V \otimes \det V^b, \text{ then } W(n) \cong \mathrm{Sym}^{a-b} V \otimes \det^{b+n(a+b)} V.$$

Observe that the identity map $W(n) \rightarrow W$ together with $T_n: \mathrm{GL}(2) \rightarrow \mathrm{GL}(2)$ induces a map

$$e_n: [W(n)/\mathrm{GL}(2)] \rightarrow [W/\mathrm{GL}(2)].$$

Given $w \in W$, the $\mathrm{GL}(2)$ -orbit closure of w in W under ρ is equal to that of w in $W(n)$. But to distinguish the ambient representations, we denote them by $\mathrm{Orb}(w)$ and $\mathrm{Orb}(w)(n)$, respectively. Then

$$\mathrm{Orb}(w)(n) = e_n^{-1}(\mathrm{Orb}(w)),$$

and hence

$$[\mathrm{Orb}(w)(n)] = e_n^*([\mathrm{Orb}(w)]) \in A_{\mathrm{GL}(2)}.$$

The map

$$e_n^*: A_{\mathrm{GL}(2)} \rightarrow A_{\mathrm{GL}(2)}$$

is easy to describe. Thinking of $A_{\mathrm{GL}(2)}$ as the subring of $\mathbf{Q}[v_1, v_2]$ consisting of symmetric polynomials, it is given by

$$(12) \quad e_n^*: v_1 \mapsto v_1 + n(v_1 + v_2) \text{ and } v_2 \mapsto v_2 + n(v_1 + v_2).$$

Let Γ and $\Gamma(n)$ be the stabilisers of w under ρ and $\rho \circ T_n$, respectively. Then we have the sequence

$$1 \rightarrow \mu_{n+2} \rightarrow \Gamma(n) \rightarrow \Gamma \rightarrow 1.$$

In particular, we have $|\Gamma(n)| = (n+2)|\Gamma|$.

Given $u \in \mathbf{P}^1$, let Λ^u be the Newton polygon associated to $w \in W$ and $\Lambda^u(n)$ the Newton polygon associated to $w \in W(n)$. Using (11), it follows that $\Lambda^u(n) \subset \mathbf{R}^2$ is obtained from $\Lambda^u \subset \mathbf{R}^2$ by applying the transformation

$$(13) \quad (x, y) \mapsto \frac{1}{n+2}(x+1, y+1).$$

Let Q be the polynomial on the right-hand side of (2) in the main theorem for $w \in W$ and $Q(n)$ the corresponding polynomial for $w \in W(n)$. Using (12) and (13), it is easy to check that

$$e_n^*(Q) = Q(n)/(n+2).$$

Since we also have

$$e_n^*(|\Gamma|[\mathrm{Orb}(w)]) = |\Gamma(n)|[\mathrm{Orb}(w)(n)]/(n+2),$$

the main theorem holds for $w \in W$ if and only if it holds for $w \in W(n)$. Thus, in the proof, we are free to replace W by $W(n)$ for any n . In particular, by choosing a sufficiently large n , we may assume without loss of generality that

$$(14) \quad W \cong \bigoplus \mathrm{Sym}^{a_i-b_i} V \otimes \det^{b_i} V \text{ with } a_i \geq b_i \geq 0.$$

5. COMPLETE ORBIT PARAMETRISATIONS OF $\mathrm{GL}(2)$ -ORBITS

Recall that we have a 2-dimensional vector space V and

$$W = W_1 \oplus \cdots \oplus W_n, \text{ where } W_i = \mathrm{Sym}^{a_i-b_i} V \otimes \det V^{b_i}.$$

We set $d_i = a_i + b_i$, which we call the *weight* of W_i , and assume $d_i > 0$ for all i . Also assume that $b_i \geq 0$; this can be achieved after twisting W as in Section 4. Consider the central $\mathbf{G}_m \rightarrow \mathrm{GL} V$ given by $t \mapsto t \cdot I$. Observe that $t \in \mathbf{G}_m$ scales the elements of W_i by t^{d_i} . If U is another 2-dimensional vector space, then by $W_i(U)$ we mean the representation

$$W_i(U) = \mathrm{Sym}^{a_i-b_i} U \otimes \det^{b_i} U,$$

and by $W(U)$ the direct sum

$$W(U) = \bigoplus_i W_i(U).$$

Let $\mathcal{P}W$ be the weighted projective stack

$$\mathcal{P}W = [W - 0 / \mathbf{G}_m]$$

Let U be another two-dimensional vector space and set

$$M = \mathbf{P} \mathrm{Hom}(U, V).$$

Fix a non-zero $w \in W(U)$. Let $w_i \in W_i(U)$ be the i -th component of w .

Let $I = \{i \mid w_i \neq 0\}$ and $J = \{1, \dots, n\} - I$. Set $W_I = \bigoplus_{i \in I} W_i$ and similarly for W_J . Let w_I be the projection of w to W_I . Plainly, we have $\mathrm{Orb}(w) = \mathrm{Orb}(w_I) \times \{0\} \subset W_I \oplus W_J$, and hence

$$[\mathrm{Orb}(w)] = c_{\dim W_J} (W_J) [\mathrm{Orb}(w_I)].$$

Using this, we see that it suffices to prove the main theorem when $J = \emptyset$. So, assume that $w_i \neq 0$ for all i .

We have a rational map

$$(15) \quad M \dashrightarrow \mathcal{P}W$$

defined by

$$m \mapsto [m \cdot w].$$

It is defined on the locus of m such that $m \cdot w \neq 0$. More formally, on $M = \mathbf{P} \mathrm{Hom}(U, V)$, we have the universal homomorphism

$$e: U \otimes \mathcal{O}_M(-1) \rightarrow V \otimes \mathcal{O}_M,$$

which induces

$$W_i(U) \otimes \mathcal{O}_M(-d_i) \rightarrow W_i(V) \otimes \mathcal{O}_M.$$

By pre-composing with the section

$$w_i: \mathcal{O}_M \rightarrow W_i(U_i) \otimes \mathcal{O}_M,$$

we get the map

$$(16) \quad \mathcal{O}_M(-d_i) \rightarrow W_i(V) \otimes \mathcal{O}_M.$$

The maps in (16) define a morphism to $\mathcal{P}W$ on the open subset of M where at least one of the maps is non-zero. Observe that this open subset includes all points of M corresponding to invertible homomorphisms.




FIGURE 1. The scheme theoretic zero locus of the map (16) is cut out locally by an ideal of the form $I_{K_u}^r \cdot I_\Delta^b$, where $\Delta \subset M$ is the determinant quadric and $K_u \subset \Delta$ are certain lines on it.

We now describe the scheme theoretic zero locus of the map (16). It is supported on the determinant quadric

$$\Delta = \{m \in M \mid \det m = 0\},$$

and it has embedded primes supported on lines of one ruling of this quadric (see Proposition 5.1 and Figure 1). Given a point $u \in \mathbf{PU}$, let $K_u \subset M$ be the line defined by

$$K_u = \{m \in M \mid mu = 0\}.$$

Observe that as u varies in $\mathbf{PU} \cong \mathbf{P}^1$, the lines K_u sweep out one of the two rulings of Δ .

Proposition 5.1. *Suppose $w_i = f \otimes \delta^{b_i}$, where $f \in \text{Sym}^{a_i - b_i}(U)$ and $\delta \in \det U$ are non-zero. Take $m \in \Delta \subset M$ and let $u \in \mathbf{PU}$ be the kernel of m . Suppose f vanishes to order r at u . Then, in a neighbourhood of m , the scheme theoretic zero locus of the map*

$$e: \mathcal{O}_M(-d_i) \rightarrow W_i(V) \otimes \mathcal{O}_M,$$

defined in (16), is cut out by the ideal $I_{K_u}^r \cdot I_\Delta^{b_i}$.

Proof. Since i is fixed, we omit it from the subscript in a_i, b_i , and d_i , and do a local calculation. Denote by $u_2 \in U$ a lift of $u \in \mathbf{PU}$. We choose a linearly independent vector $u_1 \in U$ and take (u_1, u_2) as a basis of U . In this basis, the point $u \in \mathbf{PU}$ is given by $[0 : 1]$.

Choose a basis (v_1, v_2) of V and suppose that in the chosen bases, the map $m: U \rightarrow V$ is given by

$$m = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

We can take $\delta = u_1 \wedge u_2$. Consider the affine neighbourhood of $m \in M$ given by matrices of the form

$$\begin{pmatrix} 1 & x \\ z & y + xz \end{pmatrix}.$$

Up to multiplication by a non-zero scalar, the element $f = f(u_1, u_2)$ has the form

$$(17) \quad f(u_1, u_2) = u_1^{a-b-r} u_2^r + * \cdot u_1^{a-b-r-1} u_2^{r+1} + \dots$$

Substituting $u_1 \mapsto v_1 + zv_2$ and $u_2 \mapsto xv_1 + (y+xz)v_2$ yields

$$(18) \quad \begin{aligned} (Mf)(v_1, v_2) &= (v_1 + zv_2)^{a-b-r} (xv_1 + (y+xz)v_2)^r \\ &\quad + * \cdot (v_1 + zv_2)^{a-b-r-1} (xv_1 + (y+xz)v_2)^{r+1} + \dots, \end{aligned}$$




FIGURE 2. For $W = \text{Sym}^3 V \oplus \text{Sym}^2 V \otimes \det V$, the Newton polygon Λ^u at $u \in \mathbf{P}^1$ with the vanishing orders $r_1 = 2$ and $r_2 = 0$. The short normal vectors (dashed) represent β^{can} and the longer ones (dotted) represent β^{res} .

and,

$$(19) \quad (M\delta)(v_1, v_2) = y(v_1 \wedge v_2).$$

Observe that the ideal generated by the coefficients of Mf is $\langle x, y \rangle^r$. The ideal $\langle x, y \rangle$ is precisely the ideal I_{K_u} and the ideal $\langle y \rangle$ is precisely the ideal I_Δ . So the ideal generated by the coefficients of $M(f \otimes \delta^b)$ is $I_{K_u}^r \cdot I_\Delta^b$, as required. \square

We now resolve the rational map $M \dashrightarrow \mathcal{P}W$ using a stacky blow-up. Let the components of w be $w_i = f_i \otimes \delta^{b_i}$. Let $A \subset \mathbf{P}U$ be any finite set that includes the common zeros of f_i for i that realise the minimum $\min_i b_i/d_i$. Let $\text{Bl}_A M \rightarrow M$ be the blow-up of M along the lines K_u for $u \in A$. Let $E_u \subset \text{Bl}_A M$ be the exceptional divisor over K_u and $D \subset \text{Bl}_A M$ the proper transform of $\Delta \subset M$. Let r_i^u be the order of vanishing of f_i at u . Let

$$e: \mathcal{O}_{\text{Bl}_A M}(-d_i) \rightarrow W_i(V) \otimes \mathcal{O}_{\text{Bl}_A M}$$

be the pull-back of (16). By Proposition 5.1, the ideal generated by the components of this map is $I_{E_u}^{r_i^u + b_i} I_D^{b_i}$.

Fix $u \in A \subset \mathbf{P}U$. Let $\Lambda = \Lambda^u \subset \mathbf{R}^2$ be the Newton polygon defined by the set of weighted monomials $\{(x^{r_i^u + b_i} y^{b_i}, d_i) \mid i = 1, \dots, n\}$. Let $\overline{\mathcal{X}}_{\Lambda, \beta} \rightarrow [\mathbf{A}^2/\mathbf{G}_m^2]$ be the blow-up defined in Section 2 (see Remark 2.1) for $\beta = \beta^{\text{can}}$ and $\beta = \beta^{\text{res}}$. Let $\text{Bl}_A M \rightarrow [\mathbf{A}^2/\mathbf{G}_m^2]$ be the map defined by the divisors E_u and D and let $\mathcal{M}_u^{\text{res}} \rightarrow \text{Bl}_A M$ and $\mathcal{M}_u^{\text{can}} \rightarrow \text{Bl}_A M$ be the pullbacks of $\overline{\mathcal{X}}_{\Lambda, \beta^{\text{res}}} \rightarrow [\mathbf{A}^2/\mathbf{G}_m^2]$ and $\overline{\mathcal{X}}_{\Lambda, \beta^{\text{can}}} \rightarrow [\mathbf{A}^2/\mathbf{G}_m^2]$. Since E_u and D are smooth, normal crossings divisors, the map $\text{Bl}_A M \rightarrow [\mathbf{A}^2/\mathbf{G}_m^2]$ is smooth, and therefore both $\mathcal{M}_u^{\text{res}}$ and $\mathcal{M}_u^{\text{can}}$ are smooth. See Figure 2 for an example of Λ^u with β^{can} and β^{res} .

We can write local charts for $\mathcal{M}_u^{\text{res}}$ and $\mathcal{M}_u^{\text{can}}$ by simply substituting local equations of E_u and D in the local charts described in Section 2. Let $\lambda = (\lambda_1, \lambda_2)$ be the vertex of the Newton polyhedron $\Lambda = \Lambda^u$ with the smallest second coordinate. Then $\lambda_2 = \min_i (b_i/d_i)$. Suppose $\lambda_2 = b/d$ where $\gcd(b, d) = 1$. Note that $\lambda + \mathbf{R}_{\geq 0} \times 0$ is a ray of Λ . Its associated divisor is the vanishing locus of y , which pulls back to $D \subset \text{Bl}_A M$. The functional β_1^{can} associated to this ray is the projection $p_2: (a, b) \mapsto b$. On the other hand, the functional β_1^{res} is $d \cdot p_2$. As a result, $\mathcal{M}_u^{\text{res}} \rightarrow \text{Bl}_A M$ over the complement of E_u is the root stack along D of order d . Note that d is independent of $u \in A$.

Let $\mathcal{M}^{\text{res}} \rightarrow \text{Bl}_A M$ and $\mathcal{M}^{\text{can}} \rightarrow \text{Bl}_A M$ be the blow-ups as above carried out for all $u \in A$ at once. That is, for all $u \in A$, in a neighbourhood of E_u , the map $\mathcal{M}^{\text{res}} \rightarrow \text{Bl}_A M$ is the blow-up

$\mathcal{M}_u^{\mathrm{res}} \rightarrow \mathrm{Bl}_A M$, and similarly for $\mathcal{M}^{\mathrm{can}} \rightarrow \mathrm{Bl}_A M$. We have maps

$$\mathcal{M}^{\mathrm{res}} \rightarrow \mathcal{M}^{\mathrm{can}} \rightarrow \mathrm{Bl}_A M.$$

The map $\mathcal{M}^{\mathrm{can}} \rightarrow \mathrm{Bl}_A M$ is an isomorphism away from the union of the lines $E_u \cap D$ for $u \in A$. The map $\mathcal{M}^{\mathrm{res}} \rightarrow \mathrm{Bl}_A M$ is an isomorphism away from the union of the divisors E_u for $u \in A$ and D . Over the complement of the union of E_u for $u \in A$, it is the root stack of order d along D .

Let \mathcal{M} be $\mathcal{M}^{\mathrm{can}}$ or $\mathcal{M}^{\mathrm{res}}$. For every $g \in \mathrm{GL} V$, it is easy to check that the action map $g: M \rightarrow M$ lifts to a morphism $g: \mathcal{M} \rightarrow \mathcal{M}$. For $g, h \in \mathrm{GL} V$, the two morphisms $h \circ g$ and gh agree on a dense open subscheme in \mathcal{M} . Since \mathcal{M} is normal and separated, [9, Appendix A] implies that there exists a unique 2-morphism $h \circ g \Rightarrow gh$. As a result, the maps $g: \mathcal{M} \rightarrow \mathcal{M}$ for $g \in \mathrm{GL} V$ give an action of $\mathrm{GL} V$ on \mathcal{M} .

Proposition 5.2. *The rational map*

$$M \dashrightarrow \mathcal{P} W$$

extends to a regular map

$$\iota: \mathcal{M}^{\mathrm{res}} \rightarrow \mathcal{P} W,$$

which is a complete orbit parametrisation of the orbit of $[w] \in \mathcal{P} W$.

Proof. The extension exists due to Proposition 2.3 (see Corollary 2.4). It is immediate that ι gives a complete orbit parametrisation. \square

6. ATIYAH–BOTT LOCALISATION

Proposition 3.1 gives a formula for $[\mathrm{Orb}(w)]$ as an integral. We compute the integral in Proposition 3.1 using the Atiyah–Bott localisation formula for stacks [13, § 5.3]. In this section, we use \mathcal{M} to denote either $\mathcal{M}^{\mathrm{res}}$ or $\mathcal{M}^{\mathrm{can}}$. A claim about \mathcal{M} is understood to hold for both $\mathcal{M}^{\mathrm{res}}$ and $\mathcal{M}^{\mathrm{can}}$. Most such claims will be on the level of points or rational Chow groups, both of which are identical for the two stacks.

Fix a basis (v_1, v_2) of V . Let $T \subset \mathrm{GL} V$ be the diagonal torus with respect to the chosen basis. The T -fixed locus in M is the disjoint union of the two lines L_i for $i = 1, 2$ defined by

$$L_i = \{m \mid \mathrm{Image}(m) \subset \langle v_i \rangle\}.$$

These are lines on Δ of the opposite ruling compared to the lines K_u (see Figure 1).

6.1. Fixed points of the T -action on \mathcal{M} . Let $\mathcal{L}_i^{\mathrm{res}} \subset \mathcal{M}^{\mathrm{res}}$ and $\mathcal{L}_i^{\mathrm{can}} \subset \mathcal{M}^{\mathrm{can}}$ be the proper transforms of $L_i \subset M$ (with the reduced scheme structure). We use $\mathcal{L}_i \subset \mathcal{M}$ to refer to either one of these.

Fix a $u \in A \subset \mathbf{P} U$. Choose a basis u_1, u_2 of U such that $u = [u_2]$. Consider the affine open chart $\mathbf{A}_{x,y,z}^3 \subset M$ consisting of matrices of the form

$$\begin{pmatrix} 1 & x \\ z & y + xz \end{pmatrix}.$$

In this basis, the line L_1 , the line K_u , and the determinant Δ are cut out by

$$\begin{aligned} L_1 : z = y = 0, \\ K_u : x = y = 0, \\ \Delta : y = 0. \end{aligned}$$

The line L_2 is absent from this chart. Thinking of x, y, z as regular functions on this chart, we see that the T -action is given by

$$(20) \quad (t_1, t_2): (x, y, z) \mapsto (x, t_1 t_2^{-1} y, t_1 t_2^{-1} z).$$

The blow-up $\text{Bl}_{K_u} M$ has the local description

$$\{(x, y, z, [X : Y]) \mid Xy = xY\} \subset \text{Spec } \mathbf{k}[x, y, z] \times \mathbf{P}^1.$$

On the blow-up, the proper transform of L_1 is cut out by $z = 0$ and $Y = 0$. The only T -fixed points on the blow-up are the points of the proper transform of L_1 and the point p_1^u with coordinates $((0, 0, 0), [0 : 1])$.

The proper transform of Δ is defined by $Y = 0$, and is thus contained in the affine chart of the blow-up given by $X \neq 0$. This chart is given by

$$\{(x, y, z, [1 : Y]) \mid y = xY\} \cong \text{Spec } \mathbf{k}[x, Y, z].$$

The T -action is given by

$$(t_1, t_2): (x, Y, z) \mapsto (x, t_1^{-1} t_2 Y, t_1^{-1} t_2 z).$$

The stacky blow-up of this chart is defined by the weighted monomials $(x^{r_i^u + b_i} Y^{b_i}, d_i)$. Let Λ^u be the Newton polyhedron defined by these weighted monomials (see Section 2). Since z is absent from the monomials, we may think of Λ^u as a subset of \mathbf{R}^2 . Let $\lambda(0), \dots, \lambda(k)$ be the vertices of Λ^u arranged from the bottom-right to the top-left. That is, using subscripts to denote first and second coordinates, we have

$$\lambda(0)_1 > \dots > \lambda(k)_1 \text{ and } \lambda(0)_2 < \dots < \lambda(k)_2.$$

Note that the point corresponding to the bottom-right vertex $\lambda(0)$ lies on the proper transform of L_1 . It is easy to check that the only T -fixed points on the stacky blow-up of this chart are:

- (1) points of the proper transform of L_1 ,
- (2) points corresponding to the vertices $\lambda(1), \dots, \lambda(k)$.

For $j = 1, \dots, k$, we label the point corresponding to $\lambda(j)$ as $p_{1,j}^u$.

We have analogous points p_2^u and $p_{2,j}^u$ over the line $L_2 \subset M$.

Summarising the discussion above, we see that the T -fixed locus of \mathcal{M} is the disjoint union of

- (1) $\mathcal{L}_1 \sqcup \mathcal{L}_2$
- (2) $\{p_1^u, p_2^u\}$ for $u \in A$.
- (3) $\{p_{1,j}^u, p_{2,j}^u \mid j = 1, \dots, k = k^u\}$ for $u \in A$.

6.2. Ingredients of the localisation formula. Recall that we have the map

$$\iota: \mathcal{M}^{\text{res}} \rightarrow \mathcal{P}W,$$

which is the complete orbit parametrisation of $[w]$. We describe the pull-back of $\mathcal{O}(-1)$ and the normal bundles to the components of the fixed locus of the T -action as elements of the corresponding (rational) T -equivariant Grothendieck groups.

Let $M_T = \text{Hom}(T, \mathbf{G}_m) \otimes \mathbf{Q}$ and $K_T = \mathbf{Z}[M_T]$. We use \oplus to denote the formal sums in K_T . By a *rational T -representation*, we mean a representation of a finite cover of T . Every rational T -representation has a class in K_T . In particular, for $m, n \in \mathbf{Q}$, we have classes $\chi(m, n) \in K_T$ of rational characters. See the discussion before [13, Proposition 5.3.4] for the need to accommodate rational representations.

Proposition 6.1. *Fix $u \in A \subset \mathbf{PU}$. Suppose i realises the minimum $\min_i(\frac{1}{d_i}(r_i^u + b_i))$. Then the map*

$$(21) \quad \iota^* \mathcal{O}(-d_i) \rightarrow W_i \otimes \mathcal{O}_{\mathcal{M}^{\mathrm{res}}}$$

is non-zero at p_1^u and its image is spanned by $v_1^{a_i-b_i-r_i} v_2^{r_i} \otimes (v_1 \wedge v_2)^{b_i}$.

Proof. In the local coordinates introduced in Section 6.1, the point p_1^u lies in the chart

$$\{(x, y, z, [X : 1]) \mid x = yX\} \cong \mathrm{Spec} \mathbf{k}[X, y, z]$$

of the blow-up $\mathrm{Bl}_{K_u} M$. The stacky blow-up $\mathcal{M}^{\mathrm{res}}$ of this chart is defined by the weighted monomials $(y^{r_i^u+b_i}, d_i)$. Suppose the minimum in the statement is c/d , where $\gcd(c, d) = 1$. From (3) and (4), we get the following local chart of $\mathcal{M}^{\mathrm{res}} \rightarrow \mathrm{Bl}_{K_u} M$ at p_1^u :

$$[\mathrm{Spec} \mathbf{k}[u_1, u_2, u_3]/\mu_d] \rightarrow \mathrm{Spec} \mathbf{k}[X, y, z],$$

where the map is defined by

$$X \mapsto u_1, \quad y \mapsto u_2^d, \quad z \mapsto u_3.$$

Let $r = r_i^u$ and $b = b_i$. From the proof of Corollary 2.4, we know that on $\mathrm{Spec} \mathbf{k}[u_1, u_2, u_3]$, the map (21) is y^{-r-b} times the original map e studied in Proposition 5.1. From (18) and (19), we see that the map e is given by

$$y^b ((v_1 + zv_2)^{a-b-r} y^r (Xv_1 + (1 + Xz)v_2)^r + \cdots) \otimes (v_1 \wedge v_2)^b.$$

Multiplying by y^{-r-b} and setting $u_1 = u_2 = u_3 = 0$ yields the result. \square

Let $\mathcal{O}(-1)_{p_1^u}$ be the class in K_T of the fiber of $\iota^* \mathcal{O}(-1)$ at p_1^u . Similarly, let $N_{p_1^u}$ be the class in K_T of the normal bundle of p_1^u in $\mathcal{M}^{\mathrm{can}}$. Let $r^u = \min_i(r_i^u + b_i)/d_i$.

Proposition 6.2. *With the notation above, we have*

$$\begin{aligned} \mathcal{O}(-1)_{p_1^u} &= \chi(1 - r^u, r^u), \text{ and} \\ N_{p_1^u} &= \chi(1, -1) \oplus \chi(-1, 1) \oplus \chi(-1, 1). \end{aligned}$$

Proof. Suppose i realises the minimum $\min_i(r_i^u + b_i)/d_i$. Proposition 6.1 identifies the fiber of $\iota^* \mathcal{O}(-d_i)$ at p_1^u with the span of $v_1^{a_i-r_i} v_2^{r_i} \otimes (v_1 \wedge v_2)^{b_i}$, on which T acts by weights $a_i - r_i$ and $b_i + r_i$. Dividing through by d_i yields the first equality.

The map $\mathcal{M}^{\mathrm{can}} \rightarrow \mathrm{Bl}_A M$ is an isomorphism near p_1^u . The normal space at p_1^u is spanned by $\frac{\partial}{\partial X}$, $\frac{\partial}{\partial y}$, and $\frac{\partial}{\partial z}$, on which the T acts by weights $(1, -1)$, $(-1, 1)$, and $(-1, 1)$, respectively. \square

Proposition 6.3. *Fix $u \in A \subset \mathbf{PU}$. Let λ be a vertex of Λ^u , say $\lambda = \frac{1}{d_i}(r_i^u + b_i, b_i)$. Then the map*

$$(22) \quad \iota^* \mathcal{O}(-d_i) \rightarrow W_i \otimes \mathcal{O}_{\mathcal{M}^{\mathrm{res}}}$$

is non-zero at $p_{1,j}^u$ and its image is spanned by $v_1^{a_i-b_i} \otimes (v_1 \wedge v_2)^{b_i}$.

Proof. The proof is parallel to the proof of Proposition 6.1. In the local coordinates introduced in Section 6.1, consider the chart of $\mathrm{Bl}_{K_u} M$ given by

$$\{(x, y, z, [1 : Y]) \mid y = xY\} \cong \mathrm{Spec} \mathbf{k}[x, Y, z].$$

Consider the chart of $\mathcal{M}^{\mathrm{res}} \rightarrow \mathrm{Bl}_A M$ given by (3) and (4):

$$[\mathbf{k}[u_1, u_2, u_3]/\mu] \rightarrow \mathbf{k}[x, Y, z].$$

Let $r = r_i^u$ and $b = b_i$. From the proof of Corollary 2.4, we know that on $\text{Spec } \mathbf{k}[u_1, u_2, u_3]$, the map (22) is $x^{-r-b}Y^{-b}$ times the original map e studied in Proposition 5.1. From (18) and (19), we see that the map e is given by

$$x^b Y^b ((v_1 + zv_2)^{a-b-r} x^r (v_1 + (1 + Yz)v_2)^r + \cdots) \otimes (v_1 \wedge v_2)^b.$$

Multiplying by $x^{-r-b}Y^{-b}$ and setting $u_1 = u_2 = u_3 = 0$ yields the result. \square

Let $\mathcal{O}(-1)_{p_{\ell,j}^u}$ be the class in K_T of the fiber of $\iota^*\mathcal{O}(-1)$ at $p_{\ell,j}^u$. Similarly, let $N_{p_{1,j}^u}$ be the class in K_T of the normal bundle of $p_{1,j}^u$ in \mathcal{M}^{can} . Let η and ζ be the shortest integral normal vectors to the two rays of Λ^u at the vertex $\lambda(j)$. Let $N = \det(\eta, \zeta)$, so that $|N|$ is the index of the sub-lattice $\langle \eta, \zeta \rangle \subset \mathbf{Z}^2$.

Proposition 6.4. *With the notation above, we have*

$$\begin{aligned} \mathcal{O}(-1)_{p_{1,j}^u} &= \chi(1 - \lambda(j)_2, \lambda(j)_2), \text{ and} \\ N_{p_{1,j}^u} &= \chi(\zeta_1/N, -\zeta_1/N) \oplus \chi(-\eta_1/N, \eta_1/N) \oplus \chi(-1, 1). \end{aligned}$$

Proof. We use the notation in Proposition 6.3. In particular, we let i be such that $\lambda(j) = \frac{1}{d_i}(a_i + r_i^u, b_i)$. Proposition 6.3 shows that $\mathcal{O}(-1)_{p_{1,j}^u} = \frac{1}{d_i}\chi(a_i, b_i)$, by the same argument as Proposition 6.2. Since $d_i = a_i + b_i$, we can re-write this as $\chi(1 - \lambda(j)_2, \lambda(j)_2)$.

For the second equality, we write $\mathcal{M}^{\text{can}} \rightarrow \text{Bl}_A M$ in charts at $p_{1,j}^u$ using (3) and (4):

$$[\text{Spec } \mathbf{k}[u_1, u_2, u_3]/\mu] \rightarrow \text{Spec } \mathbf{k}[x, Y, z],$$

where the map is given by

$$x \mapsto u_1^{\eta_1} u_2^{\zeta_1}, \quad Y \mapsto u_1^{\eta_2} u_2^{\zeta_2}, \quad z \mapsto u_3.$$

The torus T acts on x, Y , and z by weights $(0, 0)$, $(1, -1)$, and $(1, -1)$, respectively. It follows that it must act on u_1, u_2 , and u_3 by weights $\frac{1}{N}(-\zeta_1, \zeta_1)$, $\frac{1}{N}(\eta_1, -\eta_1)$, and $(1, -1)$, respectively. Since the normal space to $p_{1,j}^u$ is spanned by $\frac{\partial}{\partial u_1}$, $\frac{\partial}{\partial u_2}$, and $\frac{\partial}{\partial u_3}$, the second equality follows. \square

Remark 6.5. In Proposition 6.4, suppose $\lambda = \lambda(0)$ is the bottom right vertex of Λ^u . Then one of the two rays incident at λ is $\lambda + \mathbf{R}_{\geq 0}$, so we may choose $\eta = (0, 1)$. In that case, we have a trivial summand $\chi(0, 0)$ in $N_{p_{1,0}^u}$. This summand corresponds to the normal direction in \mathcal{L}_1 , which is fixed by T .

We have now described all the ingredients of the localisation formula for the isolated fixed points. We now turn to the T -fixed lines \mathcal{L}_i . Note that the coarse space of \mathcal{L}_i is \mathbf{P}^1 . Use K to denote the numerical Grothendieck group (two classes considered equal if they have the same Chern character). For a finite cover $\tilde{T} \rightarrow T$, a \tilde{T} -equivariant bundle on \mathcal{L}_i has a class in $K_T \otimes K(\mathcal{L}_1)$. For $a \in \mathbf{Q}$, the notation $\mathcal{O}(a)$ denotes the class in $K(\mathcal{L}_i)$ of a line bundle of degree a .

For $u \in A$, we denote by $\lambda^u(j)$ the j -th vertex of $\Lambda^u \subset \mathbf{R}^2$ with the convention that the vertices are arranged from the bottom-right to the top-left (in the increasing order by the second coordinate). Recall that $b = \min_i(b_i/d_i)$ and $r_{\text{gen}}^u = \lambda^u(0)_1 - b$ and $r_{\text{gen}} = \sum_{u \in A} r_{\text{gen}}^u$.

Proposition 6.6. *The class of $\iota^*\mathcal{O}(-1)$ restricted to \mathcal{L}_1 in $K_T \otimes K(\mathcal{L}_1)$ is given by*

$$\iota^*\mathcal{O}(-1)|_{\mathcal{L}_1} = \chi(1 - b, b) \otimes \mathcal{O}(-1 + 2b + r_{\text{gen}}).$$

Proof. Recall that the points $p_{1,0}^u$ lie on \mathcal{L}_1 . Proposition 6.4 applied to $j = 0$ shows that the fiber of $\iota^*\mathcal{O}(-1)$ at $p_{1,0}^u$, as a rational T -representation, is $\chi(1-b, b)$. So, the class of $\iota^*\mathcal{O}(-1)$ restricted to \mathcal{L}_1 is $\chi(1-b, b) \otimes \mathcal{O}(a)$ for some $a \in \mathbf{Q}$, which is simply the degree of $\iota^*\mathcal{O}(-1)$ on \mathcal{L}_1 .

Let $\pi: \mathcal{M}^{\text{res}} \rightarrow M$ be the natural map. Then

$$\iota^*\mathcal{O}(-1) = \pi^*\mathcal{O}_M(-1) \otimes \mathcal{O}(E),$$

where $E \subset \mathcal{M}^{\text{res}}$ is an effective divisor. The divisor is characterised by the property that in a neighbourhood of a point $p \in \mathcal{M}^{\text{res}}$ at which the map $\iota^*\mathcal{O}(-d_i) \rightarrow W_i \otimes \mathcal{O}_{\mathcal{M}^{\text{res}}}$ is non-zero, the divisor $d_i E$ is the vanishing locus of $e: \pi^*\mathcal{O}_M(-d_i) \rightarrow W_i \otimes \mathcal{O}_{\mathcal{M}^{\text{res}}}$. We use this characterisation at $p_{1,0}^u$ for every $u \in A \subset \mathbf{PU}$. Given $u \in A$, let i be such that $\lambda^u(0) = \frac{1}{d_i}(r_i^u + b_i, b_i)$. By Proposition 6.3, the map $\iota^*\mathcal{O}(-d_i) \rightarrow W_i \otimes \mathcal{O}_{\mathcal{M}^{\text{res}}}$ is non-zero. The proof of Proposition 6.3 shows that the vanishing locus of $e: \pi^*\mathcal{O}_M(-d_i) \rightarrow W_i \otimes \mathcal{O}_{\mathcal{M}^{\text{res}}}$ is cut out in the local coordinates by $x^{r_i^u+b_i}Y^{b_i} = x^{r_i^u}y^{b_i}$. The divisor cut out by y is the pre-image of the determinant $\Delta \subset M$. The divisor cut out by x is the pre-image of the exceptional divisor $E_u \subset \text{Bl}_A M$. Therefore, in a neighbourhood of $p_{1,0}^u$, we have

$$\begin{aligned} E &= \frac{b_i}{d_i}\Delta + \frac{r_i^u}{d_i}E_u \\ &= b\Delta + r_{\text{gen}}^u E_u \end{aligned}$$

Considering all $u \in A \subset \mathbf{PU}$, we see that in a neighbourhood of \mathcal{L}_1 , we have

$$E = b\Delta + \sum_{u \in A} r_{\text{gen}}^u E_u.$$

On \mathcal{L}_1 , the degree of Δ is 2 and the degree of E_u is 1. The result follows. \square

Let $N_1 \in K_T \otimes K(\mathcal{L}_1)$ be the class of the normal bundle of $\mathcal{L}_1^{\text{can}} \subset \mathcal{M}^{\text{can}}$. For $u \in A \subset \mathbf{PU}$, if Λ^u has at least two vertices, set

$$s^u = 1 - \frac{\lambda^u(0)_1 - \lambda^u(1)_1}{\lambda^u(0)_2 - \lambda^u(1)_2}.$$

Otherwise, set $s^u = 1$. Let $s = \sum_{u \in A} s^u$.

Proposition 6.7. *With the notation above, the class of the normal bundle N_1 is equal to*

$$\chi(-1, 1) \otimes (\mathcal{O} \oplus \mathcal{O}(2-s)).$$

Proof. For simplicity, we drop the superscript “can”, but alert the reader that it is important that we are working with \mathcal{M}^{can} and not \mathcal{M}^{res} . We use the local coordinates introduced in Section 6.1. At a generic point $(x, 0, 0) \in \mathcal{L}_1$, the normal bundle is spanned by $\frac{\partial}{\partial y}$ and $\frac{\partial}{\partial z}$ on which T acts by weights $(-1, 1)$. Therefore, the class of N_1 is $\chi(-1, 1)$ times the class in $K(\mathcal{L}_1)$ of the normal bundle. We simply need to find the degree of N_1 .

Let $\pi: \mathcal{M} \rightarrow \text{Bl}_A M$ be the natural map. Let $\tilde{L}_1 \subset \text{Bl}_A M$ be the proper transform of L_1 . Consider the sequence

$$0 \rightarrow N_1 \xrightarrow{d\pi} \pi^*N_{\tilde{L}_1/\text{Bl}_A M} \rightarrow Q \rightarrow 0,$$

so that Q is supported on $\{p_{1,0}^u \mid u \in A\}$. Let $\eta = (0, 1)$ and ζ be the shortest integer normal vectors to the two rays of Λ^u at $\lambda(0)$. Then, in a neighbourhood of $p_{1,0}^u$, the map π is

$$[\text{Spec } \mathbf{k}[u_1, u_2, u_3]/\mu] \rightarrow \text{Spec } \mathbf{k}[x, Y, z],$$

where

$$x \mapsto u_2^{\zeta_1}, \quad Y \mapsto u_1 u_2^{\zeta_2}, \quad z \mapsto u_3,$$

and μ is a cyclic group of order ζ_1 . At $p_{1,0}^u$, the dual of $N_{\tilde{L}_1/\mathrm{Bl}_A M}$ is spanned by dY and dz , whereas the dual of N_1 is spanned by du_1 and du_3 . On \mathcal{L}_1 , which is cut out by $u_1 = u_3 = 0$, we have

$$dY = u_2^{\zeta_2} du_1 \text{ and } dz = du_3.$$

Therefore, the pull-back of Q to $\mathbf{k}[u_1, u_2, u_3]$ has length ζ_2 . But since the order of μ is ζ_1 , the degree of Q at $p_{1,0}^u$ is ζ_2/ζ_1 . If Λ^u has only one vertex, then $\zeta = (1, 0)$, so $\zeta_2 = 0$. Otherwise, ζ_2/ζ_1 is the negative of the reciprocal of the slope of the line joining $\lambda(0)$ and $\lambda(1)$, which is precisely $s^u - 1$. Therefore, we conclude that

$$\deg N_1 = \deg N_{\tilde{L}_1/\mathrm{Bl}_A M} - \sum_{u \in A} (s^u - 1).$$

But we also know that

$$\deg N_{\tilde{L}_1/\mathrm{Bl}_A M} = \deg N_{L_1/M} - \sum_{u \in A} 1 = 2 - \sum_{u \in A} 1.$$

Combining the two yields the proposition. \square

6.3. Proof of the main theorem. We now have the tools to prove Theorem 1.3. Let $T \subset \mathrm{GL}(V)$ be a maximal torus. Let $N = \dim W$. By Proposition 3.1, we have

$$|\Gamma| \cdot [\mathrm{Orb}(w)] = \int_{\mathcal{M}^{\mathrm{res}}} \frac{c_N(W)}{\iota^* c_1 \mathcal{O}_{\mathcal{P}W}(-1)}.$$

The pull-back along $\mathcal{M}^{\mathrm{res}} \rightarrow \mathcal{M}^{\mathrm{can}}$ identifies the rational Chow groups, and the push-forward of the fundamental class of $\mathcal{M}^{\mathrm{res}}$ is the fundamental class of $\mathcal{M}^{\mathrm{can}}$. Therefore, we may replace $\mathcal{M}^{\mathrm{res}}$ in the integral by $\mathcal{M}^{\mathrm{can}}$.

From Section 6.1, recall that the T -fixed points of $\mathcal{M}^{\mathrm{can}}$ consist of the line $\mathcal{L}_1^{\mathrm{can}}$, the points p_1^u for $u \in A$, the points $p_{1,j}^u$ for $u \in A$ and $j = 1, \dots, k^u$, where k^u is the number of vertices of the Newton polygon Λ^u , and their analogues where the subscript 1 is replaced by 2. For $\ell = 1, 2$, we denote by N_ℓ the normal bundle of $\mathcal{L}_\ell^{\mathrm{can}}$, by $N_{p_\ell^u}$ the normal space of p_ℓ^u , and by $N_{p_{\ell,j}^u}$ the normal space of $p_{\ell,j}^u$. Let us write $\xi = c_N(W)/\iota^* c_1 \mathcal{O}_{\mathcal{P}W}(-1)$. By the localisation formula, we have the equality of T -equivariant classes

$$(23) \quad \int_{\mathcal{M}^{\mathrm{can}}} \xi = \int_{\mathcal{L}_1^{\mathrm{can}}} \frac{\xi}{c_2(N_1)} + \sum_{u \in A} \int_{p_1^u} \frac{\xi}{c_3(N_{p_1^u})} + \sum_{u \in A} \sum_{j=1}^{k^u} \int_{p_{1,j}^u} \frac{\xi}{c_3(N_{p_{1,j}^u})} + \dots$$

where \dots denotes the sum of analogous integrals over $\mathcal{L}_2^{\mathrm{can}}$ and p_2^u and $p_{2,j}^u$.

Let us now evaluate each term in (23), starting with the integral over $\mathcal{L}_1^{\mathrm{can}}$. Denote by $h \in A^1(\mathbf{P}^1)$ the class of a point. Using Proposition 6.6 and Proposition 6.7 (and the notation there), we have

$$\begin{aligned} \frac{1}{c_N(W)} \int_{\mathcal{L}_1^{\mathrm{can}}} \frac{\xi}{c_2(N_1)} &= \int \frac{1}{c_1(\mathcal{O}(-1)) \cdot c_2(N_1)} \\ &= \int ((1-b)v_1 + bv_2 + (2b + r_{\mathrm{gen}} - 1)h)^{-1} (v_2 - v_1)^{-1} (v_2 - v_1 + (2-s)h)^{-1}. \end{aligned}$$

The integral is the coefficient of h in the expansion of the integrand as a power series in h . To find it, we formally differentiate with respect to h and set $h = 0$ to obtain

$$(24) \quad \frac{1}{c_N(W)} \int_{\mathcal{L}_1^{\text{can}}} \frac{\xi}{c_2(N_1)} = (2-s)((1-b)v_1 + bv_2)^{-1}(v_1 - v_2)^{-3} \\ - (2b + r_{\text{gen}} - 1)((1-b)v_1 + bv_2)^{-2}(v_1 - v_2)^{-2}.$$

The analogous integral over $\mathcal{L}_2^{\text{can}}$ is obtained by switching v_1 and v_2 .

Let us turn to the integral over p_1^u . By Proposition 6.2 (and the notation there), we have

$$(25) \quad \frac{1}{c_N(W)} \int_{p_1^u} \frac{\xi}{c_3(N_1)} = \int ((1-r^u)v_1 + r^u v_2)^{-1}(v_1 - v_2)^{-3} \\ = ((1-r^u)v_1 + r^u v_2)^{-1}(v_1 - v_2)^{-3}$$

The analogous integral over p_2^u is obtained by switching v_1 and v_2 .

Finally, let us compute the integral over $p_{1,j}^u$. By Proposition 6.4 (and the notation there), we have

$$(26) \quad \frac{1}{c_N(W)} \int_{p_{1,j}^u} \frac{\xi}{c_3(N_1)} = \int N^2((1-\lambda(j)_2)v_1 + \lambda(j)_2 v_2)^{-1}(v_1 - v_2)^{-3} \zeta_1^{-1} \eta_1^{-1} \\ = |N| \zeta_1^{-1} \eta_1^{-1} ((1-\lambda(j)_2)v_1 + \lambda(j)_2 v_2)^{-1}(v_1 - v_2)^{-3}.$$

In the last equality, we have used that $p_{1,j}^u \in \mathcal{M}^{\text{can}}$ has a stabiliser of order $|N|$, and hence the integral divides the integrand by $|N|$. The analogous integral over $p_{2,j}^u$ is obtained by switching v_1 and v_2 .

The expression in Theorem 1.3 is the sum of the contributions from (24), (25), (26), and their analogues with v_1 and v_2 switched.

7. APPLICATIONS

7.1. Orbits of elliptic fibrations. Recall that an element $(A, B) \in \text{Sym}^{4n}(V) \oplus \text{Sym}^{6n}(V)$ gives rise to an elliptic fibration

$$\pi: E \rightarrow \mathbf{P}^1$$

defined locally by the Weierstrass equation

$$y^2 = x^3 + Ax + B.$$

Given $u \in \mathbf{P}^1$, recall that r_1^u is the order of vanishing of A at u and r_2^u is the order of vanishing of B at u . We are now ready to prove Theorem 1.1, which computes the degree of the orbit closure of (A, B) .

Proof of Theorem 1.1. Let $w = (A, B) \in W = \text{Sym}^{4n}(V) \oplus \text{Sym}^{6n}(V)$ be non-zero. In the notation of Theorem 1.3, we have $b = 0$. For every $u \in \mathbf{P}^1$, the Newton polygon Λ^u has only one possible shape. It is a translated quadrant $\lambda + \mathbf{R}_{\geq 0}$ whose vertex λ is

$$\lambda = \left(\min \left(\frac{1}{4n} r_1^u, \frac{1}{6n} r_2^u \right), 0 \right) = \left(\frac{c(u)}{2n}, 0 \right).$$

In the notation of Theorem 1.3, we have $r^u = r_{\text{gen}}^u = c(u)/2n$ and $s^u = 1$.

Note that $\mathbf{P}W$ is the quotient of $W - 0$ by the \mathbf{G}_m acting by weights 2 and 3. The central $\mathbf{G}_m \subset GL V$ acts by weights $4n$ and $6n$. Therefore, the equivariant class for the first \mathbf{G}_m is obtained

from the $\mathrm{GL} V$ -equivariant class by the specialisation $v_1 = v_2 = \frac{h}{2n}$. With these substitutions, Theorem 1.1 follows from Theorem 1.3. \square

7.2. Orbits of rational self maps. Recall that elements in a Zariski open subset of $\mathrm{Hom}(V, \mathrm{Sym}^n V)$ give rise to maps $f: \mathbf{P}V \rightarrow \mathbf{P}V$ of degree n . We have an isomorphism of $\mathrm{GL} V$ -representations

$$(27) \quad \mathrm{Hom}(V, \mathrm{Sym}^n V) = \mathrm{Sym}^{n-1} V \oplus \mathrm{Sym}^{n+1} V \otimes \det V^{-1}.$$

The first projection $\mathrm{Hom}(V, \mathrm{Sym}^n V) \rightarrow \mathrm{Sym}^{n-1} V$ is the contraction. The second projection arises as the composite

$$\mathrm{Hom}(V, \mathrm{Sym}^n V) \otimes \det V = V \otimes \mathrm{Sym}^n V \rightarrow \mathrm{Sym}^{n+1} V$$

where the first map arises from the isomorphism $V^* \otimes \det V = V$ and the second map is the multiplication. The element in $\mathrm{Sym}^{n+1} V$ in the second projection defines the scheme theoretic fixed locus of f . I do not know a similar geometric interpretation of the first projection.

Fix a basis x, y of V with the dual basis x^*, y^* of V^* .

Proposition 7.1. *Let $f = x^* \otimes F(x, y) + y^* \otimes G(x, y)$, where $F, G \in \mathrm{Sym}^n V$ are polynomials of degree n . Let f correspond to $(I, J \otimes x^* \wedge y^*) \in \mathrm{Sym}^{n-1} V \otimes \mathrm{Sym}^n V \otimes \det V^{-1}$. Then, up to non-zero scalar multiples, we have*

$$I = \frac{\partial}{\partial x} F(x, y) + \frac{\partial}{\partial y} G(x, y) \text{ and } J = yF(x, y) - xG(x, y).$$

In the other direction, we have

$$\frac{1}{n+1} F = \frac{\partial J}{\partial y} + xI \text{ and } \frac{1}{n+1} G = yI - \frac{\partial J}{\partial x}.$$

Proof. It is enough to check that the construction of I and J is $\mathrm{GL}(2)$ -equivariant. We leave this to the reader. The other direction follows from the first using Euler's formula

$$x \frac{\partial *}{\partial x} + y \frac{\partial *}{\partial y} = \deg(*) \cdot *.$$

\square


Fix a non-zero $\delta \in \det V$.

Proposition 7.2. *Suppose $f \in \mathrm{Hom}(V, \mathrm{Sym}^n V)$ defines a rational map $\mathbf{P}V \rightarrow \mathbf{P}V$ of degree n and corresponds to $(I, J \otimes \delta^{-1})$ under an isomorphism (27). If J vanishes to order at least 2 at $u \in \mathbf{P}V$, then I does not vanish at u .*

Proof. Write $f = x^* \otimes F(x, y) + y^* \otimes G(x, y)$ in coordinates. Since f defines a map of degree n , the polynomials F and G have no common factor. If J vanishes to order at least 2 at u , then both partials of J vanish to order at least 1 at u . Since at least one of F or G does not vanish at u , we see from Proposition 7.1 that I cannot vanish at u . \square

We are now ready to prove Theorem 1.2, which computes the equivariant orbit class of a rational map.

Proof of Theorem 1.2. Let $f \in \mathrm{Hom}(V, \mathrm{Sym}^n V)$ define a rational map $\mathbf{P}V \rightarrow \mathbf{P}V$ of degree n . Let f correspond to $(I, J \otimes \delta^{-1})$ under an isomorphism (27). We apply Theorem 1.3, taking $A = V(J)$ to be the set of fixed points of $\mathbf{P}V \rightarrow \mathbf{P}V$. We have $b = -1/(n-1)$. By Proposition 7.2, for $u \in A$, the Newton polygon Λ^u has two possible shapes (see Figure 3). If u is a simple fixed point, then

FIGURE 3. Λ^u for a simple fixed point u (left) and a fixed point of order $j \geq 2$ (right)

its only vertex is

$$\lambda^u(0) = \frac{1}{d_2}(r_2^u + b_2, b_2) = \frac{1}{n-1}(0, -1).$$

In this case, $r_{\text{gen}}^u = 1/(n-1)$ and $r^u = 0$ and $s^u = 1$. If u is a fixed point of order $j^u \geq 2$, then the vertices of Λ^u are

$$\begin{aligned} \lambda^u(0) &= \frac{1}{d_2}(r_2^u + b_2, b_2) = \frac{1}{n-1}(j^u - 1, -1) \text{ and} \\ \lambda^u(1) &= \frac{1}{d_1}(b_1, b_1) = (0, 0). \end{aligned}$$

In this case, $r_{\text{gen}}^u = j^u/(n-1)$ and $r^u = 0$ and $s^u = j^u$. In the notation of Theorem 1.3, we have

$$\begin{aligned} F &= 2(n-1)(nv_1 - v_2)^{-1}(v_1 - v_2)^{-3} + (n+1)(nv_1 - v_2)^{-2}(v_1 - v_2)^{-2}, \text{ and} \\ G^u &= v_1^{-1}(v_1 - v_2)^{-3} - j^u(n-1)(nv_1 - v_2)^{-1}(v_1 - v_2)^{-3} - j^u(nv_1 - v_2)^{-2}(v_1 - v_2)^{-2}, \end{aligned}$$

and for a higher order fixed point u ,

$$H^u(1) = (j^u - 1)v_1^{-1}(v_1 - v_2)^{-3}.$$

Summing up and multiplying by $c_N(W)$ yields the class

$$n(n+1)(n-1)^2 \prod_{j=1}^{n-2} (jv_1 + (n-1-j)v_2) \prod_{j=1}^n ((j-1)v_1 + (n-j)v_2).$$

Note that $\mathbf{P} \text{Hom}(V, \text{Sym}^n V)$ is the quotient of $\text{Hom}(V, \text{Sym}^n V) - 0$ by \mathbf{G}_m acting by weight one. The central $\mathbf{G}_m \subset GL V$ acts by weight $n-1$. Therefore, the weight one \mathbf{G}_m equivariant class is obtained from the $GL V$ -equivariant class by specialising to $v_1 = v_2 = 1/(n-1)$. The stabiliser group $\Gamma \subset GL V$ in Theorem 1.3 and the stabiliser group $\bar{\Gamma} \subset PGL V$ in Theorem 1.2 are related by the exact sequence

$$1 \rightarrow \mu_{n-1} \rightarrow \Gamma \rightarrow \bar{\Gamma} \rightarrow 1.$$

So we must divide the class given by Theorem 1.3 by $(n-1)$. Specialising to $v_1 = v_2 = 1/(n-1)$ and dividing by $(n-1)$ gives $n(n+1)(n-1)$. \square

APPENDIX A. EQUIVARIANT CLASSES OF TORUS ORBITS

Fix an algebraic torus $T = \mathbf{G}_m^d$. We compute T -equivariant fundamental classes of orbits in T -representations. Let $M = \text{Hom}(T, \mathbf{G}_m)$ be the character group of M and set $N = \text{Hom}(M, \mathbf{Z})$. Identify $A_T = \text{Sym}(M_{\mathbf{Q}})$.

Fix a T -representation W and a $w \in W$. Write $W = \bigoplus_{i=1}^n W_i$, where W_i is one-dimensional on which T acts by the character $\chi_i \in M$. Fix a non-zero $w = (w_1, \dots, w_n) \in W$.

We recall the notion of equivariant multiplicity from [5]. Given a polyhedral rational pointed cone $\sigma \subset M_{\mathbf{R}}$, denote by $\sigma^{\vee} \subset N_{\mathbf{R}}$ the dual cone. Since σ is pointed, σ^{\vee} has non-empty interior. Given λ in the interior of σ^{\vee} , let $P_{\sigma}(\lambda)$ be the convex polytope

$$P_{\sigma}(\lambda) = \{x \in \sigma \mid \langle x, \lambda \rangle \leq 1\}.$$

There exists a unique rational function $e_{\sigma} \in \text{frac Sym}(M_{\mathbf{Q}})$ such that for every λ in the interior of σ^{\vee} , we have

$$e_{\sigma}(\lambda) = d! \cdot \text{Vol } P_{\sigma}(\lambda).$$

The function e_{σ} is called the *equivariant multiplicity* associated to σ (see [5, § 5.2]).

Let $\sigma \subset M_{\mathbf{R}}$ be the closed convex cone spanned by $\{\chi_i \mid w_i \neq 0\}$. Let $\text{Orb}(w) \subset W$ be the closure of the T -orbit of w and $[\text{Orb}(w)]$ its fundamental class in $A_T(W) = A_T$.

Theorem A.1. *In the setup above, if σ contains a line, then $[\text{Orb}(w)] = 0$. Otherwise,*

$$[\text{Orb}(w)] = e_{\sigma} \cdot c_n(W).$$

Proof. If σ contains a line, then $0 \in W$ is not in $\text{Orb}(w)$. As a result, the pull-back of $[\text{Orb}(w)]$ to $A_T(0)$ vanishes. But the pull-back $A_T(W) \rightarrow A_T(0)$ is an isomorphism, so $[\text{Orb}(w)]$ vanishes.

Assume that σ contains no line, that is, it is pointed. Let X be the affine toric variety

$$X = \text{Spec } \mathbf{k}[M \cap \sigma].$$

It is easy to check that the map $T \rightarrow W$ that sends $t \in T$ to $t \cdot w$ extends to a proper morphism $i: X \rightarrow W$. Then $[\text{Orb}(w)] = i_*[X]$.

To compute the push-forward, we use localisation [5, § 4.2 Corollary]. Let $0_X \in X$ and $0_W \in W$ be the origins. Then we have

$$[X] = e_{\sigma} \cdot [0_X],$$

and hence

$$i_*[X] = e_{\sigma} \cdot [0_W].$$

Since $[0_W] = c_n(W)[W] \in A_T(W)$, the theorem follows. \square

Example A.2. Let $T = \mathbf{G}_m^3$ act on $W = \mathbf{C}^4$ by the characters $(0, 0, 1), (0, 1, 1), (1, 0, 1)$, and $(1, 1, 1)$. Take $w = (1, 1, 1, 1)$. Let x, y, z be the standard basis vectors of $M = \text{Hom}(T, \mathbf{G}_m)$. Let $\sigma \subset M_{\mathbf{R}}$ be the cone spanned by the four characters. Given $\lambda = (a, b, c) \in \sigma^{\vee} \subset N_{\mathbf{R}}$, we compute

$$3! \cdot \text{vol}(P_{\sigma}(a, b, c)) = \frac{1}{c(b+c)(a+b+c)} + \frac{1}{c(a+c)(a+b+c)}.$$

Since $a = \langle x, \lambda \rangle$ and $b = \langle y, \lambda \rangle$ and $c = \langle z, \lambda \rangle$, the equivariant multiplicity function is

$$\begin{aligned} e_{\sigma} &= \frac{1}{z(y+z)(x+y+z)} + \frac{1}{z(x+z)(x+y+z)} \\ &= \frac{x+y+2z}{z(x+z)(y+z)(x+y+z)}. \end{aligned}$$

By Theorem A.1, the equivariant fundamental class of $\text{Orb}(w)$ is $x + y + 2z$. Indeed, in this case, $\text{Orb}(w) \subset W$ is the quadric hypersurface cut out by $w_1w_4 - w_2w_3$, a polynomial with character $(1, 1, 2)$.

REFERENCES

- [1] P. Aluffi and C. Faber. Linear orbits of d -tuples of points in \mathbb{P}^1 . *J. Reine Angew. Math.*, 445:205–220, 1993.
- [2] P. Aluffi and C. Faber. Linear orbits of smooth plane curves. *J. Algebr. Geom.*, 2(1):155–184, 1993.
- [3] P. Aluffi and C. Faber. Linear orbits of arbitrary plane curves. *Michigan Math. J.*, 48:1–37, 2000.
- [4] L. A. Borisov, L. Chen, and G. G. Smith. The orbifold Chow ring of toric Deligne-Mumford stacks. *J. Am. Math. Soc.*, 18(1):193–215, 2005.
- [5] M. Brion. Equivariant Chow groups for torus actions. *Transform. Groups*, 2(3):225–267, 1997.
- [6] A. Deopurkar and A. Patel. Orbits of linear series on the projective line, 2022. Pre-print, arXiv:2211.16603.
- [7] A. Deopurkar, A. Patel, and D. Tseng. A universal formula for counting cubic surfaces, 2021. Pre-print, arXiv:2109.12672.
- [8] F. Enriques and G. Fano. Sui gruppi continui di trasformazioni Cremoniane dello spazio. *Annali di Mat. (2)*, 26:59–98, 1898.
- [9] B. Fantechi, E. Mann, and F. Nironi. Smooth toric Deligne-Mumford stacks. *J. Reine Angew. Math.*, 648:201–244, 2010.
- [10] L. M. Fehér and R. Rimányi. Thom series of contact singularities. *Ann. Math. (2)*, 176(3):1381–1426, 2012.
- [11] L. M. Fehér, R. Rimányi, and A. Weber. Characteristic classes of orbit stratifications, the axiomatic approach. In *Schubert calculus and its applications in combinatorics and representation theory. Selected papers presented at the “International Festival in Schubert Calculus”, Guangzhou, China, November 6–10, 2017*, pages 223–249. Singapore: Springer, 2020.
- [12] W. Fulton. *Intersection theory.*, volume 2 of *Ergeb. Math. Grenzgeb., 3. Folge*. Berlin: Springer, 2nd ed. edition, 1998.
- [13] A. Kresch. Cycle groups for Artin stacks. *Invent. Math.*, 138(3):495–536, 1999.
- [14] M. Lee, A. Patel, and D. Tseng. Equivariant degenerations of plane curve orbits. *Trans. Am. Math. Soc.*, 376(10):6799–6843, 2023.
- [15] J. Milnor. Geometry and dynamics of quadratic rational maps (with an appendix by J. Milnor and Tan Lei). *Exp. Math.*, 2(1):37–83, 1993.
- [16] R. Miranda. *The basic theory of elliptic surfaces. Notes of lectures*. Pisa: ETS Editrice, 1989.
- [17] M. H. Quek. Around the motivic monodromy conjecture for non-degenerate hypersurfaces. *Manuscr. Math.*, 173(3-4):1015–1059, 2024.
- [18] M. Rees. A partial description of parameter space of rational maps of degree two. I. *Acta Math.*, 168(1-2):11–87, 1992.
- [19] R. Rimányi. Thom polynomials, symmetries and incidences of singularities. *Invent. Math.*, 143(3):499–521, 2001.
- [20] G. Segal. The topology of spaces of rational functions. *Acta Math.*, 143:39–72, 1979.
- [21] J. H. Silverman. The space of rational maps on \mathbf{P}^1 . *Duke Math. J.*, 94(1):41–77, 1998.

MATHEMATICAL SCIENCES INSTITUTE, AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA, AUSTRALIA