

## Extinction and scattering of nebular emission in Orion

WILLIAM J. HENNEY<sup>1</sup>

<sup>1</sup>*Instituto de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Apartado Postal 3-72, 58090 Morelia, Michoacán, Mexico*

### Abstract

I compare several different methods for estimating the dust extinction of diffuse emission from H II regions. Using archival data for the Orion Nebula, I show that apparent discrepancies between the different methods are powerful diagnostics of (1) emission line scattering from dusty PDRs; (2) the presence of dust layers sandwiched between two emitting gas layers; and (3) the presence of deeply embedded ionized gas that is invisible at optical and near-infrared wavelengths.

**Keywords:** Atomic physics; Radiative transfer; Photodissociation regions

### 1. INTRODUCTION


### 2. OTHER MEASURES OF EXTINCTION

#### 2.1. Diffuse interstellar bands

5781 Å is the cleanest. There is also 6283 Å, which is stronger, but it is affected by the telluric O<sub>2</sub> absorption.

#### 2.2. Forbidden metal doublets

We can use two lines that share an upper level. Small wavelength range, but high S/N and rock-solid intrinsic ratio.




**Figure 1.** Scatter plot of the infrared/optical extinction anomaly versus the radio/infrared extinction anomaly. Plot symbol color and size indicate the optical–radio extinction (larger values are darker and larger).

[O III] 4959 and 5007 Å is a 1% change in wavelength. We see about a 1% variation in the ratio.

[Ar III] 7751 and 7136 Å is a 9% change in wavelength. We see about a 15% variation in the ratio.

These give the gradient,  $dA_\lambda/d\lambda$

### REFERENCES



**Figure 2.** Three-color image of extinction in the inner Orion Nebula. Extinction derived from the optical band reddening of the Balmer decrement (4886 Å to 6563 Å) is shown in green. The infrared/optical extinction anomaly is shown in blue and the radio/infrared extinction anomaly is shown in red.