

Kinematics of the Turtle Nebula (Will's part)

William J. Henney[★]

Instituto de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Apartado Postal 3-72, 58090 Morelia, Michoacán, Mexico

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Material to be incorporated into the main paper by Teresa.

Key words: Atomic physics – Radiative transfer – Photodissociation regions




Figure 1. Proper motions derived from two HST [O III] images (F502N filter) separated by 10.45 years, using the FLCT algorithm with a Gaussian window width of 10 pixels. The key at bottom left shows a proper motion of 10 mas yr^{-1} , corresponding to 95 km s^{-1} for an assumed distance of 2 kpc.




Figure 2. As Fig. 1 but for two HST [N II] images (F658N filter). Note that the field of view is cropped slightly smaller than for [O III].

1 INTRODUCTION

2 OBSERVATIONS AND DATA REDUCTION

3 LONGSLIT SPECTROSCOPY

4 PROPER MOTIONS

Proper motions are calculated from HST WFPC2 imaging at two epochs separated by approximately 10 years, using the FLCT method (Welsch et al. 2004; Fisher & Welsch 2008).¹ Results are shown in Figures 1 and 2 for [O III] and [N II], respectively. In both cases, the images were remapped to a uniform square pixel grid at the WFC resolution of $0.1 \text{ arcsec pix}^{-1}$ before applying the algorithm. The resultant per-pixel motions between the two epochs are found to be of order 0.5 pix ($\approx 5 \text{ mas yr}^{-1}$) and these raw re-

¹ We used version 1.07 of FLCT, obtained from <http://cgem.ssl.berkeley.edu/cgi-bin/cgem/FLCT/home>, together with version 1.04 of the Python wrapper pyflct, obtained from <https://github.com/PyDL/pyflct>.

[★] w.henney@irya.unam.mx

sults were then corrected by applying a global shift to force the motion of the central star to be zero. The systematic error from the global alignment of the two epochs is estimated to be 1.5 mas yr^{-1} , which is expected to dominate the proper motion uncertainties in the brighter parts of the nebula. In fainter and more featureless regions of the nebula, the proper motions are increasingly affected by random noise, which can be seen in parts of the lobes in Figure 1.

The corrected results, as shown in the figures, can be seen to display motions that are predominantly radial from the central star. To convert the angular motions into transverse velocities, we assume a distance of 2 kpc, so that 10 mas yr^{-1} is equivalent to 95 km s^{-1} . From the [O III] images (Fig. 1), the fastest plane-of-sky motions are of order 60 km s^{-1} , and are chiefly along the NNW–SSE direction, including the projected major axis of the inner peanut shell, the NW knot, the N jet, and the end-cap of the S lobe. Motions along the perpendicular ENE–WSW direction are typically slower, of order 30 km s^{-1} . Note that proper motions are unavailable for the end cap of the N lobe since the second epoch HST image does not cover this region.

The [N II] images (Fig. 2) show a similar expansion pattern for the features that are visible in both lines. Remarkably low plane-of-sky velocities of $\leq 15 \text{ km s}^{-1}$ are seen for the [N II]-bright knot complexes immediately north and west of the central star.

5 KINEMATIC COMPONENTS

5.1 Inner shells

5.2 Knot complexes

5.3 Intermediate shell

5.4 Outer lobes

5.5 Haloes

6 DISCUSSION

6.1 Flow axes

6.2 Dynamical ages

References

- Fisher G. H., Welsch B. T., 2008, in Howe R., Komm R. W., Balasubramaniam K. S., Petrie G. J. D., eds, *Astronomical Society of the Pacific Conference Series Vol. 383, Subsurface and Atmospheric Influences on Solar Activity*. San Francisco, p. 373
- Welsch B. T., Fisher G. H., Abbett W. P., Regnier S., 2004, *ApJ*, **610**, 1148

This paper has been typeset from a \LaTeX file prepared by the author.




Figure 3. Reconstructed velocity channel maps from the [N II] slit spectra, showing the red-shifted (left panel) and blue-shifted (right panel) knot complexes. Note that channel maps have not been spatially interpolated, so that the individual slit positions can be seen. Each color image is constructed from 3 channels, each of width 20 km s^{-1} , as indicated on the figure. All velocities are with respect to the nominal heliocentric systemic velocity of -40 km s^{-1} .




Figure 4. Velocity components and systems in the outer lobes.