

# $e^+e^-$ Scattering to Hadrons Notes

---

**Dirk Hornung<sup>a</sup>**

*<sup>a</sup>Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology,  
Campus UAB, 08193 Bellaterra (Barcelona) Spain*

*E-mail:* [dirkhornung91@gmail.com](mailto:dirkhornung91@gmail.com)

ABSTRACT: Abstract...

KEYWORDS: Keywords...

---


## Contents

<b>1</b>	<b>Tau Decays</b>	<b>2</b>
<b>2</b>	<b>Theory</b>	<b>3</b>
2.1	The theoretical description of the correlator $\Pi(s)$	4
2.1.1	The perturbative contribution	4
<b>3</b>	<b>Theory</b>	<b>6</b>
3.1	Theory equals Experiment	9
3.2	Transform Two Point Function to Adler Function via Complex Contour Integration by Parts	9
3.3	Pearson's $\chi^2$ test	10
<b>4</b>	<b>Adler function Contributions</b>	<b>10</b>
4.1	Dimension Zero	10
<b>5</b>	<b>The Experimental Spectral Moments</b>	<b>10</b>
<b>A</b>	<b>Derivation of the <math>\tau</math>-decay ratio</b>	<b>12</b>
<b>B</b>	<b>Coefficients</b>	<b>14</b>
B.1	$\beta$ function	15
B.2	$\gamma$ -function	15
B.3	Adler function	15
<b>C</b>	<b>Numerical Analysis</b>	<b>15</b>
C.1	Newton-Raphson Method	15
C.2	Newton-Cotes	15
C.3	Gauss Quadrature	15
C.4	Discrepancies between Matthias and my Code	15

---

## 1 Tau Decays

The  $\tau$ -lepton is the only lepton heavy enough to decay into hadrons. It is an ideal candidate to perform a QCD analysis at low to intermediate energies<sup>1</sup>, extracting and probing Standard model parameters like the strong coupling  $\alpha_s$ . Due to its W-Boson mediator it allows a study of the  $q\bar{q}$  vector and axial vector current, whereas  $e^+e^- \rightarrow \gamma^* \rightarrow (\text{hadrons})$  scattering only covers the vector ones. For the tree level of the SM the tau decays into a  $\nu_\tau$  neutrino and a W-Boson, which then decays into either  $(\nu_\mu, \mu)$  and  $(\nu_e, e)$  lepton pairs or  $(\bar{u}, d)$ , and  $(\bar{u}, s)$  quark pairs, as illustrated in Figure 1. Other quark pair generation is forbidden either due to the  $\tau$ -mass of 1776.86 MeV [4] or the confinement of weak isospin ( $T_3$ )<sup>2</sup> Due to color confinement<sup>3</sup> the detected quark pairs are hadrons (pions,  $\rho$  mesons, kaons and others).



**Figure 1.** Standard Model  $\tau$  decays.

The data is gathered via the ALEPH experiment in CERN and expressed through the ratio

$$R_\tau = \frac{\mathcal{B}_\tau}{\mathcal{B}_{e^-}} = \frac{\Gamma(\tau^- \rightarrow \text{hadrons}^- \nu_\tau)}{\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)}, \quad (1.1)$$

which can be connected to the vacuum polarization  $\Pi(s)$  of the W-Boson<sup>4</sup>.  $\mathcal{B}_\tau$  and  $\mathcal{B}_{e^-}$  are the branching fractions of the non strange hadronic and electronic decays and the  $\Gamma$ s the respective decay rates.

Our aim is to match the experimental data with a theory model in a computer program to analyse and extract parameters of the SM. Therefore we will start connecting the experimental data with quantum field theory. Then describe the needed theory and experimental sides in detail. And lastly discuss the results.

<sup>1</sup>At high energies we have no valid theory because  $\alpha_s \gg 1$  for high-energies.

<sup>2</sup>E.g. the  $(\bar{d}, u)$  pair would lead to  $(\tau \rightarrow \nu_\tau + \bar{d} + u)$ , which represents an inequality in terms of weak isospin conservation ( $-1/2 \neq +1/2 + 1/2 - 1/2$ ).

<sup>3</sup>Quarks have never been seen as single particle, but only in compounds as mesons or baryons.

<sup>4</sup>Also referred to as the weak current two-point correlation function.

## 2 Theory

Starting point of our analysis will be the definition of the vector and axial vector QCD correlator

$$\Pi_{\mu\nu,ij}^{V/A}(p) \equiv i \int dx e^{ipx} \langle \Omega | T \{ J_{\mu,ij}^{V/A}(x) J_{\nu,ij}^{V/A}(0)^\dagger \} | \Omega \rangle, \quad (2.1)$$

with  $|\Omega\rangle$  being the physical vacuum and  $i, j$  representing the light quark flavours: up, down and strange<sup>5</sup>. The V and A currents are given by

$$J_{\mu,ij}^{V/A}(x) = [\bar{q}_j \gamma_\mu (\gamma_5) q_i](x). \quad (2.2)$$

The before introduced correlator can be decomposed in several ways starting by the standard longitudinal and transversal decomposition (omitting the flavour indices  $i, j$  from now on, if not necessary):

$$\begin{aligned} \Pi_{\mu\nu}(p)^{V/A} &= (p_\mu p_\nu - g_{\mu\nu} p^2) \Pi^{V/A,(1)}(p^2) + p_\mu p_\nu \Pi^{V/A,(0)}(p^2) \\ &= (p_\mu p_\nu - g_{\mu\nu} p^2) \Pi^{V/A,(1+0)}(p^2) + g_{\mu\nu} p^2 \Pi^{V/A,(0)}(p^2) \end{aligned} \quad (2.3)$$

, where the upper indice (1) denotes the transversal contribution and (0) the longitudinal one. Besides the Lorentz decomposition, the currents contributin to hadronic  $\tau$  decays can also be decomposed according to their light quark flavour content:

$$\Pi_{V/A}^{(J)}(s) = |V_{ud}|^2 \left[ \Pi_{ud}^{V,(J)}(s) + \Pi_{ud}^{A,(J)}(s) \right] + |V_{us}|^2 \left[ \Pi_{us}^{V,(J)}(s) + \Pi_{us}^{A,(J)}(s) \right], \quad (2.4)$$

with  $V_{ij}$  being the corresponding elemtns of the Cabibbo-Kobayashi-Maskawas (CKM) quark-mixing matrix. Another useful decomposition of the V (A) correlator is the separation of vector (axial vector) and scalar (pseudoscalar) components

$$\begin{aligned} \Pi_{\mu\nu}^{V/A}(p) &= (p_\mu p_\nu - g_{\mu\nu} p^2) \Pi^{V/A}(p^2) \\ &+ \frac{g_{\mu\nu}}{p^2} (m_i \mp m_j)^2 \Pi^{S,P}(p^2) + g_{\mu\nu} \frac{m_i \mp m_j}{p^2} [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle]. \end{aligned} \quad (2.5)$$

This implies the following relations between the Lorentz scalar function of Eqs. (2.3) and (2.5)

$$s^2 \Pi^{V/A,(0)}(p^2) = (m_i \mp m_j) \Pi^{S,P}(p^2) + (m_i \mp m_j) [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle], \quad (2.6)$$

$$\Pi^{V/A,(1+0)}(p^2) \equiv \Pi^{V/A,(1)}(p^2) + \Pi^{V/A,(0)}(p^2) = \Pi^{V/A}(p^2). \quad (2.7)$$

Consequently, in the chiral limit ( $m_i \rightarrow 0$ ) the longitudinal (Spin-0) component, proportional to the quark masses, vanishes. With these definitions one can then derive the inclusive decay ratio:

$$R_\tau = 12\pi S_{EW} \int_0^{m_\tau^2} \frac{ds}{m_\tau^2} \left( 1 - \frac{s}{m_\tau^2} \right)^2 \left[ \left( 1 + 2 \frac{s}{m_\tau^2} \right) \text{Im} \Pi^{(T)}(s) + \text{Im} \Pi^{(L)}(s) \right], \quad (2.8)$$

---

<sup>5</sup>As mentioned before, due to it's mass the  $\tau$ -lepton cannot decay into the heavier quarks top, bottom and charm.

where  $S_{EW}$  is an electroweak correction factor.

With the help of the QCD Sum Rules, more specifically the FESR we can then rewrite the tau decay ratio (2.8) into

$$R_\tau = 6\pi i S_{EW} \oint_{|s|=m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \left[ \left(1 + 2\frac{s}{m_\tau^2}\right) \text{Im} \Pi^{(T)}(s) + \text{Im} \Pi^{(L)}(s) \right]. \quad (2.9)$$

Normally, using the FESR, we need to pay attention to the integral, which runs along the real axis where the perturbation theory is not valid anymore. This problems close to the real axis are believed to be caused by duality violations. The normal procedure would be to introduce so called pinched weight function, which vanishes at the real axis. However, in the case of the  $\tau$ -decay ratio we naturally derived a pinched weight function (App. ??)

$$w_\tau(s) = \left(1 - \frac{s}{m_\tau^2}\right) \left(1 + 2\frac{s}{m_\tau^2}\right), \quad (2.10)$$

which introduce a double zero at  $s = m_\tau^2$ . The sum rules using pinched weight functions are also referred to as pinched weighted finite energy sum rules or pFESR. It is convenient to rewrite Eq. (??) as

$$R_\tau = 6\pi i S_{EW} \oint_{|s|=m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \left[ \left(1 + 2\frac{s}{m_\tau^2}\right) \text{Im} \Pi^{(T+L)}(s) - \left(2\frac{s}{m_\tau^2}\right) \text{Im} \Pi^{(L)}(s) \right], \quad (2.11)$$

where we simply reordered the spin composition

$$\Pi^{(T+L)}(s) = \Pi^{(T)}(s) + \Pi^{(L)}(s). \quad (2.12)$$

Unfortunately the two-points functions  $\Pi^{(J)}$  are not RG invariant and contains scale and scheme dependent subtractions constants. Noticing that the decay ratio integral (Eq. (??)) is RG invariant we can define the so called Adler function, which by itself is RG invariant

$$D^{(L+T)}(s) \equiv -s \frac{d}{ds} \left( \Pi^{(L+T)}(s) \right), \quad D^{(L)}(s) = \frac{s}{m_\tau^2} \frac{d}{ds} \left( s \Pi^{(L)}(s) \right). \quad (2.13)$$

Making use of integration by parts yields

$$R_\tau = -\pi i \oint_{|s|=m_\tau^2} \frac{ds}{s} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left[ 3 \left(1 + \frac{s}{m_\tau^2}\right) D^{(L+T)}(s) + 4 D^{(L)}(s) \right], \quad (2.14)$$

which can be calculated within QCD.

## 2.1 The theoretical description of the correlator $\Pi(s)$

### 2.1.1 The perturbative contribution

We will perform our analysis in the chiral limit, which cancels the longitudinal part of the two point function  $\Pi^{(0)}(s) = 0$ <sup>6</sup>. This leaves us with with the  $\Pi_V^{(1+0)}(s)$  component, which can generally be expressed as [1]

$$\Pi_V^{(1+0)}(s) = -\frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_\mu^n \sum_{k=0}^{n+1} c_{n,k} L^k, \quad (2.15)$$

---

<sup>6</sup>The longitudinal part is proportional to the quark masses. Hence zero in the limit of zero quark masses

with

$$L \equiv \log \left( \frac{-s}{\mu} \right) \quad \text{and} \quad a_\mu \equiv a(\mu^2) = \frac{a_s(\mu)}{\pi} \quad (2.16)$$

and  $\mu$  being the renormalisation scale. With the help of Eq. (2.13) we can rewrite the former expression into terms of the RG invariant Adler function

$$D_V^{(1+0)}(s) = \frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_\mu^n \sum_{k=1}^{n+1} k c_{n,k} L^{k-1} \quad (2.17)$$

, where the coefficients  $c_{n,k}$  are given in App. B.

### 3 Theory

$\Pi(q^2)$  itself is not a physical quantity. However, we can define two physical quantities out of the two-point correlator. Setting  $s = q^2$ , the spectral function  $\rho(s)$  and the Adler function  $D(s)$  are given by

$$\rho(s) \equiv \frac{1}{\pi} \text{Im} \Pi(s) \quad D(s) \equiv -s \frac{d\Pi(s)}{ds} \quad (3.1)$$

Using Kramers-Kronig relation we can connect the theoretical, non physical quantity  $\Pi(s)$  with the experimental measurable spectral function  $\rho(s)$ . With Kramer-Kronic relation we can display the real part of a function as its imaginary part and vice verse. Let  $f(x) = \text{Re}f(x) + i\text{Im}f(x)$  be a complex function of the variable  $s$ . Suppose this function is analytic in the closed upper half-plane of  $s$  and vanishes like  $1/|s|$  as  $|s| \rightarrow \infty$ . Then the real part of  $f$  can be given as integral over the complex part of  $f$

$$\text{Re}f(s) = \frac{1}{i\pi} P \int_{-\infty}^{\infty} \frac{\text{Im}f(s')}{s' - s} ds' \quad (3.2)$$

To derive the Kramers-Kronig relation we start from Cauchy's theorem. Given an analytic function  $f(s)$  in the closed upper half plane. The function  $s' \rightarrow f(s')/(s' - s)$ , where  $s$  and  $s'$  are real values, is also be analytic in the upper half of the plane. The Cauchy's residum theorem then consequently states that

$$\oint \frac{f(s')}{s' - s} ds' = 0. \quad (3.3)$$

**include graph** Following the contour integral in **figure above** we are left with

$$\begin{aligned} \oint \frac{f(s')}{s' - s} ds' &= \int_{-R}^{-\epsilon} \frac{f(s')}{s' - s} ds' + \int_{\gamma_{\epsilon+}} \frac{f(s')}{s' - s} ds' + \int_{\epsilon}^R \frac{f(s')}{s' - s} ds' + \int_{\gamma_R}^+ \frac{f(s')}{s' - s} ds' \\ &= \oint_{-\infty}^{\infty} \frac{f(s')}{s' - s} ds', \end{aligned} \quad (3.4)$$

where the integral along upper semicircle vanishes due to the fact, that in the limit  $R \rightarrow \infty$  the function  $f(s')$  vanishes faster than  $1/|s'|$ . Furthermore we passed the half-circle to zero. **why possible?** The real line integral we are left with can be solved using the Sokhotski-Plemelj theorem. Starting by the integral

$$\begin{aligned} \lim_{\epsilon \rightarrow 0^+} \int_a^b \frac{f(x)}{x + i\epsilon} dx &= \lim_{\epsilon \rightarrow 0^+} \int_a^b \frac{x - i\epsilon}{(x + i\epsilon)(x - i\epsilon)} \\ &= -i\pi \lim_{\epsilon \rightarrow 0^+} \int_a^b \frac{f(x)}{\pi(x^2 + \epsilon^2)} + \lim_{\epsilon \rightarrow 0^+} \int_a^b \frac{x^2}{x^2 + \epsilon^2} \frac{f(x)}{x} dx, \end{aligned} \quad (3.5)$$

where the first term in the second expression is the nasent delta function and therefore approaches a dirac delta function and turns into  $-i\pi f(0)$ . The second term of the second identity is approaches 1 for  $|x| \ll \epsilon$  and 0 for  $|x| \gg \epsilon$ . Additionally it is totally symmetrix about 0 (which is important for the Cauchy principal value) and turns in the limit into Cauchys principal value integral given by

$$P \int_L f(x) dx = \lim_{\epsilon \rightarrow 0} \int_{L(\epsilon)} f(x) dx \quad (3.6)$$

Consequently we get the Sokhotski-Plemelj theorem

$$\lim_{\epsilon \rightarrow 0^+} \int_a^b \frac{f(x)}{x - i\epsilon} dx = -i\pi f(x). \quad (3.7)$$

Finally we can we just have to plug in the theorem and change the formula

$$\begin{aligned} 0 &= \oint \frac{f(s')}{s' - s} ds' = P \int_{-\infty}^{\infty} \frac{f(s')}{s' - s} - i\pi f(s) \\ \Rightarrow f(s) &= \frac{P}{i\pi} \int_{-\infty}^{\infty} \frac{f(s')}{s' - s}. \end{aligned} \quad (3.8)$$

Remembering the definition of the spectral function  $\rho(s)$  we can now express the two-point correlator  $\Pi(s)$  as

$$\Pi(s) = \int_0^{\infty} \frac{\rho(s')}{s' - s - i0} ds' + P(s) \quad (3.9)$$

what happen with the Cauchy principal value (symmetry about 0?), what is P(s) (carries required scale dependence???) think about this once again, at least one more step to  $\rho$  function The starting point for a QCD analysis of the  $e^+e^-$  scattering is the finite energy sum rule (FESR) cite?

$$\frac{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow q\bar{q})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} \equiv R_q(s) = \int_0^{s_0} w(s)\rho(s)ds = -\frac{1}{2\pi i} \oint_{|s|=s_0} w(s)\Pi(s)ds \quad (3.10)$$

how are now 1.15 and 1.16 connected? Why can we write  $R_q$  as the integral from 0 to  $s_0$  over the  $w(s)$  times  $p(s)$ ? In the massless case the vector correlation function  $\Pi(s)$  can be written as follows

$$\Pi(s) = -\frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_{\mu}^n \sum_{k=0}^{n+1} c_{nk}(L)^k, \quad \text{where} \quad L \equiv \ln \frac{-s}{\mu^2} \quad (3.11)$$

and  $a_{\mu} = a = \alpha_s/\pi$ . Consequently the perturbative expansion of the Adler function  $D(s)$  can be writte as

$$D(s) = \frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_{\mu}^n \sum_{k=0}^{n+1} c_{nk} k(L)^{k-1}. \quad (3.12)$$

The spectral function  $\rho(s)$  and the Adler function  $D(s)$  are physical quantities, so that they have to follow the homogenous renormalisation group equation (RGE)

$$-\mu \frac{d}{d\mu} \left\{ \frac{D(s)}{\rho(s)} \right\} = \left[ 2 \frac{\partial}{\partial L} + \beta(a) \frac{\partial}{\partial a} \right] \left\{ \frac{D(s)}{\rho(s)} \right\} = 0, \quad (3.13)$$

where we simply used the known form of the homogenous RGE, substitutet  $\mu \rightarrow L$  and multiplied by -1. The known RGE is given by

$$\left\{ \mu \frac{\partial}{\partial \mu} - \beta(a_s) \frac{\partial}{\partial a_s} - \gamma(a_s) m \frac{\partial}{\partial m} \right\} R(q, a_s, m) = 0, \quad (3.14)$$



where  $R(q, a_s, m)$  represents an arbitrary physical quantity. In our case these physical quantity is represented by the Adler or the Spectral function. As we neglected the mass the partial derivative of the mass  $\partial/\partial m$  vanishes. Furthermore the substitution is given by

$$L = \ln \frac{-s}{\mu^2} = \ln(-s) - 2 \ln(\mu) \quad \Rightarrow \quad \frac{\partial L}{\partial \mu} = -\frac{2}{\mu}. \quad (3.15)$$

Inserting the substitution in eq. 3.14 and multiplying with -1 then yields eq. 3.13.

In the RGE we we have used the  $\beta$ -function, which is defined as

$$-\mu \frac{da}{d\mu} \equiv \beta(a) = \beta_1 a^2 + \beta_2 a^3 + \beta_3 a^4 + \beta_4 a^5 + \dots \quad (3.16)$$

and known numerically for  $N_c = 3$  in the  $\bar{M}\bar{S}$ -scheme [cite, jamin 27](#) up to fourth coefficient by [cite, jamin 28-30](#)

$$\begin{aligned} \beta_1 &= \frac{11}{2} - \frac{1}{3}N_f, & \beta_2 &= \frac{51}{4} - \frac{19}{12}N_f, & \beta_3 &= \frac{2857}{64} - \frac{5033}{576}N_f + \frac{325}{1728}N_f^2, \\ \beta_4 &= \frac{149753}{768} + \frac{891}{32}\zeta_3 - \left( \frac{1078361}{20736} + \frac{1627}{864}\zeta_3 \right) N_f + \left( \frac{50065}{20736} + \frac{809}{1296}\zeta_3 \right) N_f^2 + \frac{1093}{93312}N_f^3. \end{aligned} \quad (3.17)$$

Now we want to have a closer look to the coefficients  $c_{nk}$  used in the perturbative expansion of the Adler and the Spectral function. Obviously the coefficients  $c_{n0}$  have to vanish in both, the Spectral function (it is purely imaginary and the coefficient is real) and the Adler function (it has a factor  $k$ ), thus the  $c_{n0}$  are unphysically and do not appear in measurable quantities. Furthermore the homogenous RGE (eq. ??) puts constraints on the coefficients  $c_{nk}$ . Considering  $c_{n1}$  to be independent all other coefficients  $c_{nk}$  with  $k = 2, \dots, n+1$  can be expressed with  $c_{n1}$  and the  $\beta$ -function up to order  $\alpha_s^2$ . Hence the RG constraints lead to

$$\begin{aligned} c_{22} &= -\frac{\beta_1}{4}c_{11}, & c_{33} &= \frac{\beta_1^2}{12}c_{11}, & c_{32} &= -\frac{1}{4}(\beta_2 c_{11} + 2\beta_1 c_{21}), \\ c_{44} &= -\frac{\beta_1^3}{32}c_{11}, & c_{43} &= \frac{\beta_1}{24}(5\beta_2 c_{11} + 6\beta_1 c_{21}), & c_{42} &= -\frac{1}{4}(\beta_3 c_{11} + 2\beta_2 c_{21} + 3\beta_1 c_{31}). \end{aligned} \quad (3.18)$$

Additionally the coefficients,  $c_{n,n+1} = 0$  for  $n \geq 1$ . The independent coefficients  $c_{n1}$  are known analytically up to order  $\alpha_s^3$  [cite, jamin 31,32](#) and at  $N_c = 3$  in the  $\bar{M}\bar{S}$ -scheme take the following values

$$\begin{aligned} c_{01} &= c_{11} = 1, & c_{21} &= \frac{365}{24} - 11\zeta_3 - \left( \frac{11}{12} - \frac{2}{3}\zeta_3 \right) N_f, \\ c_{31} &= \frac{87029}{288} - \frac{1103}{4}\zeta_3 + \frac{275}{6}\zeta_5 - \left( \frac{7847}{216} - \frac{262}{9}\zeta_3 + \frac{25}{9}\zeta_5 \right) N_f + \left( \frac{151}{162} - \frac{19}{27}\zeta_3 \right) N_f^2. \end{aligned} \quad (3.19)$$

### 3.1 Theory equals Experiment

$$\begin{aligned}
\int_{C_2} ds \Pi(s) &= - \int_{C_1} ds \Pi(s) \\
&= - \int_{0+i\epsilon}^{s_0+i\epsilon} \Pi(s) ds - \int_{0-i\epsilon}^{s_0-i\epsilon} \Pi(s) ds \\
&= - \int_0^{s_0} \Pi(s' + i\epsilon) ds' - \int_0^{s_0} \Pi(s' - i\epsilon) ds' \\
&= - \int_0^{s_0} (\Pi(s' + i\epsilon) - \Pi(s' - i\epsilon)) ds' \\
&= - \int_0^{s_0} 2i \operatorname{Im} \Pi(s' + i\epsilon) ds'
\end{aligned} \tag{3.20}$$

$$- \int_0^{s_0} 2 \operatorname{Im} \Pi(s) ds = \oint_{|s|=s_0} \Pi(s) ds \tag{3.21}$$

$$- \int_0^{s_0} \rho_{exp}(s) ds = - \frac{1}{2\pi i} \oint_{|s|=s_0} \Pi(s) ds \tag{3.22}$$

### 3.2 Transform Two Point Function to Adler Function via Complex Contour Integration by Parts

Remember the definition of the **Adler function**:

$$D \equiv -s \frac{d}{ds} \Pi(s). \tag{3.23}$$

Having a look at the **Fundamental theorem of calculus** we can write for a path  $\gamma : [0, 1] \rightarrow \mathbb{C}$

$$\int_{\gamma} f(z) dz = F(z_1) - F(z_0) \quad \text{or} \quad f(\gamma(1)) - f(\gamma(0)) = \int_{\gamma} f'(z) dz. \tag{3.24}$$

Using the definition of the contour integration

$$\int_{\gamma} f(z) dz = \int_0^1 f(\gamma(t)) \gamma'(t) dt = \int_0^1 (f \circ \gamma)'(t) dt \tag{3.25}$$

the **complex integration by parts** is given by

$$\begin{aligned}
\int_{\gamma} f'(z) g(z) dz &= \int_0^1 f'(\gamma(t)) g(\gamma(t)) \gamma'(t) dt \\
&= \int_0^1 (f \circ \gamma)'(t) (g \circ \gamma)(t) dt \\
&= [(f \circ \gamma)(t) (g \circ \gamma)(t)]_0^1 - \int_0^1 (f \circ \gamma)(t) (g \circ \gamma)'(t) dt \\
&= - \int_0^1 f(\gamma(t)) g'(\gamma(t)) dt
\end{aligned} \tag{3.26}$$

for a closed path (the boundary term vanished for same start and end point). Applying **complex integration by parts** let's us substitute the **Two-Point function** by the **Adler function**

$$\oint_{|s|=s_0} w(s) \Pi(s) = - \oint_{|s|=s_0} \hat{w}(s) \frac{d\Pi(s)}{ds} = \oint_{|s|=s_0} \frac{\hat{w}(s)}{s} D(s). \tag{3.27}$$

### 3.3 Pearson's $\chi^2$ test

$$\chi^2(\alpha) = [I_i^{exp} - I_i^{th}(\alpha)] C_{ij}^{-1} [I_j^{exp} - I_j^{th}(\alpha)] \quad (3.28)$$

with

$$\begin{aligned} I_i^{exp} &= - \int_0^{s_0} \rho_{exp}(s) ds \\ I_i^{th}(\alpha) &= \oint_{|s|=s_0} \Pi(s) ds \\ C &= \begin{pmatrix} \sigma_1^2 & \rho_{12}\sigma_1\sigma_2 & \rho_{13}\sigma_1\sigma_3 & \cdots \\ \rho_{21}\sigma_1\sigma_2 & \sigma_2^2 & \rho_{23}\sigma_2\sigma_3 & \cdots \\ \rho_{31}\sigma_1\sigma_2 & \rho_{32}\sigma_2\sigma_3 & \sigma_3^2 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \end{aligned} \quad (3.29)$$

## 4 Adler function Contributions

$$\beta_1 = \frac{11}{2} - \frac{1}{3}N_f, \quad \beta_2 = \frac{51}{4} - \frac{19}{12}N_f, \quad \beta_3 = \frac{2857}{64} - \frac{5033}{576}N_f + \frac{325}{1728}N_f^2, \quad (4.1)$$

$$\beta_4 = \frac{149753}{768} + \frac{891}{32}\zeta_3 - \left( \frac{1078361}{20736} + \frac{1627}{864}\zeta_3 \right) N_f + \left( \frac{50065}{20736} + \frac{809}{1296}\zeta_3 \right) N_f^2 + \frac{1093}{93312}N_f^3 \quad (4.2)$$

### 4.1 Dimension Zero

$$D(s) = \frac{1}{4\pi^2} \sum_{n=0} \tilde{K}_n(\mu) a^n (-\mu s^2) \quad (4.3)$$

$$\tilde{K}_0(\mu) = K_0, \quad \tilde{K}_1(\mu) = K_1, \quad \tilde{K}_2(\mu) = K_2 - \beta_1 K_1 \ln(\mu) \quad (4.4)$$

$$\tilde{K}_3(\mu) = K_3 - [\beta_2 K_1 + 2\beta_1 K_2] \ln \mu + \beta_1^2 K_1 \ln^2 \mu \quad (4.5)$$

$$K_0 = K_1 = 1, \quad K_2 = \frac{295}{24} - 9\zeta_3, \quad K_3 = \frac{58057}{288} - \frac{779}{4}\zeta_3 - \frac{75}{2}\zeta_5 \quad (4.6)$$

## 5 The Experimental Spectral Moments

The general form of the spectral moment can be defined as:

$$I_{exp} = \int_0^{s_0} \frac{ds}{s_0} \omega_\tau \left( \frac{s}{s_0} \right) \rho_{exp}(s). \quad (5.1)$$

This is the experimental side we need to build our  $\chi^2$  method to fit our parameters to the data. The spectral-function of the inclusive Hadron  $\tau$  decays is connected to the total decay rate

$$R_\tau = 12\pi^2 \int_0^{s_\tau} \frac{ds}{s_\tau} \left( 1 - \frac{s}{s_\tau} \right)^2 \left[ \left( 1 + 2\frac{s}{s_\tau} \right) \rho_{V+A}^{(1+0)} - 2\frac{s}{s_\tau} \rho_{V+A}^{(0)}(s) \right] \quad (5.2)$$

where  $s_\tau = m_\tau^2$  and  $\rho^{(1)}(s)$  as well as  $\rho^{(0)}(s)$  being the transversal and the longitudinal spectral functions, which are related to the theoretical vector/axialvector (V/A) correlators and implicitly contain the CKM ( $|V_{ud}|^2$ ) and electro-weak ( $S_{EW}$ ) correction factors.

The total decay rate  $R$  is measured within the experiments and is provided by the *Aleph group* as a Fortran subroutine, outputting the binned, normalized invariant mass squared distribution

$$\int_0^{m_\tau^2} ds \text{sfm2}_{V/A}(s) = \int_0^{m_\tau^2} ds B_i \left( \frac{dN_i}{N_i ds} \right) \times 100 = B_i \times 100 = R_{\tau,i} B_e \times 100 \quad (5.3)$$

of the non-strange hadronic tau decays for the respective channel  $i=\{\text{V-channel, the A-channel, the combined V+A and V-A channels}\}$ .

Knowing that the contribution of the longitudinal spectral function is neglect-able<sup>7</sup> we can rewrite the total decay width into

$$R_{\tau,V/A} = 12\pi^2 \int_0^{s_\tau} \frac{ds}{s_\tau} \left( 1 - \frac{s}{s_\tau} \right)^2 \left( 1 + 2 \frac{s}{s_\tau} \right) \rho_{V+A}^{(1+0)}, \quad (5.4)$$

which we can connect to the binned experimental data

$$\text{sfm2}(sbin) = 100 \times \frac{12\pi^2 B_e}{s_\tau} \underbrace{\int_{sbin-0.5dsbin}^{sbin+0.5dsbin} ds \left( 1 - \frac{s}{s_\tau} \right)^2 \left( 1 + 2 \frac{s}{s_\tau} \right) \rho_{exp}(sbin)}_{\omega_\tau} \quad (5.5)$$

$$= 100 \times \frac{12\pi^2 B_e}{s_\tau} \omega_\tau \rho_{exp}(sbin), \quad (5.6)$$

where  $sbin$  denotes the center of the data bin with a given bin width  $dsbin$ <sup>8</sup>. The needed spectral function is therefore given by:

$$\rho_{exp} = \frac{s_\tau}{100 \times 12\pi^2 B_e \omega_\tau} \text{sfm2}(sbin) \quad (5.7)$$

As we related now the data (sfm2) to the spectral function ( $\rho_{exp}$ ) we can write out the final expression used for the weighted experimental spectral moments

$$I_{exp,\omega} = \sum_{sbin \leq s_0} \left( \int_{sbin-0.5dsbin}^{sbin+0.5dsbin} ds \omega \left( \frac{s}{s_0} \right) \rho_{exp}(sbin) \right) \quad (5.8)$$

$$i\mathcal{M} = (-i)^2 \frac{g^2}{2} \bar{u}(p_\tau) \gamma_\mu \left( \frac{1 - \gamma^5}{2} \right) u(p_\nu) \frac{-i}{M_W^2} \left( \frac{i}{4} \frac{g^2}{2} \Pi^{\mu\nu}(q) \right) \times \frac{-i}{M_W^2} \quad (5.9)$$

<sup>7</sup>Apart from the pion-pion pole contribution, which we'll subtract from the data.

<sup>8</sup>The bin widths given in the Aleph data are not equidistant, but given

## A Derivation of the $\tau$ -decay ratio

One of the central formulas of our QCD  $\tau$ -decay analysis is the  $\tau$ -decay ratio

$$R_\tau = 12\pi S_{EW} \int_0^{m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \left[ \left(1 + 2\frac{s}{m_\tau^2}\right) \text{Im} \Pi^{(1)}(s) + \text{Im} \Pi^{(0)}(s) \right]. \quad (\text{A.1})$$

It was originally derived by Tsai in 1971 [8] and will be probed in the following the guide of [6].

First we need to remember the Lorentz-decomposition of the QCD correlator

$$\Pi_{\mu\nu}(q) = (q_\mu q_\nu - q^2 g_{\mu\nu}) \Pi^{(T)}(q^2) + q^\mu q^\nu \Pi^{(L)}(q^2). \quad (\text{A.2})$$

The desired decay-ratio can be expressed as

$$R_\tau = \frac{\Gamma(\tau \rightarrow \text{hadrons})}{\Gamma(\tau \rightarrow e\nu_\tau \bar{\nu}_e)} \quad (\text{A.3})$$

so we need to make use of the general decay-rate defined as

$$\Gamma = \frac{1}{2M_A} \int \left( \Pi_f \frac{d^3p}{(2\pi)^3} \frac{1}{2E_f} \right) |\mathcal{M}(A \rightarrow \sum_f f)|^2 (2\pi)^4 \delta(p_A - \sum_f p_f), \quad (\text{A.4})$$

where  $|\mathcal{M}|^2$  is the squared decay matrix element. As we do not know all the corrections of quark interactions within the  $\tau$ -decay we want to make use of the optical theorem. The optical theorem states that the before mentioned squared decay matrix element  $|\mathcal{M}|^2$ , as in Fig. A b), is twice the imaginary part of two times the diagram, like in Fig. A a). Consequently we will calculate the squared decay matrix element of a) in A and use its imaginary part, dividing by a factor of two to get to the needed  $\tau$  decay rate. Within the  $\tau$ -decay rate we furthermore have to sum over all possible hadronic final states and integrate over their phase space factors to deal with the momentum- $\delta$ , leaving us with an integration over the  $\tau$ -neutrino.

Starting with the diagrammatical evaluation of Fig. A a), making use of the Feynman rules, we get

$$\begin{aligned} i\mathcal{M} &= (-i)^2 \frac{g^2}{2} \bar{u}(p_\tau) \gamma_\nu \left( \frac{1 - \gamma^5}{2} \right) u(p_{\nu_\tau}) \frac{-i}{M_W^2} \left( \frac{i}{4} \frac{g^2}{2} \Pi^{\mu\nu}(q) \right) \\ &\times \frac{-i}{M_W^2} \bar{(p_{\nu_\tau})} \gamma_\nu \left( \frac{1 - \gamma^5}{2} \right) u(p_\tau). \end{aligned} \quad (\text{A.5})$$


Focusing on the spinors, while summing over the spins we can evaluate the traces

$$\frac{1}{2} \sum_{\text{spins}} \bar{u}(p_\tau) \gamma_\mu \left( \frac{1 - \gamma^5}{2} \right) u(p_{\nu_\tau}) \bar{u}(p_{\nu_\tau}) \gamma_\nu \left( \frac{1 - \gamma^5}{2} \right) u(p_\tau) \quad (\text{A.6})$$

$$= \frac{1}{2} \text{Tr} \left[ \not{p}_\tau \gamma_\mu \left( \frac{1 - \gamma^5}{2} \right) \not{p}_{\nu_\tau} \gamma_\nu \left( \frac{1 - \gamma^5}{2} \right) \right] \quad (\text{A.7})$$

$$= \frac{1}{2} \text{Tr} \left[ \not{p}_\tau \gamma_\mu \not{p}_{\nu_\tau} \gamma_\nu \left( \frac{1 - \gamma^5}{2} \right) \right] \quad (\text{A.8})$$

$$= (p_\tau^\mu p_{\nu_\tau}^\nu + p_\tau^\nu p_{\nu_\tau}^\mu - g^{\mu\nu} p_{\nu_\tau} \cdot p_\tau - i\epsilon^{\alpha\mu\beta\nu} p_{\tau,\alpha} p_{\nu_\tau,\beta}), \quad (\text{A.9})$$



**Figure 2.** The optical theorem relates the imaginary part of a) to the hadronic  $\tau$  decay b). The blue areas, indicate higher order interactions.

which then have to be dotted into (A.5). We notice that  $\epsilon^{\alpha\mu\beta\nu} p_{\tau,\alpha} p_{\nu,\beta}$  drops out due to symmetry. We will start regarding only the transversal contribution of the Lorentz-decomposed QCD correlator (A.2) and later infer the missing longitudinal one. Thus we obtain

$$\begin{aligned} \frac{1}{2} \sum_{spins} |\mathcal{M}|^2 &= \frac{g^4}{16M_W^2} (p_\tau^\mu p_\nu^\nu + p_\tau^\nu p_\nu^\mu - g^{\mu\nu} p_\nu p_\tau) (q_\mu q_\nu - q^2 g_{\mu\nu}) \Pi^{(T)}(q^2) \\ &= \frac{g^4}{16M_W^2} [2(p_\tau \cdot q)(p_\nu \cdot q) + (p_\nu \cdot p_\tau) q^2] \Pi^{(T)}(q^2). \end{aligned} \quad (\text{A.10})$$

In the center of mass frame we then can find the needed kinematic expressions

$$2p_\tau \cdot p_\nu = 2p_\nu \cdot q = M_\tau^2 - s \quad (\text{A.11})$$

$$2p_\tau \cdot q = M_\tau^2 + s \quad (\text{A.12})$$

and get the full expression of the transversal amplitude

$$\begin{aligned} &\frac{g^4}{16M_W^2} \frac{1}{2} [(M_\tau^2 + s)(M_\tau^2 - s) + (M_\tau^2 - s)s] \Pi^{(T)}(q^2) \\ &= \frac{g^4}{16M_W^4} \left[ \left(1 - \frac{s}{M_\tau^2}\right) \left(1 + \frac{2s}{M_\tau^2}\right) \right] \text{Im} \Pi^{(T)}(q^2). \end{aligned} \quad (\text{A.13})$$

Noting from (A.10) and (A.2) that the longitudinal contribution differs only by a missing

summand  $-q^2 g_{\mu\nu}$  we can infer the longitudinal amplitude

$$\begin{aligned}
& \frac{g^4}{16M_W^4} (p_\tau^\mu p_\nu^\nu + p_\tau^\nu p_\nu^\mu - g^{\mu\nu} p_\nu p_\tau) (q_\mu q_\nu \Pi^{(L)}(q^2)) \\
&= \frac{g^4}{16M_W^4} [2(p_\tau \cdot q)(p_\nu \cdot q) - (p_\nu \cdot p_\tau)s] \Pi^{(L)}(s) \quad . \\
&= \frac{g^4}{16M_W^4} \left[ M_\tau^4 \left( 1 - \frac{s}{M_\tau^2} \right) \right] \Pi^{(L)}(s)
\end{aligned} \tag{A.14}$$

To get to the inclusive hadron decay we have to deal with the integration

$$\int \frac{d^3 p_\nu}{(2\pi)^3} \frac{1}{2E_\nu} . \tag{A.15}$$

Due to the massless  $\tau$ -neutrino and the kinematics we notice that

$$|p_\nu| = \frac{1}{2} \frac{M_\tau^2 - s}{M_\tau} \quad \text{and} \quad E_\nu = p_\nu \tag{A.16}$$

so that the new integrand can be written as:  $dp = -\frac{ds}{2M_\tau}$ . Thus we get a factor of

$$\begin{aligned}
\int \frac{d^3 p_\nu}{(2\pi)^3} \frac{1}{2E_\nu} &= - \int \frac{d\theta d\phi ds}{(2\pi)^3} \frac{1}{2M_\tau} \frac{p_\nu^2}{2p_\nu} \sin^2 \theta \\
&= - \int \frac{d\Omega}{(2\pi)^3} \frac{ds}{4\dot{2}} \left( \frac{M_\tau^2 - s}{M_\tau^2} \right) \quad . \\
&= \int_0^{M_\tau^2} \frac{4\pi M_\tau^2 ds}{(2\pi)^3 8} \left( 1 - \frac{s}{M_\tau^2} \right)
\end{aligned} \tag{A.17}$$

for the integration part of the  $\tau$ -decay rate. Now we just have to combine the longitudinal and transversal contribution with the before evaluated integration factor yielding the final decay rate:

$$\begin{aligned}
\Gamma(\tau \rightarrow \text{hadrons}) &= \frac{1}{2M_\tau} \int_0^{M_\tau^2} \frac{g^4 M_\tau^6}{M_\tau^2 2^8 \pi^2 M_W^4} \left( 1 - \frac{s}{M_\tau^2} \right)^2 \\
&\times \left[ \left( 1 + 2 \frac{s}{M_\tau^2} \right) \text{Im} \Pi^{(T)}(s) + \text{Im} \Pi^{(L)}(s) \right] .
\end{aligned} \tag{A.18}$$

The  $\tau$  decay rate into electron-neutrinos is

$$\Gamma(\tau \rightarrow e \nu_\tau \bar{\nu}_e) = \frac{G_F^2 M_\tau^5}{192 \pi^3} = \frac{g^4 M_\tau^5}{32 \cdot 192 M_W^4 \pi^3} , \tag{A.19}$$

so that the desired ratio is given by

$$R_\tau = 12\pi \int_0^{M_\tau^2} \frac{ds}{M_\tau^2} \left( 1 - \frac{s}{M_\tau^2} \right) \left[ \left( 1 + 2 \frac{s}{M_\tau^2} \right) \text{Im} \Pi^{(T)}(s) + \text{Im} \Pi^{(L)}(s) \right] \tag{A.20}$$

## B Coefficients

Taken from [2]

### B.1 $\beta$ function

$$\begin{aligned}\beta_1 &= \frac{1}{6}(11N_c - 2N_f), \quad \beta_2 = \frac{1}{12}(17N_c^2 - 5N_cN_f - 3C_fN_f), \\ \beta_3 &= \frac{1}{32} \left( \frac{2857}{54}N_c^3 - \frac{1415}{54}N_c^2N_f + \frac{79}{54}N_cN_f^2 - \frac{205}{18}N_cC_fN_f + \frac{11}{9}C_fN_f^2 + C_f^2N_f \right), \quad (\text{B.1}) \\ \beta_4 &= \frac{140599}{2304} + \frac{445}{16}\zeta(3),\end{aligned}$$

with  $C_f = (N_c^2 - 1)/2N_c$ .

### B.2 $\gamma$ -function

$$\begin{aligned}\gamma_1 &= \frac{3}{2}C_f, \quad \gamma_2 = \frac{C_f}{48}(97N_c + 9C_f - 10N_f), \\ \gamma_3 &= \frac{C_f}{32} \left[ \frac{11413}{108}N_c^2 - \frac{129}{4}N_cC_f - \left( \frac{278}{27} + 24\zeta(3) \right) N_cN_f + \frac{129}{2}C_f^2 \right. \\ &\quad \left. - (23 - 24\zeta(3)) C_fN_f - \frac{35}{27}N_f^2 \right], \\ \gamma_4 &= \frac{2977517}{20736} - \frac{9295}{216}\zeta(3) + \frac{135}{8}\zeta(4) - \frac{125}{6}\zeta(5).\end{aligned} \quad (\text{B.2})$$

### B.3 Adler function

$$\begin{aligned}c_{11} &= 1, \quad c_{21} = \frac{365}{24} - 11\zeta_3 - \left( \frac{11}{12} - \frac{2}{3}\zeta(3) \right) N_f = 1.640, \\ c_{31} &= \frac{87029}{288} - \frac{1103}{4}\zeta_3 + \frac{275}{6}\zeta_5 \\ &\quad - \left( \frac{7847}{216} - \frac{262}{9}\zeta_3 + \frac{25}{9}\zeta_5 \right) N_f + \left( \frac{151}{162} - \frac{19}{27}\zeta_3 \right) N_f^2 = 6.371, \\ c_{41} &= \frac{78631453}{20736} - \frac{1704247}{432}\zeta_3 + \frac{4185}{8}\zeta_3^2 + \frac{34165}{96}\zeta_5 - \frac{1995}{16}\zeta_7 = 49.076\end{aligned} \quad (\text{B.3})$$

from [3]

$$\begin{aligned}c_{23} &= 0, \quad c_{22} = \frac{\beta_1 c_{11}}{4}, \\ c_{34} &= 0, \quad c_{33} = \frac{\beta_1^2}{12}c_{11}, \quad c_{32} = -\frac{1}{4}(\beta_2 c_{11} + 2\beta_1 c_{21}), \\ c_{42} &= -\frac{1}{4}(\beta c_{11} + 2\beta_2 c_{21} + 3\beta_1 c_{31}).\end{aligned} \quad (\text{B.4})$$

## C Numerical Analysis

[7] [5]



### **C.1 Newton-Raphson Method**

### **C.2 Newton-Cotes**

### **C.3 Gauss Quadrature**

### **C.4 Discrepancies between Matthias and my Code**

- My spectral moments are slightly lower than Matthias ones (of order  $10^{-2}$ ). Could be caused by the different treatment of weight functions. Mine are integrated with Gaussian Quadratures. Matthias are taken at the center and multiplied with the bin width.

## References

- [1] M. Beneke and M. Jamin.  $\alpha_s$  and the  $\tau$  hadronic width: fixed-order, contour- improved and higher-order perturbation theory. *JHEP*, 09, 044, 2008. URL: [0806.3156](#).
- [2] Diogo Boito. *QCD phenomenology with  $\tau$  and charm decays*. PhD thesis, Universitat Autnoma de Barcelona, September 2011.
- [3] M. Jamin. Qcd and renormalisation group methods. Herbstschule fu r Hochenergiephysik Maria Laach, 2006.
- [4] K.A. u.a. Olive. Review of particle physics. *Chin. Phys. C38 (2014), S. 090001.*, DOI 10.1088/1674-1137/38/9/090001, 2014. URL: <http://dx.doi.org/10.1088/1674-1137/38/9/090001>.
- [5] William H. Press, Saul A. Teukolsky William, T. Vetterling, and Brian P. Flannery. *Numerical Recipes*. Cambridge, 2007.
- [6] Felix Schwab. Strange quark mass determination from sum rules for hadronic  $\tau$ -decays. Master’s thesis, University of Heidelberg, 2002.
- [7] Raymond P. Canale Steven C. Chapra. *Numerical Methods for Engineers*. McGraw Hill, 2010.
- [8] Y. Tsai. Decay correlations of heavy leptons in  $e^+e^- \rightarrow l^+l^-$ . *Phys. Rev.*, D4, 1971.