

Have coaches changed how they select which players to give more minutes to?

Derek Corcoran^{*,1}, Nicholas M. Watanabe¹

^a *Department, Street, City, State, Zip*

^b *Department, Street, City, State, Zip*

Abstract

Since the NBA adopted the three point line in 1979 the league has had several key rule changes (eg hand checking rules, allowing zone defense) that have altered the value of different skill-sets players may have. With the creation of the three point shot, a shot made from behind the arc was worth more, which made long distance shooting more valuable. We explore which stats better explain the minutes per game played in every season. Mostly we see that assist per 100 possessions and points per 100 possessions are the two stats have the most explanation for minutes per game played. We notice that since 2009, rebounding per 100 possessions stopped being an important variable selected by coaches

Text based on elsarticle sample manuscript, see <http://www.elsevier.com/author-schemas/latex-instructions#elsarticle>

Methods

The basketball reference player season finder was used to extract the per 100 team possessions stats, single season, during the three point era (since season 1979-80), during the regular season (“Player Season Finder Basketball-Reference.com” 2017), and that information was coupled with the minutes per game played by each player, again extracted from the basketball reference player season finder but now in the per game stats. We divided that data-set into 37 data-sets, one for each season from the 1979-80 season to the 2015-16 season.

Queries

- For single seasons, played in the NBA/BAA, in the regular season, from 1979-80 to 2016-17, played G or G-F, qualified for Minutes Per Game Leaderboard, sorted by descending Win Shares
- For single seasons, played in the NBA/BAA, in the regular season, from 1979-80 to 2016-17, played F or F-G or F-C, qualified for Minutes Per Game Leaderboard, sorted by descending Win Shares
- For single seasons, played in the NBA/BAA, in the regular season, from 1979-80 to 2016-17, played C or C-F, qualified for Minutes Per Game Leaderboard, sorted by descending Win Shares

For each season, the per 100 team possession stats was used to fit a global glm model (Nelder and Baker 1972) that explains the minutes per game of each player based on the following variables: Two point shot attempts per 100 possessions, two point shot percentage, three point shots attempts per 100 possessions, three point shot percentage, free throw attempts per 100 possessions, free throw percentage, total rebounds per 100 possessions, assists per 100 possessions, steals per 100 possessions, blocks per 100 possessions, turnovers per 100 possessions, points per 100 possessions and effective field goal percentage.

In order to be able to compare the strength of relationship of every variable on the same scale, all of them were scaled and centered (Bro and Smilde 2003) using the caret package (Kuhn and Johnson 2013).

*Corresponding Author

Email addresses: derek.corcoran.barrios@gmail.com (Derek Corcoran), nmwatana@olemiss.edu (Nicholas M. Watanabe)

For each season, we tested variables for collinearity. Then we fitted every possible first order model not allowing models to coexist if they had a Pearson correlation coefficient equal or higher than 0.7 (Dormann et al. 2013). Then the models were ranked based on Akaike's Information Criteria for small sample sizes (AICc) (Cavanaugh 1997) using the MuMin Package (Bartoń 2013; Burnham and Anderson 2002). We didn't use model averaging since even though collinear variables were prohibited to coexist in the same model, these might coexist in the average model (Cade 2015), thus we selected the best possible model for each season selecting by AICc (Burnham and Anderson 2002). All of the analyses using R statistical Software (Team 2016),

Results

Centers

As we can see in figure 1, the Season is the variable that appears in most seasons being selected in 38 of 37 seasons, followed by TwoPA and AST being selected in 19 and 12 seasons respectively

Forwards

As we can see in figure 1, the Season is the variable that appears in most seasons being selected in 38 of 37 seasons, followed by STL and TwoPA being selected in 28 and 22 seasons respectively

Guards

As we can see in figure 1, the TRB is the variable that appears in most seasons being selected in 37 of 37 seasons, followed by AST and FTA being selected in 34 and 33 seasons respectively

Functionality. The Elsevier article class is based on the standard article class and supports almost all of the functionality of that class. In addition, it features commands and options to format the

- document style
- baselineskip
- front matter
- keywords and MSC codes
- theorems, definitions and proofs
- labels of enumerations
- citation style and labeling.

Front matter

The author names and affiliations could be formatted in two ways:

- (1) Group the authors per affiliation.
- (2) Use footnotes to indicate the affiliations.

See the front matter of this document for examples. You are recommended to conform your choice to the journal you are submitting to.

Bibliography styles

There are various bibliography styles available. You can select the style of your choice in the preamble of this document. These styles are Elsevier styles based on standard styles like Harvard and Vancouver. Please use BibTeX to generate your bibliography and include DOIs whenever available.

Here are two sample references: Bartoń (2013; Cade 2015).

References

- Bartoń, Kamil. 2013. “MuMIn: Multi-Model Inference. R Package Version 1.9. 13.” *The Comprehensive R Archive Network (CRAN)*, Vienna, Austria.
- Bro, Rasmus, and Age K Smilde. 2003. “Centering and Scaling in Component Analysis.” *Journal of Chemometrics* 17 (1). Wiley Online Library: 16–33.
- Burnham, KP, and DR Anderson. 2002. “Information and Likelihood Theory: A Basis for Model Selection and Inference.” *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach 2*. Springer-Verlag New York: 49–97.
- Cade, Brian S. 2015. “Model Averaging and Muddled Multimodel Inferences.” *Ecology* 96 (9). Wiley Online Library: 2370–82.
- Cavanaugh, Joseph E. 1997. “Unifying the Derivations for the Akaike and Corrected Akaike Information Criteria.” *Statistics & Probability Letters* 33 (2). Elsevier: 201–8.
- Dormann, Carsten F, Jane Elith, Sven Bacher, Carsten Buchmann, Gudrun Carl, Gabriel Carré, Jaime R García Márquez, et al. 2013. “Collinearity: A Review of Methods to Deal with It and a Simulation Study Evaluating Their Performance.” *Ecography* 36 (1). Wiley Online Library: 27–46.
- Kuhn, Max, and Kjell Johnson. 2013. *Applied Predictive Modeling*. Vol. 26. Springer.
- Nelder, John A, and R Jacob Baker. 1972. “Generalized Linear Models.” *Encyclopedia of Statistical Sciences*. Wiley Online Library.
- “Player Season Finder Basketball-Reference.com.” 2017. Accessed February 27. http://www.basketball-reference.com/play-index/psl_finder.cgi.
- Team, R Core. 2016. “R: A Language and Environment for Statistical Computing. R Development Core Team, Vienna.”

List of Figures

1	Strength of relationship by season for Centers	5
2	Strength of relationship by season for Forwards	6
3	Strength of relationship by season for Guards	7




Figure 1: Strength of relationship by season for Centers




Figure 2: Strength of relationship by season for Forwards




Figure 3: Strength of relationship by season for Guards