The transverse rigidity of the particle arrangement: hypotheses and tests

D. Teaney[1]

[1]*Department of Physics and Astronomy, Stony Brook University, Stony Brook, New York 11794, USA*

(Dated: November 9, 2026)

Abstract

The equilibrium runs (`campaigns.tex`) show transverse sound: the shear mode oscillates with $c_t = 0.022$ instead of diffusing. This note separates what is measured from what is conjectured, states the conjectures with the tests that can falsify them (all in the campaign `rigidity`, now running), and lists ways to remove the rigidity, to be discussed before any is implemented.

I. WHAT IS MEASURED

All at $h/a = 2$, Wendland C2 kernel, $\delta v = 0.043$ ($s_a = 541$), $\eta/s = 0.2$ ($\nu = \eta/w = 2.2 \times 10^{-3}$).

1. $C_T(k,\tau)$ is fitted, within its errors, by the velocity correlator of a damped oscillator,

$$C_T(k,\tau) = A\, e^{-\gamma\tau}\Big[\cos\omega\tau - \frac{\gamma}{\omega}\sin\omega\tau\Big], \qquad \omega^2 + \gamma^2 = c_t^2 k^2, \tag{1}$$

   with $c_t = 0.022$ for $kh = 0.13$–0.74, in boxes 1.2 and 2.4, from two starts and two versions of the code. The fit recovers c_t and γ from synthetic data with known values, and returns $c_t = 0$ for a synthetic fluid.

2. $2\gamma/\nu k^2 = 1.1$–1.2 at $kh \geq 0.5$, rising to ≈ 2 at $kh = 0.13$.

3. The local disorder (rms change of nearest-neighbour distances) saturates at $0.3a$: the particles do not exchange neighbours.

4. Ideal SPH alone (no viscous step), box 0.6: from the lattice with thermal momenta the kinetic energy grows from 0.9 to 6.8 times its equilibrium value by $t = 20$, still rising, while H is conserved to 10^{-7}.

5. Ideal SPH alone from random positions (smoothed entropy made uniform): the kinetic energy reaches 420 times its equilibrium value by $t = 5$. The full scheme from random positions heats by 64% in H by $t = 10$.

II. HYPOTHESIS A: A BROKEN RELABELING SYMMETRY GIVES A SHEAR MODULUS

a. The continuum. The energy of a perfect fluid depends on the density and entropy fields only. A volume-preserving relabeling of the fluid elements leaves the fields, and so the energy, unchanged: a transverse displacement $\boldsymbol{u}$ with $\nabla \cdot \boldsymbol{u} = 0$ costs no energy, and there is no shear modulus.

b. SPH. The smoothed entropy at particle a is $S_a = \sum_b s_b W(\boldsymbol{x}_a - \boldsymbol{x}_b)$. On a lattice with equal $s_b = sV$,

$$S_a = s \sum_{\boldsymbol{G}} \tilde{W}(\boldsymbol{G})\, e^{i\boldsymbol{G}\cdot\boldsymbol{x}_a}, \tag{2}$$

a sum over the reciprocal lattice. The term $\boldsymbol{G} = 0$ is the exact density; the others, set by $\tilde{W}$ at $|\boldsymbol{G}| \geq b = 4\pi/(\sqrt{3}a)$, are the lattice error. A shear at fixed area changes the $|\boldsymbol{G}|$, hence the sum, at second order in the strain ϵ: the energy rises by $\frac{1}{2}G\epsilon^2$ per volume with, by this estimate,

$$\frac{G}{p} \sim C\,(bh)^2\,\left|\tilde{W}''(bh)\right| \sim C\,m(m+1)\,\left|\tilde{W}(bh)\right|, \tag{3}$$

where $\tilde{W} \propto (bh)^{-m}$ asymptotically and C is a number of order the count of shortest $\boldsymbol{G}$ (six). This is an order-of-magnitude argument, not a calculation; in particular the forces in SPH depend on ∇W as well, and the stochastic step may add to or remove from G.

c. Predictions. $\tilde{W}$ at the first reciprocal vector:

kernel	$\langle r^2\rangle/h^2$	$\tilde{W}(bh)$, $h/a = 2$	2.5	3
Wendland C2	0.56	1.2×10^{-4}	3.7×10^{-5}	1.5×10^{-5}
Wendland C4	0.42	3.5×10^{-5}	7.6×10^{-6}	2.2×10^{-6}
Wendland C6	0.34	1.6×10^{-5}	2.3×10^{-6}	5.0×10^{-7}

If $G \propto \tilde{W}(bh)$, then $c_t = (G/w)^{1/2} \propto \tilde{W}(bh)^{1/2}$, and from $c_t = 0.022$ for C2 at $h/a = 2$:

run	c_t if $G \propto \tilde{W}$	c_t if $G \propto \tilde{W}^2$
C2, $h/a = 2.5$	0.012	0.0068
C2, $h/a = 3$	0.0078	0.0028
C4, $h/a = 2$	0.012	0.0064
C4, $h/a = 2.5$	0.0055	0.0014
C6, $h/a = 2$	0.0081	0.0030

d. Falsified if: c_t does not fall with h/a and with the kernel order; or C6 at $h/a = 2$ and C2 at $h/a = 3$ (equal $\tilde{W}(bh)$, different h/a) give different c_t; or the rigidity is absent when the positions are frozen *and* present only with the stochastic step (then it is not a property of H).

e. Second test of the same quantity. The runs record the displacements $\boldsymbol{u}_a$ from the initial sites. For an elastic medium $\langle|\dot{u}_T|^2\rangle = c_t^2 k^2 \langle|u_T|^2\rangle$, and $\boldsymbol{M}_{\boldsymbol{k}} \simeq wV\dot{\boldsymbol{u}}_{\boldsymbol{k}}$, so the statics alone give

$$c_t^2 = \frac{\langle|M_T(\boldsymbol{k})|^2\rangle}{(wV)^2 k^2 \langle|u_T(\boldsymbol{k})|^2\rangle}. \tag{4}$$

In a fluid $\langle|u_T|^2\rangle$ grows linearly in time without bound. Agreement with the fit of eq. (1) is a check that the oscillation is elastic.

III. HYPOTHESIS B: THE ARRANGEMENT CARRIES ENERGY

Facts 4 and 5 show that at fixed s_a the SPH energy depends on the arrangement, by a lot for a random one ($\sim 400T$ per particle released) and by a little for the lattice ($\sim 6T$ per particle by $t = 20$, still growing). The conjecture is that at temperature T the equilibrium of the positions lies near a minimum of this energy, so the particles form a solid whatever the start, and a random arrangement is not an equilibrium state.

This is *not* established. Fact 4 says that the hexagonal lattice is not itself the minimum (ideal SPH moves away from it, releasing energy), so the relevant minimum is some nearby disordered arrangement, and why the full scheme stays at a disorder of $0.3a$ is not explained. The ideal-SPH runs at $h/a = 2$, 2.5, 3 and for C4, C6 (to $t = 50$) show whether the kinetic energy saturates, and how the released energy scales with $\tilde{W}(bh)$. If it scales like G, A and B are the same effect.

IV. WAYS OUT, IF A HOLDS

a. Smaller lattice error. A fluid needs $c_t k \ll \nu k^2/2$ at the smallest k of interest,

$$c_t \ll \frac{\nu k_{\min}}{2} = \frac{\pi \nu}{L}, \tag{5}$$

about 3×10^{-3} for $L = 2.4$: a factor 7 in c_t, 50 in G. Whether a kernel and h/a reach this at acceptable cost is what the runs measure.

b. Noise in $d\boldsymbol{x}/dt$. Particles that exchange neighbours relax shear stress in a Maxwell time $\tau_M \sim a^2/D_x$ (diffusion constant D_x of the positions relative to the fluid), turning the solid into a fluid with an added viscosity $G\tau_M$. This is small compared with η when

$$D_x \gg \frac{c_t^2 a^2}{\nu} \approx 0.06\,\nu. \tag{6}$$

Three versions, with their expected outcomes:

1. *Displacements alone* (s_a, $\boldsymbol{m}_a$ fixed): each move raises the arrangement energy (Hypothesis B), which ends up as heat; expected to heat strongly.

2. *Displacements with a Metropolis test on e^S*: samples the true equilibrium of the positions; if B holds that is the solid, and G is unchanged.

3. *Relabeling*: displace a particle and refit s_a, $\boldsymbol{m}_a$, e_a so that the smoothed fields are unchanged, as molecular self-diffusion leaves the hydrodynamic fields unchanged. The energy change is then only the error of the refit, which a Metropolis test can absorb. This restores the relabeling symmetry to the accuracy of the refit. It needs a local linear solve per move that conserves E and $\boldsymbol{M}$, and a check that it does not spoil the detailed balance of the stochastic step.

A quick test of each at $L = 0.6$ (heating and c_t against D_x) would settle 1 and 2.

c. Not proposed. Lowering s_a to melt the arrangement: the scheme blows up at $\delta v = 0.14$ (`campaigns.tex`).