

Almost everything you need to know about PLS

Part 1: Background, Theory, and Examples

Jenny Rieck & Derek Beaton

October 24, 2017

The BIG outline

- Part 1: Background & Examples

The BIG outline

- Part 1: Background & Examples
 - RIGHT NOW

The BIG outline

- Part 1: Background & Examples
 - RIGHT NOW
 - Introduce everything we need

The BIG outline

- Part 1: Background & Examples
 - RIGHT NOW
 - Introduce everything we need
- Part 2: PLS in Matlab & R

The BIG outline

- Part 1: Background & Examples
 - RIGHT NOW
 - Introduce everything we need
- Part 2: PLS in Matlab & R
 - Tuesday November 21, 10:00-12:00 Worstman Hall

The BIG outline

- Part 1: Background & Examples
 - RIGHT NOW
 - Introduce everything we need
- Part 2: PLS in Matlab & R
 - Tuesday November 21, 10:00-12:00 Worstman Hall
 - Put knowledge into practice

Part 1 outline

- Theory & Background

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)
- Short break

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)
- Short break
- Examples (with ADNI)

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
 - Standard PLS (& inference)

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
 - Standard PLS (& inference)
 - Discriminant PLS

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
 - Standard PLS (& inference)
 - Discriminant PLS
 - “Seed” PLS

Part 1 outline

- Theory & Background
 - Principal component analysis (PCA)
 - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
 - Standard PLS (& inference)
 - Discriminant PLS
 - “Seed” PLS
- And beyond!

Principal Component Analysis

Background

Where did PCA come from?

- Modern form

Where did PCA come from?

- Modern form
 - Hotelling (1933)

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)
 - Galton (1859)

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)
 - Galton (1859)
 - Pearson (1901)

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)
 - Galton (1859)
 - Pearson (1901)
- Modern overviews

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)
 - Galton (1859)
 - Pearson (1901)
- Modern overviews
 - S. Wold et al., (1987)

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)
 - Galton (1859)
 - Pearson (1901)
- Modern overviews
 - S. Wold et al., (1987)
 - Jolliffe (2002)

Where did PCA come from?

- Modern form
 - Hotelling (1933)
- Traces back to
 - Cauchy (1829)
 - Galton (1859)
 - Pearson (1901)
- Modern overviews
 - S. Wold et al., (1987)
 - Jolliffe (2002)
 - Abdi & Williams (2010)

What is PCA?

- Visualize high dimensional data

What is PCA?

- Visualize high dimensional data
- Orthogonal transformation

What is PCA?

- Visualize high dimensional data
- Orthogonal transformation
- Dimensionality reduction

What is PCA?

- Find “components”

What is PCA?

- Find “components”
 - Components are new variables that are combinations of old variables

What is PCA?

- Find “components”
 - Components are new variables that are combinations of old variables
- Components explain maximum possible variance

What is PCA?

- Find “components”
 - Components are new variables that are combinations of old variables
- Components explain maximum possible variance
 - Conditional to orthogonality

Visual example






Figure 1: The kind of data we usually expect for PCA


Visual example




Visual example




Visual example




Visual example




Visual example




Visual example




Visual example



Visual example



Visual example



What is PCA?

- The basis of modern techniques

What is PCA?

- The basis of modern techniques
 - Factor analyses

What is PCA?

- The basis of modern techniques
 - Factor analyses
 - Independent components analysis

What is PCA?

- The basis of modern techniques
 - Factor analyses
 - Independent components analysis
 - Partial least squares

What is PCA?

- The basis of modern techniques
 - Factor analyses
 - Independent components analysis
 - Partial least squares
 - Discriminant analyses

What is PCA?

- The basis of modern techniques
 - Factor analyses
 - Independent components analysis
 - Partial least squares
 - Discriminant analyses
 - Multi-table (e.g., MFA, GCCA)

What is PCA?

- A special case of the singular value decomposition (SVD)

What is PCA?

- A special case of the singular value decomposition (SVD)
- Which means (almost) everything else is, too

Formalization

Singular value decomposition

The SVD is one of the most ubiquitous and important tools

SVD

- We'll go into *enough* formalization

SVD

- We'll go into *enough* formalization
- If you want more:

SVD

- We'll go into *enough* formalization
- If you want more:
 - S. Wold et al., (1987)

SVD

- We'll go into *enough* formalization
- If you want more:
 - S. Wold et al., (1987)
 - Jolliffe (2002)

SVD

- We'll go into *enough* formalization
- If you want more:
 - S. Wold et al., (1987)
 - Jolliffe (2002)
 - Abdi & Williams (2010)

SVD

- We'll go into *enough* formalization
- If you want more:
 - S. Wold et al., (1987)
 - Jolliffe (2002)
 - Abdi & Williams (2010)
 - And many others...

SVD



Figure 2: The shape of the data

SVD




Figure 3: SVD breaks down the data

SVD

Notation

SVD

Notation

- x - a scalar

SVD

Notation

- x - a scalar
- \mathbf{a} - a vector

SVD

Notation

- x - a scalar
- \mathbf{a} - a vector
- \mathbf{A} - a matrix

SVD

Notation

- x - a scalar
- \mathbf{a} - a vector
- \mathbf{A} - a matrix
- \mathbf{A}^T - transpose

SVD

Notation

- x - a scalar
- \mathbf{a} - a vector
- \mathbf{A} - a matrix
- \mathbf{A}^T - transpose
- \mathbf{AB} - multiplication

SVD

- Think back to PCA

SVD

- Think back to PCA
- We want to find *the* principal component

SVD

- Think back to PCA
- We want to find *the* principal component
 - a.k.a. maximum source of variance

SVD

Given a matrix \mathbf{X} we generally assume that

SVD

Given a matrix \mathbf{X} we generally assume that

- column-wise centered

SVD

Given a matrix \mathbf{X} we generally assume that

- column-wise centered
- column-wise scaled (e.g., z-scores or sums of squares = 1)

SVD

Given \mathbf{X} of size $I \times J$

SVD

Given \mathbf{X} of size $I \times J$

We want to find vectors

SVD

Given \mathbf{X} of size $I \times J$

We want to find vectors

- \mathbf{u} of size $I \times 1$
- \mathbf{v} of size $J \times 1$

SVD

Given \mathbf{X} of size $I \times J$

We want to find vectors

- \mathbf{u} of size $I \times 1$
- \mathbf{v} of size $J \times 1$

such that

$$\operatorname{argmax}_{\mathbf{u}, \mathbf{v}} \delta = \mathbf{u}^T \mathbf{X} \mathbf{v} \text{ conditional to } \mathbf{u}^T \mathbf{u} = 1 = \mathbf{v}^T \mathbf{v}$$

SVD

Which gives us the following equivalencies:

- $\mathbf{X}\mathbf{v} = \mathbf{u}\delta$
- $\mathbf{X}^T \mathbf{u} = \mathbf{v}\delta$

SVD

Which gives us the following equivalencies:

- $\mathbf{X}\mathbf{v} = \mathbf{u}\delta$
- $\mathbf{X}^T\mathbf{u} = \mathbf{v}\delta$

where

- $\mathbf{X}_1 = \delta\mathbf{u}\mathbf{v}^T$

SVD

Which gives us the following equivalencies:

- $\mathbf{X}\mathbf{v} = \mathbf{u}\delta$
- $\mathbf{X}^T \mathbf{u} = \mathbf{v}\delta$

where

- $\mathbf{X}_1 = \delta \mathbf{u} \mathbf{v}^T$
 - \mathbf{X}_1 is \mathbf{X} as represented by *source of maximum variance*

SVD

Given $\mathbf{X}_1 = \delta \mathbf{p} \mathbf{q}^T$ we can find the other components

SVD

Given $\mathbf{X}_1 = \delta \mathbf{p} \mathbf{q}^T$ we can find the other components

- $\mathbf{X}_{2:L} = \mathbf{X} - \mathbf{X}_1$
- $\mathbf{X}_{2:L}$ is orthogonal to \mathbf{X}_1

SVD

Given $\mathbf{X}_1 = \delta \mathbf{p} \mathbf{q}^T$ we can find the other components

- $\mathbf{X}_{2:L} = \mathbf{X} - \mathbf{X}_1$
- $\mathbf{X}_{2:L}$ is orthogonal to \mathbf{X}_1

$$\operatorname{argmax}_{\mathbf{p}, \mathbf{q}} \delta = \mathbf{p}^T \mathbf{X}_{2:L} \mathbf{q} \text{ conditional to } \mathbf{p}^T \mathbf{p} = 1 = \mathbf{q}^T \mathbf{q}$$

SVD

The SVD of \mathbf{X} of size $I \times J$:

SVD

The SVD of \mathbf{X} of size $I \times J$:

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \quad (1)$$

SVD

The SVD of \mathbf{X} of size $I \times J$:

$$\mathbf{X} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^T \quad (1)$$

\mathbf{U} and \mathbf{V} are orthonormal such that

SVD

The SVD of \mathbf{X} of size $I \times J$:

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \quad (1)$$

\mathbf{U} and \mathbf{V} are orthonormal such that

$$\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V} \quad (2)$$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- where rank is L (i.e., number of extractable components)

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- where rank is L (i.e., number of extractable components)
- \mathbf{U} is $I \times L$ (left singular vectors; rows of \mathbf{X})

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- where rank is L (i.e., number of extractable components)
- \mathbf{U} is $I \times L$ (left singular vectors; rows of \mathbf{X})
- \mathbf{V} is $J \times L$ (right singular vectors; columns of \mathbf{X})

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{\Delta}$ is $L \times L$ diagonal matrix

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{\Delta}$ is $L \times L$ diagonal matrix
- $\text{diag}\{\mathbf{\Delta}\} = \boldsymbol{\delta}$ are singular values (decreasing)

SVD

$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T$ such that $\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$

- $\mathbf{\Delta}$ is $L \times L$ diagonal matrix
- $\text{diag}\{\mathbf{\Delta}\} = \boldsymbol{\delta}$ are singular values (decreasing)
- $\boldsymbol{\lambda} = \boldsymbol{\delta}^2$ are the eigenvalues (variance)

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

SVD

$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T$ such that $\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$

- $\mathbf{x}_1 = \mathbf{U}_1\mathbf{\Delta}_1\mathbf{V}_1^T$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{X}_1 = \mathbf{U}_1\mathbf{\Delta}_1\mathbf{V}_1^T$
- $\mathbf{X}_{2:L} = \mathbf{U}_{2:L}\mathbf{\Delta}_{2:L}\mathbf{V}_{2:L}^T$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta}$ (row component scores)

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta}$ (row component scores)
- $\mathbf{F}_J = \mathbf{V}\mathbf{\Delta}$ (column component scores)

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

SVD

$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T$ such that $\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta} = \mathbf{X}\mathbf{V}$

SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta} = \mathbf{X}\mathbf{V}$
- $\mathbf{F}_J = \mathbf{V}\mathbf{\Delta} = \mathbf{X}^T\mathbf{U}$

SVD

- Phew.

SVD

- Phew.
- Enough nerd stuff

SVD

- Phew.
- Enough nerd stuff
- Let's get back to PCA

Toy example

- In R with ExPosition packages

- In R with ExPosition packages
 - Sneak preview of newest version

- In R with ExPosition packages
 - Sneak preview of newest version
- <https://github.com/derekbeaton/ExPosition-Family/>

- In R with ExPosition packages
 - Sneak preview of newest version
- <https://github.com/derekbeaton/ExPosition-Family/>
- <https://cran.r-project.org/web/packages/ExPosition/index.html>

PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)

PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)

PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)

PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)
 - We'll use this one for PCA

PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)
 - We’ll use this one for PCA
 - We’ll use both for PLS

PCA Subjective Measures

	fruity	floral	vegetal	spicy
Chili_red_merlot	6	2	1	4
Chili_red_cabernet	5	3	2	3
Chili_red_shiraz	7	1	2	6
Canada_red_pinot	4	2	3	1
Canada_white_chardonnay	4	3	2	1
Canada_white_sauvignon	8	4	3	2
USA_rose_cabernet	8	3	3	3
USA_rose_pinot	6	1	1	2
USA_rose_syrah	9	3	2	5

PCA - Wine

```
pca.res <- epPCA(  
  wine.subjective,  
  scale = T,  
  center = T,  
  graphs = F)
```

PCA - Wine

- PCA maximizes variance

PCA - Wine

- PCA maximizes variance
- So let's visualize the variance per component

PCA - Wine

- PCA maximizes variance
- So let's visualize the variance per component
 - Use the eigenvalues

PCA - Wine

- PCA maximizes variance
- So let's visualize the variance per component
 - Use the eigenvalues
 - "Scree plot"

PCA - Wine

- Many ways to present the results

PCA - Wine

- Many ways to present the results
- We prioritize visualization

PCA - Wine

- Many ways to present the results
- We prioritize visualization
 - Use numbers and tables when needed

PCA - Wine

- Many ways to present the results
- We prioritize visualization
 - Use numbers and tables when needed
- PCA is half


PCA - Wine

- Many ways to present the results
- We prioritize visualization
 - Use numbers and tables when needed
- PCA is half
 - Stats


PCA - Wine

- Many ways to present the results
- We prioritize visualization
 - Use numbers and tables when needed
- PCA is half
 - Stats
 - Art


PCA - Wine




PCA - Wine



PCA - Wine



PCA - Wine



PCA - Wine




Figure 4: Variables & Observations

PCA

- If you know PCA you know about 90% of the multivariate stats in use

Partial least squares

Background

PLS means a lot of things

- Projection onto latent structures

PLS means a lot of things

- Projection onto latent structures
 - Probably the most accurate name

PLS means a lot of things

- Projection onto latent structures
 - Probably the most accurate name
 - But also probably too broad a definition

PLS means a lot of things

Partial least squares sounds like ordinary least squares

- When we have two matrices: \mathbf{X} and \mathbf{Y}

PLS means a lot of things

Partial least squares sounds like ordinary least squares

- When we have two matrices: \mathbf{X} and \mathbf{Y}
- OLS: $(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$

PLS means a lot of things

Partial least squares sounds like ordinary least squares

- When we have two matrices: \mathbf{X} and \mathbf{Y}
- OLS: $(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$
- PLS: $\mathbf{X}^T \mathbf{Y}$

PLS means a lot of things

- Partial least squares path modelling (PLS-PM)

PLS means a lot of things

- Partial least squares path modelling (PLS-PM)
- Partial least squares regression (PLSR)

PLS means a lot of things

- Partial least squares path modelling (PLS-PM)
- Partial least squares regression (PLSR)
- Partial least squares correlation (PLSC)

PLS means a lot of things

- Partial least squares path modelling (PLS-PM)
- Partial least squares regression (PLSR)
- Partial least squares correlation (PLSC)
 - This is the one we'll talk about today

PLSC has many...

Names

- Inter-battery (factor) analysis (Tucker, 1958)

PLSC has many...

Names

- Inter-battery (factor) analysis (Tucker, 1958)
- Covariance between two fields (Bretherton, Smith, & Wallace, 1992)

PLSC has many...

Names

- Inter-battery (factor) analysis (Tucker, 1958)
- Covariance between two fields (Bretherton, Smith, & Wallace, 1992)
- PLS-SVD (Tenenhaus, 2005)

PLSC has many...

Names

- Inter-battery (factor) analysis (Tucker, 1958)
- Covariance between two fields (Bretherton, Smith, & Wallace, 1992)
- PLS-SVD (Tenenhaus, 2005)
- co-inertia analysis (Dray, 2014)

PLSC has many...

Friends

- Reduced Rank Regression

PLSC has many...

Friends

- Reduced Rank Regression
- Canonical Correlation Analysis

PLSC has many...

Friends

- Reduced Rank Regression
- Canonical Correlation Analysis
- (Fisher's) Linear Discriminant Analysis

PLSC has many...

Friends

- Reduced Rank Regression
- Canonical Correlation Analysis
- (Fisher's) Linear Discriminant Analysis
- PLS-correspondence analysis

PLSC

History

- McIntosh, Bookstein, Haxby, & Grady (1996)

PLSC

History

- McIntosh, Bookstein, Haxby, & Grady (1996)
- Bookstein (1992)

PLSC

History

- McIntosh, Bookstein, Haxby, & Grady (1996)
- Bookstein (1992)
- Tucker (1958)

PLSC

Modern overviews

- McIntosh & Lobaugh (2004)

PLSC

Modern overviews

- McIntosh & Lobaugh (2004)
- Krishnan et al., (2011)

PLSC

It's effectively just PCA applied to the cross product of two matrices measured on the same observations:

PLSC

It's effectively just PCA applied to the cross product of two matrices measured on the same observations:

- \mathbf{X} which is $I \times J$

PLSC

It's effectively just PCA applied to the cross product of two matrices measured on the same observations:

- **X** which is $I \times J$
- **Y** which is $I \times K$

Formalization

PLSC via the SVD

Given \mathbf{X} of size $I \times J$ and \mathbf{Y} of size $I \times K$

PLSC via the SVD

Given \mathbf{X} of size $I \times J$ and \mathbf{Y} of size $I \times K$

We want to find vectors

PLSC via the SVD

Given \mathbf{X} of size $I \times J$ and \mathbf{Y} of size $I \times K$

We want to find vectors

- \mathbf{u} of size $J \times 1$
- \mathbf{v} of size $K \times 1$

To define latent variables

PLSC via the SVD

Given \mathbf{X} of size $I \times J$ and \mathbf{Y} of size $I \times K$

We want to find vectors

- \mathbf{u} of size $J \times 1$
- \mathbf{v} of size $K \times 1$

To define latent variables

- $\mathbf{l}_X = \mathbf{X}\mathbf{u}$ of size $I \times 1$
- $\mathbf{l}_Y = \mathbf{Y}\mathbf{v}$ of size $I \times 1$

PLSC via the SVD

Given \mathbf{X} of size $I \times J$ and \mathbf{Y} of size $I \times K$

We want to find vectors

- \mathbf{u} of size $J \times 1$
- \mathbf{v} of size $K \times 1$

To define latent variables

- $\mathbf{l}_X = \mathbf{X}\mathbf{u}$ of size $I \times 1$
- $\mathbf{l}_Y = \mathbf{Y}\mathbf{v}$ of size $I \times 1$

such that

$$\operatorname{argmax}_{\mathbf{u}, \mathbf{v}} \delta = \mathbf{u}^T \mathbf{X}^T \mathbf{Y} \mathbf{v} \text{ conditional to } \mathbf{u}^T \mathbf{u} = 1 = \mathbf{v}^T \mathbf{v}$$

PLSC via the SVD

Compute the relationship between \mathbf{X} and \mathbf{Y}

$$\mathbf{R} = \mathbf{X}^T \mathbf{Y} \quad (3)$$

PLSC via the SVD

Compute the relationship between \mathbf{X} and \mathbf{Y}

$$\mathbf{R} = \mathbf{X}^T \mathbf{Y} \quad (3)$$

Compute the SVD of \mathbf{R}

$$\mathbf{R} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^T \quad (4)$$

PLSC via the SVD

Compute the relationship between \mathbf{X} and \mathbf{Y}

$$\mathbf{R} = \mathbf{X}^T \mathbf{Y} \quad (3)$$

Compute the SVD of \mathbf{R}

$$\mathbf{R} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^T \quad (4)$$

Compute the latent variables

$$\mathbf{L}_\mathbf{X} = \mathbf{X} \mathbf{U} \text{ and } \mathbf{L}_\mathbf{Y} = \mathbf{Y} \mathbf{V} \quad (5)$$

PLSC via the SVD

Almost everything is the same:

- Δ are singular values

PLSC via the SVD

Almost everything is the same:

- Δ are singular values
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})

PLSC via the SVD

Almost everything is the same:

- Δ are singular values
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})
- $\mathbf{F}_K = \mathbf{V}\Delta$ (component scores for variables of \mathbf{Y})

PLSC via the SVD

Almost everything is the same:

- Δ are singular values
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})
- $\mathbf{F}_K = \mathbf{V}\Delta$ (component scores for variables of \mathbf{Y})
- So our nomenclature will align with PCA

PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$ express the individuals w.r.t. \mathbf{X}

PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$ express the individuals w.r.t. \mathbf{X}
- $\mathbf{L}_Y = \mathbf{Y}\mathbf{V}$ express the individuals w.r.t. \mathbf{Y}

PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$ express the individuals w.r.t. \mathbf{X}
- $\mathbf{L}_Y = \mathbf{Y}\mathbf{V}$ express the individuals w.r.t. \mathbf{Y}
- Not in PCA

PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$ express the individuals w.r.t. \mathbf{X}
- $\mathbf{L}_Y = \mathbf{Y}\mathbf{V}$ express the individuals w.r.t. \mathbf{Y}
- Not in PCA
 - We'll call these "latent variable scores"

PLSC via the SVD

Maximizes the latent variables

PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \mathbf{\Delta} \quad (6)$$

PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \mathbf{\Delta} \quad (6)$$

When expanded

$$(\mathbf{XU})^T (\mathbf{YV}) = \mathbf{\Delta}$$

PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{R} \mathbf{V} = \Delta$$

PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{R} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{U} \Delta \mathbf{V}^T \mathbf{V} = \Delta$$

PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{XU})^T (\mathbf{YV}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{R} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{U} \Delta \mathbf{V}^T \mathbf{V} = \Delta$$

because

$$\mathbf{U}^T \mathbf{U} = \mathbf{I} = \mathbf{V}^T \mathbf{V}$$

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values
- \mathbf{U} akin to loadings (for variables of \mathbf{X})

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values
- \mathbf{U} akin to loadings (for variables of \mathbf{X})
- \mathbf{V} akin to loadings (for variables of \mathbf{Y})

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values
- \mathbf{U} akin to loadings (for variables of \mathbf{X})
- \mathbf{V} akin to loadings (for variables of \mathbf{Y})
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values
- \mathbf{U} akin to loadings (for variables of \mathbf{X})
- \mathbf{V} akin to loadings (for variables of \mathbf{Y})
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})
- $\mathbf{F}_K = \mathbf{V}\Delta$ (component scores for variables of \mathbf{Y})

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values
- \mathbf{U} akin to loadings (for variables of \mathbf{X})
- \mathbf{V} akin to loadings (for variables of \mathbf{Y})
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})
- $\mathbf{F}_K = \mathbf{V}\Delta$ (component scores for variables of \mathbf{Y})
- $\mathbf{L}_X = \mathbf{XU}$ express the individuals w.r.t. \mathbf{X}

PLSC via the SVD

It's effectively just PCA with some new-ness:

- Δ are singular values
- \mathbf{U} akin to loadings (for variables of \mathbf{X})
- \mathbf{V} akin to loadings (for variables of \mathbf{Y})
- $\mathbf{F}_J = \mathbf{U}\Delta$ (component scores for variables of \mathbf{X})
- $\mathbf{F}_K = \mathbf{V}\Delta$ (component scores for variables of \mathbf{Y})
- $\mathbf{L}_X = \mathbf{XU}$ express the individuals w.r.t. \mathbf{X}
- $\mathbf{L}_Y = \mathbf{YV}$ express the individuals w.r.t. \mathbf{Y}

Example

PLS Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)

PLS Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)

PLS Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)

PLS Subjective Measures

	fruity	floral	vegetal	spicy
Chili_red_merlot	6	2	1	4
Chili_red_cabernet	5	3	2	3
Chili_red_shiraz	7	1	2	6
Canada_red_pinot	4	2	3	1
Canada_white_chardonnay	4	3	2	1
Canada_white_sauvignon	8	4	3	2
USA_rose_cabernet	8	3	3	3
USA_rose_pinot	6	1	1	2
USA_rose_syrah	9	3	2	5


PLS Objective Measures

	Acidity	Alcohol	Sugar	Tanin
Chili_red_merlot	5.33	13.8	2.75	559
Chili_red_cabernet	5.14	13.9	2.41	672
Chili_red_shiraz	5.16	14.3	2.20	455
Canada_red_pinot	5.70	13.3	1.70	320
Canada_white_chardonnay	6.00	13.5	3.00	35
Canada_white_sauvignon	7.50	12.0	3.50	40
USA_rose_cabernet	5.71	12.5	4.30	93
USA_rose_pinot	5.40	13.0	3.10	79
USA_rose_syrah	6.50	13.5	3.00	89

PLS - Wine

```
pls.res <- tepPLS(  
  DATA1 = wine.objective,  
  center1 = T, scale1 = T,  
  DATA2 = wine.subjective,  
  center2 = T, scale2 = T,  
  graphs=F  
)
```

PLS - Wine



PLS - Wine

While PLS is similar to PCA we should focus on what PLS maximizes:

PLS - Wine

While PLS is similar to PCA we should focus on what PLS maximizes:

$$\mathbf{L}_X^T \mathbf{L}_Y$$

PLS - Wine


While PLS is similar to PCA we should focus on what PLS maximizes:

$$\mathbf{L}_X^T \mathbf{L}_Y$$

So we'll start with the latent variable scores


PLS - Wine

PLS Wine Latent Variable Scores: LV1




PLS - Wine


PLS Wine Latent Variable Scores: LV1




PLS - Wine





PLS - Wine




PLS - Wine



PLS - Wine



PLS Wine Latent Variable Scores: LV1



Overview

We're experts now

- PCA is the core of PLS

We're experts now

- PCA is the core of PLS
- PLS generalizes PCA

We're experts now

- PCA is the core of PLS
- PLS generalizes PCA
 - If your two matrices are both the same, e.g., \mathbf{X}

We're experts now

- PCA is the core of PLS
- PLS generalizes PCA
 - If your two matrices are both the same, e.g., \mathbf{X}
 - PLS gives same results as PCA

We're experts now

- Visualize first

We're experts now

- Visualize first
- Use what you know to help construct the story from the numbers

We're experts now

- Visualize first
- Use what you know to help construct the story from the numbers
 - We'll see some additional helpers in next part today

Break!

We covered a lot!

- And we're not even at the good part yet!

We covered a lot!

- And we're not even at the good part yet!
- 7 ± 2 break

We covered a lot!

- And we're not even at the good part yet!
- 7 ± 2 break
 - Stretch, coffee

We covered a lot!

- And we're not even at the good part yet!
- 7 ± 2 break
 - Stretch, coffee
 - Pressing questions