

Simple & Multiple Correspondence Analyses

Contingency, categorical, ordinal, continuous and mixed data

Derek Beaton

Rotman Research Institute

October 28, 2019

Before we get started

Our new best friends

CONTINUOUS

measured data, can have ∞ values within possible range.



I AM 3.1" TALL
I WEIGH 34.16 grams

DISCRETE

OBSERVATIONS CAN ONLY EXIST
AT LIMITED VALUES, OFTEN
COUNTS.



I HAVE 8 LEGS
and
4 SPOTS!

@allison_horst

via @allison_horst

NOMINAL

UNORDERED DESCRIPTIONS



ORDINAL

ORDERED DESCRIPTIONS



BINARY

ONLY 2 MUTUALLY EXCLUSIVE OUTCOMES



@allison_horst

via @allison_horst

CONTINUOUS

measured data, can have oo values within possible range.



I AM 3.1" TALL
I WEIGH 34.16 grams

DISCRETE

observations can only exist at limited values, often counts.



I HAVE 8 LEGS
and
4 SPOTS!

@olivia_hart

NOMINAL

UNORDERED DESCRIPTIONS



-i'm a
TURTLE!

i'm a
snail!-



-i'm a
butterfly!

ORDINAL

ORDERED DESCRIPTIONS



-I am
unhappy



-I am
OK.



-I am
Awesome!!!

BINARY

ONLY 2 MUTUALLY
EXCLUSIVE OUTCOMES



I am
EXTINCT!



-HA.

@olivia_hart

► What do we do with all of these in a PCA like way?

CONTINUOUS

measured data, can have oo values within possible range.



I AM 3.1" TALL
I WEIGH 34.16 grams

DISCRETE

observations can only exist at limited values, often counts.



I HAVE 8 LEGS
and
4 SPOTS!

@olivia_hart

NOMINAL

UNORDERED DESCRIPTIONS



-i'm a
TURTLE!

i'm a
snail!-



-i'm a
butterfly!

ORDINAL

ORDERED DESCRIPTIONS



-I am
unhappy



-I am
OK.



-I am
Awesome!!!

BINARY

ONLY 2 MUTUALLY
EXCLUSIVE OUTCOMES



I AM
EXTINCT!



-HA.

@olivia_hart

- ▶ What do we do with all of these in a PCA like way?
- ▶ Some are very difficult and effectively ignored

CONTINUOUS

measured data, can have oo values within possible range.



I AM 3.1" TALL
I WEIGH 34.16 grams

DISCRETE

observations can only exist at limited values, often counts.



I HAVE 8 LEGS
and
4 SPOTS!

@olivia_hart

NOMINAL

UNORDERED DESCRIPTIONS



-i'm a
TURTLE!

i'm a
snail!-



-i'm a
butterfly!

ORDINAL

ORDERED DESCRIPTIONS



-I am
unhappy



-I am
OK.



-I am
Awesome!!!

BINARY

ONLY 2 MUTUALLY
EXCLUSIVE OUTCOMES



I AM
EXTINCT!



-HA.

@olivia_hart

- ▶ What do we do with all of these in a PCA like way?
- ▶ Some are very difficult and effectively ignored
 - ▶ We won't do that!

Motivation for today

- ▶ Not everything is a number

Motivation for today

- ▶ Not everything is a number
- ▶ Sometimes numbers aren't numbers!

Motivation for today

- ▶ Not everything is a number
- ▶ Sometimes numbers aren't numbers!
- ▶ We need to recognize when this happens

Motivation for today

- ▶ Not everything is a number
- ▶ Sometimes numbers aren't numbers!
- ▶ We need to recognize when this happens
 - ▶ And know what to do

Typology

- ▶ SS Stevens (not a boat!)

Typology

- ▶ SS Stevens (not a boat!)
- ▶ Levels of measurement

Typology

- ▶ SS Stevens (not a boat!)
- ▶ Levels of measurement
- ▶ Excellent examples:
https://en.wikipedia.org/wiki/Level_of_measurement

Where to find everything

- ▶ Generally: <https://github.com/derekbeaton/workshops>

Where to find everything

- ▶ Generally: <https://github.com/derekbeaton/workshops>
- ▶ Today:

Overview

- ▶ Revisit PCA

Overview

- ▶ Revisit PCA
- ▶ Looking at some data

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)
 - ▶ and how to handle various data types

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)
 - ▶ and how to handle various data types
- ▶ A whole bunch of bonuses

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)
 - ▶ and how to handle various data types
- ▶ A whole bunch of bonuses
 - ▶ Robustness

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)
 - ▶ and how to handle various data types
- ▶ A whole bunch of bonuses
 - ▶ Robustness
 - ▶ PLS

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)
 - ▶ and how to handle various data types
- ▶ A whole bunch of bonuses
 - ▶ Robustness
 - ▶ PLS
 - ▶ Networks

Overview

- ▶ Revisit PCA
- ▶ Looking at some data
- ▶ Simple correspondence analysis
 - ▶ and many of its connections
- ▶ Multiple correspondence analysis
 - ▶ generalizes CA (amongst many other things)
 - ▶ and how to handle various data types
- ▶ A whole bunch of bonuses
 - ▶ Robustness
 - ▶ PLS
 - ▶ Networks
 - ▶ Software

Revisiting PCA

What is PCA for?

- ▶ When we can compute a covariance or correlation matrix

What is PCA for?

- ▶ When we can compute a covariance or correlation matrix
- ▶ Break data into components

What is PCA for?

- ▶ When we can compute a covariance or correlation matrix
- ▶ Break data into components
 - ▶ Orthogonal


What is PCA for?

- ▶ When we can compute a covariance or correlation matrix
- ▶ Break data into components
 - ▶ Orthogonal
 - ▶ Rank ordered


What is PCA for?

- ▶ When we can compute a covariance or correlation matrix
- ▶ Break data into components
 - ▶ Orthogonal
 - ▶ Rank ordered
 - ▶ Made of bits & pieces of original measures


Eigen- and singular value decompositions




Eigen- and singular value decompositions




Eigen- and singular value decompositions



Eigen- and singular value decompositions



Eigen- and singular value decompositions



Left (row) singular
vectors

Some data


Diagnosis and education

	CN	Dementia	MCI
<i>ADV</i>	39	7	54
<i>B</i>	57	17	75
<i>B+</i>	75	19	113
<i>HS</i>	25	13	46
<i>HS+</i>	39	9	77

- ▶ Given a table, and asked for a multivariate analysis


- ▶ Given a table, and asked for a multivariate analysis
- ▶ We do what we know: PCA

PCA:
Row component scores




PCA:
Variable Component Correlations

PCA:
Row component scores



PCA:
Variable-Component Correlations



What did we analyze?

	CN	Dementia	MCI
CN	1.000	0.730	0.921
Dementia	0.730	1.000	0.652
MCI	0.921	0.652	1.000


What did PCA detect?

	CN	Dementia	MCI	<i>Row sums</i>
<i>ADV</i>	39	7	54	<i>100</i>
<i>B</i>	57	17	75	<i>149</i>
<i>B+</i>	75	19	113	<i>207</i>
<i>HS</i>	25	13	46	<i>84</i>
<i>HS+</i>	39	9	77	<i>125</i>


Let's try something different!

	ADV	B	B+	HS	HS+
<i>CN</i>	39	57	75	25	39
<i>Dementia</i>	7	17	19	13	9
<i>MCI</i>	54	75	113	46	77

PCA:
Row component scores



PCA:
Variable-Component Correlations



What did PCA analyze?

	ADV	B	B+	HS	HS+
ADV	1.000	1.000	0.995	0.935	0.963
B	1.000	1.000	0.994	0.932	0.960
B+	0.995	0.994	1.000	0.965	0.984
HS	0.935	0.932	0.965	1.000	0.996
HS+	0.963	0.960	0.984	0.996	1.000

What did PCA detect?

	ADV	B	B+	HS	HS+	<i>Row sums</i>
<i>CN</i>	39	57	75	25	39	235
<i>Dementia</i>	7	17	19	13	9	65
<i>MCI</i>	54	75	113	46	77	365

What is PCA for?

- ▶ When we can compute a *meaningful* covariance or correlation matrix

Let's take another look

	CN	Dementia	MCI	<i>Row sums</i>
<i>ADV</i>	39	7	54	100
<i>B</i>	57	17	75	149
<i>B+</i>	75	19	113	207
<i>HS</i>	25	13	46	84
<i>HS+</i>	39	9	77	125
Column sums	235	65	365	

- Tell me things about this matrix

Let's take another look

	CN	Dementia	MCI	<i>Row sums</i>
<i>ADV</i>	39	7	54	100
<i>B</i>	57	17	75	149
<i>B+</i>	75	19	113	207
<i>HS</i>	25	13	46	84
<i>HS+</i>	39	9	77	125
Column sums	235	65	365	

- ▶ Tell me things about this matrix
- ▶ What kind of problem does this look like?

Simple correspondence analysis

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature
 - ▶ From Benzecri (1964) & Escofier (1965)

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature
 - ▶ From Benzecri (1964) & Escofier (1965)
- ▶ Fully developed by Escofier (1969)

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature
 - ▶ From Benzecri (1964) & Escofier (1965)
- ▶ Fully developed by Escofier (1969)
- ▶ Explosion of the technique in France

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature
 - ▶ From Benzecri (1964) & Escofier (1965)
- ▶ Fully developed by Escofier (1969)
- ▶ Explosion of the technique in France
 - ▶ Across virtually every field (except psychology and neuroscience)

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature
 - ▶ From Benzecri (1964) & Escofier (1965)
- ▶ Fully developed by Escofier (1969)
- ▶ Explosion of the technique in France
 - ▶ Across virtually every field (except psychology and neuroscience)
- ▶ The magic of CA relies on the magic of χ^2

What is CA?

- ▶ Initially: *visualize contingency tables* (**a la PCA**, factor analyses)
 - ▶ Text (corpus) of philosophy, biblical passages, literature
 - ▶ From Benzecri (1964) & Escofier (1965)
- ▶ Fully developed by Escofier (1969)
- ▶ Explosion of the technique in France
 - ▶ Across virtually every field (except psychology and neuroscience)
- ▶ The magic of CA relies on the magic of χ^2
 - ▶ And there's some *crazy* magic here

History

- ▶ Hotelling (1933) & Thurstone (1933)

History

- ▶ Hotelling (1933) & Thurstone (1933)
- ▶ Hirschfeld (1935) & Horst (1935)

History

- ▶ Hotelling (1933) & Thurstone (1933)
- ▶ Hirschfeld (1935) & Horst (1935)
- ▶ Guttman (1941)

History

- ▶ Hotelling (1933) & Thurstone (1933)
- ▶ Hirschfeld (1935) & Horst (1935)
- ▶ Guttman (1941)
- ▶ Burt (1950)

History

- ▶ Hotelling (1933) & Thurstone (1933)
- ▶ Hirschfeld (1935) & Horst (1935)
- ▶ Guttman (1941)
- ▶ Burt (1950)
- ▶ And then Benzecri (1964) & Escofier (1965)

History

- ▶ Hotelling (1933) & Thurstone (1933)
- ▶ Hirschfeld (1935) & Horst (1935)
- ▶ Guttman (1941)
- ▶ Burt (1950)
- ▶ And then Benzecri (1964) & Escofier (1965)
- ▶ Many more very important characters to re-discover CA

- ▶ See Lebart's History & Prehistory of CA

- ▶ See Lebart's History & Prehistory of CA
 - ▶ http://www.dtmvic.com/doc/About_the_History_of_CA.pdf

- ▶ See Lebart's History & Prehistory of CA
 - ▶ http://www.dtmvic.com/doc/About_the_History_of_CA.pdf
- ▶ And Beh & Lombardo's series


- ▶ See Lebart's History & Prehistory of CA
 - ▶ http://www.dtmvic.com/doc/About_the_History_of_CA.pdf
- ▶ And Beh & Lombardo's series
 - ▶ A geneaology of CA:
<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-842X.2012.00676.x>

- ▶ See Lebart's History & Prehistory of CA
 - ▶ http://www.dtmvic.com/doc/About_the_History_of_CA.pdf
- ▶ And Beh & Lombardo's series
 - ▶ A geneaology of CA:
<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-842X.2012.00676.x>
 - ▶ A geneaology of CA 2: <http://siba-ese.unisalento.it/index.php/ejasa/article/view/19785>


We're diving in



	CN	Dementia	MCI
<i>ADV</i>	39	7	54
<i>B</i>	57	17	75
<i>B+</i>	75	19	113
<i>HS</i>	25	13	46
<i>HS+</i>	39	9	77

CA:
Row component scores



CA:
Column component scores







Want to see a cool trick?

	CN	Dementia	MCI
<i>ADV</i>	39	7	54
<i>B</i>	57	17	75
<i>B+</i>	75	19	113
<i>HS</i>	25	13	46
<i>HS+</i>	39	9	77

	ADV	B	B+	HS	HS+
<i>CN</i>	39	57	75	25	39
<i>Dementia</i>	7	17	19	13	9
<i>MCI</i>	54	75	113	46	77

What if we perform CA on this?



How did that happen?

Table 1: Data

	CN	Dementia	MCI
<i>ADV</i>	39	7	54
<i>B</i>	57	17	75
<i>B+</i>	75	19	113
<i>HS</i>	25	13	46
<i>HS+</i>	39	9	77

Table 2: Observed probabilities

	CN	Dementia	MCI
<i>ADV</i>	0.059	0.011	0.081
<i>B</i>	0.086	0.026	0.113
<i>B+</i>	0.113	0.029	0.170
<i>HS</i>	0.038	0.020	0.069
<i>HS+</i>	0.059	0.014	0.116

Table 3: Observed probabilities and margins

	CN	Dementia	MCI	<i>Row sums</i>
<i>ADV</i>	0.059	0.011	0.081	<i>0.150</i>
<i>B</i>	0.086	0.026	0.113	<i>0.224</i>
<i>B+</i>	0.113	0.029	0.170	<i>0.311</i>
<i>HS</i>	0.038	0.020	0.069	<i>0.126</i>
<i>HS+</i>	0.059	0.014	0.116	<i>0.188</i>
Column sums	<i>0.353</i>	<i>0.098</i>	<i>0.549</i>	

Table 4: Expected probabilities and margins

	CN	Dementia	MCI	<i>Row sums</i>
<i>ADV</i>	0.053	0.015	0.083	<i>0.150</i>
<i>B</i>	0.079	0.022	0.123	<i>0.224</i>
<i>B+</i>	0.110	0.030	0.171	<i>0.311</i>
<i>HS</i>	0.045	0.012	0.069	<i>0.126</i>
<i>HS+</i>	0.066	0.018	0.103	<i>0.188</i>
Column sums	<i>0.353</i>	<i>0.098</i>	<i>0.549</i>	

Table 5: Deviations: Observed - Expected

	CN	Dementia	MCI
<i>ADV</i>	0.006	-0.004	-0.001
<i>B</i>	0.007	0.004	-0.010
<i>B+</i>	0.003	-0.002	-0.001
<i>HS</i>	-0.007	0.007	0.000
<i>HS+</i>	-0.008	-0.005	0.013


Table 6: Row constraints (inverse row margins)

	ADV	B	B+	HS	HS+
<i>ADV</i>	6.65	0.000	0.000	0.000	0.00
<i>B</i>	0.00	4.463	0.000	0.000	0.00
<i>B+</i>	0.00	0.000	3.213	0.000	0.00
<i>HS</i>	0.00	0.000	0.000	7.917	0.00
<i>HS+</i>	0.00	0.000	0.000	0.000	5.32


Table 7: Column constraints (inverse column margins)

	CN	Dementia	MCI
<i>CN</i>	2.83	0.000	0.000
<i>Dementia</i>	0.00	10.231	0.000
<i>MCI</i>	0.00	0.000	1.822


What CA needs



R: Row constraints
(inverse row probabilities)



Z: Deviations



C: Column constraints
(inverse column probabilities)

► GSVD(**R**, **X**, **C**)

- ▶ GSVD(**R**, **X**, **C**)
- ▶ Uses but generalizes the SVD

- ▶ GSVD(**R**, **X**, **C**)
- ▶ Uses but generalizes the SVD
 - ▶ Uses row & column weights (constraints)

- ▶ GSVD(**R**, **X**, **C**)
- ▶ Uses but generalizes the SVD
 - ▶ Uses row & column weights (constraints)
- ▶ Gives back

- ▶ GSVD(**R**, **X**, **C**)
- ▶ Uses but generalizes the SVD
 - ▶ Uses row & column weights (constraints)
- ▶ Gives back
 - ▶ Component (factor) scores



- ▶ GSVD(**R**, **X**, **C**)
- ▶ Uses but generalizes the SVD
 - ▶ Uses row & column weights (constraints)
- ▶ Gives back
 - ▶ Component (factor) scores
 - ▶ Eigenvalues, singular values, & singular vectors

- ▶ GSVD(**R**, **X**, **C**)
- ▶ Uses but generalizes the SVD
 - ▶ Uses row & column weights (constraints)
- ▶ Gives back
 - ▶ Component (factor) scores
 - ▶ Eigenvalues, singular values, & singular vectors
 - ▶ *Generalized* singular vectors


What we really decompose



=




1/2





1/2

- A rectangle
- Deviations: Observed - Expected
 - Expected from Observed's margins



- Two squares
- Row margins and column margins



$$\frac{Z}{R^{\frac{1}{2}}C^{\frac{1}{2}}}$$

$$\frac{(O - E)}{E^{\frac{1}{2}}}$$

$$\chi^2 = \sum \frac{(\mathbf{O} - \mathbf{E})^2}{\mathbf{E}}$$

CA's first secrets

```
EDU <- amerge_subset$PTEDUCAT  
DX <- amerge_subset$DX  
edu_dx_table <- table(EDU, DX)
```

```
chisq.test(edu_dx_table)
```

```
##  
## Pearson's Chi-squared test  
##  
## data:  edu_dx_table  
## X-squared = 8.648, df = 8, p-value = 0.3729
```

```
edu_dx_ca <- epCA(edu_dx_table, graphs = F)  
sum(edu_dx_ca$ExPosition.Data$eigs) * sum(edu_dx_table)
```

```
## [1] 8.647979
```


The GSVD

Simple quick magic Then visualize it (as 3 matrices, then 1 over 2 which is just the probs not inverse) Then swing back to Chi2 Then swing to CCA Then expand it & transition to MCA

[[[pick up here and drop most of the stuff below]]]

The GSVD

- ▶ The generalized SVD

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix \mathbf{X} with I rows and J columns

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix \mathbf{X} with I rows and J columns
- ▶ $\text{SVD}(\mathbf{X})$ vs. $\text{GSVD}(\mathbf{W}_I, \mathbf{X}, \mathbf{W}_J)$

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix \mathbf{X} with I rows and J columns
- ▶ $\text{SVD}(\mathbf{X})$ vs. $\text{GSVD}(\mathbf{W}_I, \mathbf{X}, \mathbf{W}_J)$
- ▶ GSVD is

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix \mathbf{X} with I rows and J columns
- ▶ $\text{SVD}(\mathbf{X})$ vs. $\text{GSVD}(\mathbf{W}_I, \mathbf{X}, \mathbf{W}_J)$
- ▶ GSVD is
 - ▶ Matrix multiplication (by constraints on data)

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix \mathbf{X} with I rows and J columns
- ▶ $\text{SVD}(\mathbf{X})$ vs. $\text{GSVD}(\mathbf{W}_I, \mathbf{X}, \mathbf{W}_J)$
- ▶ GSVD is
 - ▶ Matrix multiplication (by constraints on data)
 - ▶ The SVD

The GSVD

- ▶ The generalized SVD
 - ▶ Constraints (weights) for rows & columns of rectangular table
 - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix \mathbf{X} with I rows and J columns
- ▶ $\text{SVD}(\mathbf{X})$ vs. $\text{GSVD}(\mathbf{W}_I, \mathbf{X}, \mathbf{W}_J)$
- ▶ GSVD is
 - ▶ Matrix multiplication (by constraints on data)
 - ▶ The SVD
 - ▶ More matrix multiplication (by constraints on vectors)

What we did to the data

- ▶ O , w_i & w_j , E , Z

What we did to the data

- ▶ O , w_i & w_j , E , Z
- ▶ Oh look that's χ^2

What we did to the data

- ▶ O , w_i & w_j , E , Z
- ▶ Oh look that's χ^2
- ▶ Sum of eigenvalues * sum of table = χ^2 .

What we did to the data

- ▶ O , w_i & w_j , E , Z
- ▶ Oh look that's χ^2
- ▶ Sum of eigenvalues * sum of table = χ^2 .
 - ▶ Each component is an additive orthogonal slice of χ^2 . WOAHH.

What we did to the data

- ▶ O , w_i & w_j , E , Z
- ▶ Oh look that's Chi^2
- ▶ Sum of eigenvalues * sum of table = Chi^2 .
 - ▶ Each component is an additive orthogonal slice of Chi^2 . WOAHH.
 - ▶ The eigenvalues are *magic*

CA visualized

- ▶ Oh look it's CCA-ish

CA visualized

- ▶ Oh look it's CCA-ish
- ▶ Oh it really really is CCA-ish!

Rules

- ▶ It's like PCA

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores
- ▶ It's unlike PCA

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores
- ▶ It's unlike PCA
 - ▶ Relative interpretation *between* sets

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores
- ▶ It's unlike PCA
 - ▶ Relative interpretation *between* sets
 - ▶ Generalized singular vectors

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores
- ▶ It's unlike PCA
 - ▶ Relative interpretation *between* sets
 - ▶ Generalized singular vectors
 - ▶ Bifactor

Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores
- ▶ It's unlike PCA
 - ▶ Relative interpretation *between* sets
 - ▶ Generalized singular vectors
 - ▶ Bifactor
 - ▶ Rows & columns treated the same


Rules

- ▶ It's like PCA
 - ▶ Variance (singular values, eigen values)
 - ▶ Directions & inter-relationships
 - ▶ Components scores
- ▶ It's unlike PCA
 - ▶ Relative interpretation *between* sets
 - ▶ Generalized singular vectors
 - ▶ Bifactor
 - ▶ Rows & columns treated the same
 - ▶ Together they help make components, as opposed to PCA


Multiple correspondence analysis

MAGIC!

Chi-squared



Chi-squared



See [here](#)

Some many bonuses!

(Some) References

See the reference sections of these

- ▶ Beaton, D., Saporta, G., Abdi, H., & Alzheimer's Disease Neuroimaging Initiative. (2019). A generalization of partial least squares regression and correspondence analysis for categorical and mixed data: An application with the ADNI data. bioRxiv, 598888.

See the reference sections of these

- ▶ Beaton, D., Saporta, G., Abdi, H., & Alzheimer's Disease Neuroimaging Initiative. (2019). A generalization of partial least squares regression and correspondence analysis for categorical and mixed data: An application with the ADNI data. bioRxiv, 598888.
- ▶ Beaton, D., Sunderland, K. M., Levine, B., Mandzia, J., Masellis, M., Swartz, R. H., ... & Strother, S. C. (2019). Generalization of the minimum covariance determinant algorithm for categorical and mixed data types. bioRxiv, 333005.

And these

- ▶ Abdi, H., Guillemot, V., Eslami, A., & Beaton, D. (2017). Canonical correlation analysis. Encyclopedia of Social Network Analysis and Mining, 1-16.

And these

- ▶ Abdi, H., Guillemot, V., Eslami, A., & Beaton, D. (2017). Canonical correlation analysis. Encyclopedia of Social Network Analysis and Mining, 1-16.
- ▶ Beaton, D., Dunlop, J., & Abdi, H. (2016). Partial least squares correspondence analysis: A framework to simultaneously analyze behavioral and genetic data. Psychological methods, 21(4), 621.

Techniques

- ▶ Greenacre, M. (2017). Correspondence analysis in practice. CRC press.

Techniques

- ▶ Greenacre, M. (2017). Correspondence analysis in practice. CRC press.
- ▶ Greenacre, M. J. (1984). Theory and Applications of Correspondence Analysis. Retrieved from <http://books.google.com/books?id=LsPaAAAAMAAJ>

Techniques

- ▶ Greenacre, M. J. (2010). Correspondence analysis. Wiley Interdisciplinary Reviews: Computational Statistics, 2(5), 613–619. <https://doi.org/10.1002/wics.114>

Techniques

- ▶ Greenacre, M. J. (2010). Correspondence analysis. Wiley Interdisciplinary Reviews: Computational Statistics, 2(5), 613–619. <https://doi.org/10.1002/wics.114>
- ▶ Lebart, L., Morineau, A., & Warwick, K. M. (1984). Multivariate descriptive statistical analysis: correspondence analysis and related techniques for large matrices. Wiley.

Techniques

- ▶ Greenacre, M. J. (2010). Correspondence analysis. Wiley Interdisciplinary Reviews: Computational Statistics, 2(5), 613–619. <https://doi.org/10.1002/wics.114>
- ▶ Lebart, L., Morineau, A., & Warwick, K. M. (1984). Multivariate descriptive statistical analysis: correspondence analysis and related techniques for large matrices. Wiley.
- ▶ Nguyen, L. H., & Holmes, S. (2019). Ten quick tips for effective dimensionality reduction. PLOS Computational Biology, 15(6), e1006907.

Data

- ▶ Escofier, B. (1978). Analyse factorielle et distances répondant au principe d'équivalence distributionnelle. *Revue de Statistique Appliquée*, 26(4), 29–37.

Data

- ▶ Escofier, B. (1978). Analyse factorielle et distances répondant au principe d'équivalence distributionnelle. *Revue de Statistique Appliquée*, 26(4), 29–37.
- ▶ Escofier, B. (1979). Traitement simultané de variables qualitatives et quantitatives en analyse factorielle. *Cahiers de l'Analyse Des Données*, 4(2), 137–146.

Data

- ▶ Escofier, B. (1978). Analyse factorielle et distances répondant au principe d'équivalence distributionnelle. *Revue de Statistique Appliquée*, 26(4), 29–37.
- ▶ Escofier, B. (1979). Traitement simultané de variables qualitatives et quantitatives en analyse factorielle. *Cahiers de l'Analyse Des Données*, 4(2), 137–146.
- ▶ Greenacre, M. (2014). Data Doubling and Fuzzy Coding. In J. Blasius & M. Greenacre (Eds.), *Visualization and Verbalization of Data* (pp. 239–253). Philadelphia, PA, USA: CRC Press.

History

- ▶ Holmes S, Josse J. Discussion of “50 Years of Data Science”. Journal of Computational and Graphical Statistics. 2017, V26(4) 768-769. <https://www.tandfonline.com/doi/full/10.1080/10618600.2017.1385471>