

# Almost everything you need to know about PLS

## Part 1: Background, Theory, and Examples

Jenny Rieck & Derek Beaton

October 24, 2017

# The BIG outline

- Part 1: Background & Examples

# The BIG outline

- Part 1: Background & Examples
  - RIGHT NOW

# The BIG outline

- Part 1: Background & Examples
  - RIGHT NOW
  - Introduce everything we need

# The BIG outline

- Part 1: Background & Examples
  - RIGHT NOW
  - Introduce everything we need
- Part 2: PLS in Matlab & R

# The BIG outline

- Part 1: Background & Examples
  - RIGHT NOW
  - Introduce everything we need
- Part 2: PLS in Matlab & R
  - Tuesday November 21, 10:00-12:00 Worstman Hall

# The BIG outline

- Part 1: Background & Examples
  - RIGHT NOW
  - Introduce everything we need
- Part 2: PLS in Matlab & R
  - Tuesday November 21, 10:00-12:00 Worstman Hall
  - Put knowledge into practice

## Part 1 outline

- Theory & Background



## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)
- Short break

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)
- Short break
- Examples (with ADNI)

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
  - Standard PLS (& inference)

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
  - Standard PLS (& inference)
  - Discriminant PLS

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
  - Standard PLS (& inference)
  - Discriminant PLS
  - “Seed” PLS

## Part 1 outline

- Theory & Background
  - Principal component analysis (PCA)
  - Partial least squares (PLS)
- Short break
- Examples (with ADNI)
  - Standard PLS (& inference)
  - Discriminant PLS
  - “Seed” PLS
- And beyond!



# Principal Component Analysis

## Background

# PCA Pet peeve

- PCA has *no principles*

# PCA Pet peeve

- PCA has *no principles*
- PCA is your *pal*

# PCA Pet peeve

- PCA has *no principles*
- PCA is your *pal*
- Don't be part of the 3.5% on Google Scholar

# PCA Pet peeve

- PCA has *no principles*
- PCA is your *pal*
- Don't be part of the 3.5% on Google Scholar
  - 70,400 (“principle”)

# PCA Pet peeve

- PCA has *no principles*
- PCA is your *pal*
- Don't be part of the 3.5% on Google Scholar
  - 70,400 (“principle”)
  - 1,940,000 (“principal”)

# Where did PCA come from?

- Modern form



# Where did PCA come from?

- Modern form
  - Hotelling (1933)

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)
  - Galton (1859)

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)
  - Galton (1859)
  - Pearson (1901)

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)
  - Galton (1859)
  - Pearson (1901)
- Modern overviews

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)
  - Galton (1859)
  - Pearson (1901)
- Modern overviews
  - S. Wold et al., (1987)

# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)
  - Galton (1859)
  - Pearson (1901)
- Modern overviews
  - S. Wold et al., (1987)
  - Jolliffe (2002)



# Where did PCA come from?

- Modern form
  - Hotelling (1933)
- Traces back to
  - Cauchy (1829)
  - Galton (1859)
  - Pearson (1901)
- Modern overviews
  - S. Wold et al., (1987)
  - Jolliffe (2002)
  - Abdi & Williams (2010)

# What is PCA?

- Visualize high dimensional data

# What is PCA?

- Visualize high dimensional data
- Orthogonal transformation

# What is PCA?

- Visualize high dimensional data
- Orthogonal transformation
- Dimensionality reduction

# What is PCA?

- Find “components”

# What is PCA?

- Find “components”
  - Components are new variables that are combinations of old variables

# What is PCA?

- Find “components”
  - Components are new variables that are combinations of old variables
- Components explain maximum possible variance

# What is PCA?

- Find “components”
  - Components are new variables that are combinations of old variables
- Components explain maximum possible variance
  - Conditional to orthogonality



## Visual example






Figure 1: The kind of data we usually expect for PCA


# Visual example




# Visual example




# Visual example




# Visual example




# Visual example



# Visual example




# Visual example






# Visual example



# Visual example



# What is PCA?

- The basis of modern techniques

# What is PCA?

- The basis of modern techniques
  - Factor analyses

# What is PCA?

- The basis of modern techniques
  - Factor analyses
  - Independent components analysis

# What is PCA?

- The basis of modern techniques
  - Factor analyses
  - Independent components analysis
  - Partial least squares

# What is PCA?

- The basis of modern techniques
  - Factor analyses
  - Independent components analysis
  - Partial least squares
  - Discriminant analyses

# What is PCA?

- The basis of modern techniques
  - Factor analyses
  - Independent components analysis
  - Partial least squares
  - Discriminant analyses
  - Multi-table (e.g., MFA, GCCA)



# What is PCA?

- A special case of the singular value decomposition (SVD)

# What is PCA?

- A special case of the singular value decomposition (SVD)
- Which means (almost) everything else is, too

## Formalization

# Singular value decomposition

The SVD is one of the most ubiquitous and important tools

# SVD

- We'll go into *enough* formalization

# SVD

- We'll go into *enough* formalization
- If you want more:

# SVD

- We'll go into *enough* formalization
- If you want more:
  - S. Wold et al., (1987)

# SVD

- We'll go into *enough* formalization
- If you want more:
  - S. Wold et al., (1987)
  - Jolliffe (2002)



# SVD

- We'll go into *enough* formalization
- If you want more:
  - S. Wold et al., (1987)
  - Jolliffe (2002)
  - Abdi & Williams (2010)

# SVD

- We'll go into *enough* formalization
- If you want more:
  - S. Wold et al., (1987)
  - Jolliffe (2002)
  - Abdi & Williams (2010)
  - And many others...

# SVD



Figure 2: The shape of the data

# SVD




Figure 3: SVD breaks down the data

# SVD

## Notation

# SVD

## Notation

- $x$  - a scalar

# SVD

## Notation

- $x$  - a scalar
- $\mathbf{a}$  - a vector

# SVD

## Notation

- $x$  - a scalar
- $\mathbf{a}$  - a vector
- $\mathbf{A}$  - a matrix



# SVD

## Notation

- $x$  - a scalar
- $\mathbf{a}$  - a vector
- $\mathbf{A}$  - a matrix
- $\mathbf{A}^T$  - transpose

# SVD

## Notation

- $x$  - a scalar
- $\mathbf{a}$  - a vector
- $\mathbf{A}$  - a matrix
- $\mathbf{A}^T$  - transpose
- $\mathbf{AB}$  - multiplication

# SVD

- Think back to PCA

# SVD

- Think back to PCA
- We want to find *the* principal component

# SVD

- Think back to PCA
- We want to find *the* principal component
  - a.k.a. maximum source of variance

# SVD

Given a matrix  $\mathbf{X}$  we generally assume that

# SVD

Given a matrix  $\mathbf{X}$  we generally assume that

- column-wise centered

# SVD

Given a matrix  $\mathbf{X}$  we generally assume that

- column-wise centered
- column-wise scaled (e.g., z-scores or sums of squares = 1)



# SVD

Given  $\mathbf{X}$  of size  $I \times J$

# SVD

Given  $\mathbf{X}$  of size  $I \times J$

We want to find vectors

# SVD

Given  $\mathbf{X}$  of size  $I \times J$

We want to find vectors

- $\mathbf{u}$  of size  $I \times 1$
- $\mathbf{v}$  of size  $J \times 1$

# SVD

Given  $\mathbf{X}$  of size  $I \times J$

We want to find vectors

- $\mathbf{u}$  of size  $I \times 1$
- $\mathbf{v}$  of size  $J \times 1$

such that

$$\operatorname{argmax}_{\mathbf{u}, \mathbf{v}} \delta = \mathbf{u}^T \mathbf{X} \mathbf{v} \text{ conditional to } \mathbf{u}^T \mathbf{u} = 1 = \mathbf{v}^T \mathbf{v}$$

# SVD

Which gives us the following equivalencies:

- $\mathbf{X}\mathbf{v} = \mathbf{u}\delta$
- $\mathbf{X}^T \mathbf{u} = \mathbf{v}\delta$

# SVD

Which gives us the following equivalencies:

- $\mathbf{X}\mathbf{v} = \mathbf{u}\delta$
- $\mathbf{X}^T\mathbf{u} = \mathbf{v}\delta$

where

- $\mathbf{X}_1 = \delta\mathbf{u}\mathbf{v}^T$

# SVD

Which gives us the following equivalencies:

- $\mathbf{X}\mathbf{v} = \mathbf{u}\delta$
- $\mathbf{X}^T\mathbf{u} = \mathbf{v}\delta$

where

- $\mathbf{X}_1 = \delta\mathbf{u}\mathbf{v}^T$ 
  - $\mathbf{X}_1$  is  $\mathbf{X}$  as represented by *source of maximum variance*

# SVD

Given  $\mathbf{X}_1 = \delta \mathbf{p} \mathbf{q}^T$  we can find the other components



# SVD

Given  $\mathbf{X}_1 = \delta \mathbf{p} \mathbf{q}^T$  we can find the other components

- $\mathbf{X}_{2:L} = \mathbf{X} - \mathbf{X}_1$
- $\mathbf{X}_{2:L}$  is orthogonal to  $\mathbf{X}_1$

## SVD

Given  $\mathbf{X}_1 = \delta \mathbf{p} \mathbf{q}^T$  we can find the other components

- $\mathbf{X}_{2:L} = \mathbf{X} - \mathbf{X}_1$
- $\mathbf{X}_{2:L}$  is orthogonal to  $\mathbf{X}_1$

$$\underset{\mathbf{p}, \mathbf{q}}{\operatorname{argmax}} \delta = \mathbf{p}^T \mathbf{X}_{2:L} \mathbf{q} \text{ conditional to } \mathbf{p}^T \mathbf{p} = 1 = \mathbf{q}^T \mathbf{q}$$

# SVD

The SVD of  $\mathbf{X}$  of size  $I \times J$ :

# SVD

The SVD of  $\mathbf{X}$  of size  $I \times J$ :

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \quad (1)$$

# SVD

The SVD of  $\mathbf{X}$  of size  $I \times J$ :

$$\mathbf{X} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^T \quad (1)$$

$\mathbf{U}$  and  $\mathbf{V}$  are orthonormal such that

# SVD

The SVD of  $\mathbf{X}$  of size  $I \times J$ :

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \quad (1)$$

$\mathbf{U}$  and  $\mathbf{V}$  are orthonormal such that

$$\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V} \quad (2)$$

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- where rank is  $L$  (i.e., number of extractable components)



# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- where rank is  $L$  (i.e., number of extractable components)
- $\mathbf{U}$  is  $I \times L$  (left singular vectors; rows of  $\mathbf{X}$ )

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- where rank is  $L$  (i.e., number of extractable components)
- $\mathbf{U}$  is  $I \times L$  (left singular vectors; rows of  $\mathbf{X}$ )
- $\mathbf{V}$  is  $J \times L$  (right singular vectors; columns of  $\mathbf{X}$ )

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{\Delta}$  is  $L \times L$  diagonal matrix

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{\Delta}$  is  $L \times L$  diagonal matrix
- $\text{diag}\{\mathbf{\Delta}\} = \boldsymbol{\delta}$  are singular values (decreasing)

# SVD

$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T$  such that  $\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$

- $\mathbf{\Delta}$  is  $L \times L$  diagonal matrix
- $\text{diag}\{\mathbf{\Delta}\} = \boldsymbol{\delta}$  are singular values (decreasing)
- $\lambda = \delta^2$  are the eigenvalues (variance)

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

# SVD

$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T$  such that  $\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$

- $\mathbf{x}_1 = \mathbf{U}_1\mathbf{\Delta}_1\mathbf{V}_1^T$



# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{X}_1 = \mathbf{U}_1\mathbf{\Delta}_1\mathbf{V}_1^T$
- $\mathbf{X}_{2:L} = \mathbf{U}_{2:L}\mathbf{\Delta}_{2:L}\mathbf{V}_{2:L}^T$

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta}$  (row component scores)

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta}$  (row component scores)
- $\mathbf{F}_J = \mathbf{V}\mathbf{\Delta}$  (column component scores)

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

# SVD

$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T$  such that  $\mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta} = \mathbf{X}\mathbf{V}$

# SVD

$$\mathbf{X} = \mathbf{U}\mathbf{\Delta}\mathbf{V}^T \text{ such that } \mathbf{U}^T\mathbf{U} = \mathbf{I} = \mathbf{V}^T\mathbf{V}$$

- $\mathbf{F}_I = \mathbf{U}\mathbf{\Delta} = \mathbf{X}\mathbf{V}$
- $\mathbf{F}_J = \mathbf{V}\mathbf{\Delta} = \mathbf{X}^T\mathbf{U}$

# SVD

- Phew.



# SVD

- Phew.
- Enough nerd stuff

# SVD

- Phew.
- Enough nerd stuff
- Let's get back to PCA

## Toy example

- In R with ExPosition packages

- In R with ExPosition packages
  - Sneak preview of newest version

- In R with ExPosition packages
  - Sneak preview of newest version
- <https://github.com/derekbeaton/ExPosition-Family/>

- In R with ExPosition packages
  - Sneak preview of newest version
- <https://github.com/derekbeaton/ExPosition-Family/>
- <https://cran.r-project.org/web/packages/ExPosition/index.html>

## PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)



## PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)

## PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)

# PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)
  - We'll use this one for PCA

## PCA Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)
  - We’ll use this one for PCA
  - We’ll use both for PLS

# PCA Subjective Measures

	fruity	floral	vegetal	spicy
Chili_red_merlot	6	2	1	4
Chili_red_cabernet	5	3	2	3
Chili_red_shiraz	7	1	2	6
Canada_red_pinot	4	2	3	1
Canada_white_chardonnay	4	3	2	1
Canada_white_sauvignon	8	4	3	2
USA_rose_cabernet	8	3	3	3
USA_rose_pinot	6	1	1	2
USA_rose_syrah	9	3	2	5

## PCA - Wine

```
pca.res <- epPCA(  
  wine.subjective,  
  scale = T,  
  center = T,  
  graphs = F)
```

# PCA - Wine

- PCA maximizes variance

# PCA - Wine

- PCA maximizes variance
- So let's visualize the variance per component



# PCA - Wine

- PCA maximizes variance
- So let's visualize the variance per component
  - Use the eigenvalues

# PCA - Wine

- PCA maximizes variance
- So let's visualize the variance per component
  - Use the eigenvalues
  - "Scree plot"

# PCA - Wine

- Many ways to present the results

# PCA - Wine

- Many ways to present the results
- We prioritize visualization

# PCA - Wine

- Many ways to present the results
- We prioritize visualization
  - Use numbers and tables when needed

## PCA - Wine

- Many ways to present the results
- We prioritize visualization
  - Use numbers and tables when needed
- PCA is half

# PCA - Wine


- Many ways to present the results
- We prioritize visualization
  - Use numbers and tables when needed
- PCA is half
  - Stats

# PCA - Wine


- Many ways to present the results
- We prioritize visualization
  - Use numbers and tables when needed
- PCA is half
  - Stats
  - Art




# PCA - Wine




# PCA - Wine



# PCA - Wine



# PCA - Wine



# PCA - Wine




Figure 4: Variables & Observations

# PCA

- If you know PCA you know about 90% of the multivariate stats in use

# Partial least squares

## Background



# PLS means a lot of things

- Projection onto latent structures

# PLS means a lot of things

- Projection onto latent structures
  - Probably the most accurate name

# PLS means a lot of things

- Projection onto latent structures
  - Probably the most accurate name
  - But also probably too broad a definition

# PLS means a lot of things

Partial least squares sounds like ordinary least squares

- When we have two matrices:  $\mathbf{X}$  and  $\mathbf{Y}$

# PLS means a lot of things

Partial least squares sounds like ordinary least squares

- When we have two matrices:  $\mathbf{X}$  and  $\mathbf{Y}$
- OLS:  $(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$

# PLS means a lot of things

Partial least squares sounds like ordinary least squares

- When we have two matrices:  $\mathbf{X}$  and  $\mathbf{Y}$
- OLS:  $(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$
- PLS:  $\mathbf{X}^T \mathbf{Y}$

# PLS means a lot of things

- Partial least squares path modelling (PLS-PM)

# PLS means a lot of things

- Partial least squares path modelling (PLS-PM)
- Partial least squares regression (PLSR)



# PLS means a lot of things

- Partial least squares path modelling (PLS-PM)
- Partial least squares regression (PLSR)
- Partial least squares correlation (PLSC)

# PLS means a lot of things

- Partial least squares path modelling (PLS-PM)
- Partial least squares regression (PLSR)
- Partial least squares correlation (PLSC)
  - This is the one we'll talk about today

# PLSC has many...

## Names

- Inter-battery (factor) analysis (Tucker, 1958)

# PLSC has many...

## Names

- Inter-battery (factor) analysis (Tucker, 1958)
- Covariance between two fields (Bretherton, Smith, & Wallace, 1992)

# PLSC has many...

## Names

- Inter-battery (factor) analysis (Tucker, 1958)
- Covariance between two fields (Bretherton, Smith, & Wallace, 1992)
- PLS-SVD (Tenenhaus, 2005)

# PLSC has many...

## Names

- Inter-battery (factor) analysis (Tucker, 1958)
- Covariance between two fields (Bretherton, Smith, & Wallace, 1992)
- PLS-SVD (Tenenhaus, 2005)
- co-inertia analysis (Dray, 2014)

# PLSC has many...

Friends

- Reduced Rank Regression

# PLSC has many...

## Friends

- Reduced Rank Regression
- Canonical Correlation Analysis



# PLSC has many...

## Friends

- Reduced Rank Regression
- Canonical Correlation Analysis
- (Fisher's) Linear Discriminant Analysis

# PLSC has many...

## Friends

- Reduced Rank Regression
- Canonical Correlation Analysis
- (Fisher's) Linear Discriminant Analysis
- PLS-correspondence analysis

# PLSC

## History

- McIntosh, Bookstein, Haxby, & Grady (1996)

# PLSC

## History

- McIntosh, Bookstein, Haxby, & Grady (1996)
- Bookstein (1992)

# PLSC

## History

- McIntosh, Bookstein, Haxby, & Grady (1996)
- Bookstein (1992)
- Tucker (1958)

# PLSC

Modern overviews

- McIntosh & Lobaugh (2004)

# PLSC

## Modern overviews

- McIntosh & Lobaugh (2004)
- Krishnan et al., (2011)

# PLSC

It's effectively just PCA applied to the cross product of two matrices measured on the same observations:



# PLSC

It's effectively just PCA applied to the cross product of two matrices measured on the same observations:

- $\mathbf{X}$  which is  $I \times J$

# PLSC

It's effectively just PCA applied to the cross product of two matrices measured on the same observations:

- **X** which is  $I \times J$
- **Y** which is  $I \times K$

## Formalization

## PLSC via the SVD

Given  $\mathbf{X}$  of size  $I \times J$  and  $\mathbf{Y}$  of size  $I \times K$

## PLSC via the SVD

Given  $\mathbf{X}$  of size  $I \times J$  and  $\mathbf{Y}$  of size  $I \times K$

We want to find vectors

## PLSC via the SVD

Given  $\mathbf{X}$  of size  $I \times J$  and  $\mathbf{Y}$  of size  $I \times K$

We want to find vectors

- $\mathbf{u}$  of size  $J \times 1$
- $\mathbf{v}$  of size  $K \times 1$

To define latent variables

## PLSC via the SVD

Given  $\mathbf{X}$  of size  $I \times J$  and  $\mathbf{Y}$  of size  $I \times K$

We want to find vectors

- $\mathbf{u}$  of size  $J \times 1$
- $\mathbf{v}$  of size  $K \times 1$

To define latent variables

- $\mathbf{l}_X = \mathbf{X}\mathbf{u}$  of size  $I \times 1$
- $\mathbf{l}_Y = \mathbf{Y}\mathbf{v}$  of size  $I \times 1$

# PLSC via the SVD

Given  $\mathbf{X}$  of size  $I \times J$  and  $\mathbf{Y}$  of size  $I \times K$

We want to find vectors

- $\mathbf{u}$  of size  $J \times 1$
- $\mathbf{v}$  of size  $K \times 1$

To define latent variables

- $\mathbf{l}_X = \mathbf{X}\mathbf{u}$  of size  $I \times 1$
- $\mathbf{l}_Y = \mathbf{Y}\mathbf{v}$  of size  $I \times 1$

such that

$$\operatorname{argmax}_{\mathbf{u}, \mathbf{v}} \delta = \mathbf{u}^T \mathbf{X}^T \mathbf{Y} \mathbf{v} \text{ conditional to } \mathbf{u}^T \mathbf{u} = 1 = \mathbf{v}^T \mathbf{v}$$



## PLSC via the SVD

Compute the relationship between  $\mathbf{X}$  and  $\mathbf{Y}$

$$\mathbf{R} = \mathbf{X}^T \mathbf{Y} \quad (3)$$

## PLSC via the SVD

Compute the relationship between  $\mathbf{X}$  and  $\mathbf{Y}$

$$\mathbf{R} = \mathbf{X}^T \mathbf{Y} \quad (3)$$

Compute the SVD of  $\mathbf{R}$

$$\mathbf{R} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^T \quad (4)$$

# PLSC via the SVD

Compute the relationship between  $\mathbf{X}$  and  $\mathbf{Y}$

$$\mathbf{R} = \mathbf{X}^T \mathbf{Y} \quad (3)$$

Compute the SVD of  $\mathbf{R}$

$$\mathbf{R} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^T \quad (4)$$

Compute the latent variables

$$\mathbf{L}_\mathbf{X} = \mathbf{X} \mathbf{U} \text{ and } \mathbf{L}_\mathbf{Y} = \mathbf{Y} \mathbf{V} \quad (5)$$

# PLSC via the SVD

Almost everything is the same:

- $\Delta$  are singular values

## PLSC via the SVD

Almost everything is the same:

- $\Delta$  are singular values
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )

# PLSC via the SVD

Almost everything is the same:

- $\Delta$  are singular values
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )
- $\mathbf{F}_K = \mathbf{V}\Delta$  (component scores for variables of  $\mathbf{Y}$ )

## PLSC via the SVD

Almost everything is the same:

- $\Delta$  are singular values
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )
- $\mathbf{F}_K = \mathbf{V}\Delta$  (component scores for variables of  $\mathbf{Y}$ )
- So our nomenclature will align with PCA

# PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$  express the individuals w.r.t.  $\mathbf{X}$



# PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$  express the individuals w.r.t.  $\mathbf{X}$
- $\mathbf{L}_Y = \mathbf{Y}\mathbf{V}$  express the individuals w.r.t.  $\mathbf{Y}$

# PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$  express the individuals w.r.t.  $\mathbf{X}$
- $\mathbf{L}_Y = \mathbf{Y}\mathbf{V}$  express the individuals w.r.t.  $\mathbf{Y}$
- Not in PCA

# PLSC via the SVD

The new-ness

- $\mathbf{L}_X = \mathbf{X}\mathbf{U}$  express the individuals w.r.t.  $\mathbf{X}$
- $\mathbf{L}_Y = \mathbf{Y}\mathbf{V}$  express the individuals w.r.t.  $\mathbf{Y}$
- Not in PCA
  - We'll call these "latent variable scores"

# PLSC via the SVD

Maximizes the latent variables

# PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

# PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \mathbf{\Delta} \quad (6)$$

When expanded

$$(\mathbf{XU})^T (\mathbf{YV}) = \mathbf{\Delta}$$

# PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

# PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{R} \mathbf{V} = \Delta$$



# PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{R} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{U} \Delta \mathbf{V}^T \mathbf{V} = \Delta$$

# PLSC via the SVD

Maximizes the latent variables

$$\mathbf{L}_X^T \mathbf{L}_Y = \Delta \quad (6)$$

When expanded

$$(\mathbf{X}\mathbf{U})^T(\mathbf{Y}\mathbf{V}) = \Delta$$

$$\mathbf{U}^T \mathbf{X}^T \mathbf{Y} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{R} \mathbf{V} = \Delta$$

$$\mathbf{U}^T \mathbf{U} \Delta \mathbf{V}^T \mathbf{V} = \Delta$$

because

$$\mathbf{U}^T \mathbf{U} = \mathbf{I} = \mathbf{V}^T \mathbf{V}$$

# PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values

# PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values
- $\mathbf{U}$  akin to loadings (for variables of  $\mathbf{X}$ )

# PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values
- $\mathbf{U}$  akin to loadings (for variables of  $\mathbf{X}$ )
- $\mathbf{V}$  akin to loadings (for variables of  $\mathbf{Y}$ )

# PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values
- $\mathbf{U}$  akin to loadings (for variables of  $\mathbf{X}$ )
- $\mathbf{V}$  akin to loadings (for variables of  $\mathbf{Y}$ )
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )

## PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values
- $\mathbf{U}$  akin to loadings (for variables of  $\mathbf{X}$ )
- $\mathbf{V}$  akin to loadings (for variables of  $\mathbf{Y}$ )
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )
- $\mathbf{F}_K = \mathbf{V}\Delta$  (component scores for variables of  $\mathbf{Y}$ )

# PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values
- $\mathbf{U}$  akin to loadings (for variables of  $\mathbf{X}$ )
- $\mathbf{V}$  akin to loadings (for variables of  $\mathbf{Y}$ )
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )
- $\mathbf{F}_K = \mathbf{V}\Delta$  (component scores for variables of  $\mathbf{Y}$ )
- $\mathbf{L}_X = \mathbf{XU}$  express the individuals w.r.t.  $\mathbf{X}$



# PLSC via the SVD

It's effectively just PCA with some new-ness:

- $\Delta$  are singular values
- $\mathbf{U}$  akin to loadings (for variables of  $\mathbf{X}$ )
- $\mathbf{V}$  akin to loadings (for variables of  $\mathbf{Y}$ )
- $\mathbf{F}_J = \mathbf{U}\Delta$  (component scores for variables of  $\mathbf{X}$ )
- $\mathbf{F}_K = \mathbf{V}\Delta$  (component scores for variables of  $\mathbf{Y}$ )
- $\mathbf{L}_X = \mathbf{XU}$  express the individuals w.r.t.  $\mathbf{X}$
- $\mathbf{L}_Y = \mathbf{YV}$  express the individuals w.r.t.  $\mathbf{Y}$

## Example

## PLS Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)

## PLS Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)

## PLS Toy Dataset - Wine

- 36 different wines (e.g., USA red cab., CAN rose syrah)
- 4 objective measures (e.g., “Alcohol”, “Acidity”)
- 9 subjective measures (e.g., “sweet”, “acidic”)

# PLS Subjective Measures

	fruity	floral	vegetal	spicy
Chili_red_merlot	6	2	1	4
Chili_red_cabernet	5	3	2	3
Chili_red_shiraz	7	1	2	6
Canada_red_pinot	4	2	3	1
Canada_white_chardonnay	4	3	2	1
Canada_white_sauvignon	8	4	3	2
USA_rose_cabernet	8	3	3	3
USA_rose_pinot	6	1	1	2
USA_rose_syrah	9	3	2	5

# PLS Objective Measures


	Acidity	Alcohol	Sugar	Tanin
Chili_red_merlot	5.33	13.8	2.75	559
Chili_red_cabernet	5.14	13.9	2.41	672
Chili_red_shiraz	5.16	14.3	2.20	455
Canada_red_pinot	5.70	13.3	1.70	320
Canada_white_chardonnay	6.00	13.5	3.00	35
Canada_white_sauvignon	7.50	12.0	3.50	40
USA_rose_cabernet	5.71	12.5	4.30	93
USA_rose_pinot	5.40	13.0	3.10	79
USA_rose_syrah	6.50	13.5	3.00	89

## PLS - Wine

```
pls.res <- tepPLS(  
  DATA1 = wine.objective,  
  center1 = T, scale1 = T,  
  DATA2 = wine.subjective,  
  center2 = T, scale2 = T,  
  graphs=F  
)
```



# PLS - Wine



# PLS - Wine

While PLS is similar to PCA we should focus on what PLS maximizes:

# PLS - Wine

While PLS is similar to PCA we should focus on what PLS maximizes:

$$\mathbf{L}_X^T \mathbf{L}_Y$$

# PLS - Wine


While PLS is similar to PCA we should focus on what PLS maximizes:

$$\mathbf{L}_X^T \mathbf{L}_Y$$

So we'll start with the latent variable scores


# PLS - Wine

PLS Wine Latent Variable Scores: LV1




# PLS - Wine


PLS Wine Latent Variable Scores: LV1



# PLS - Wine




# PLS - Wine








# PLS - Wine



# PLS - Wine



PLS Wine Latent Variable Scores: LV1



# Overview

## We're experts now

- PCA is the core of PLS

## We're experts now

- PCA is the core of PLS
- PLS generalizes PCA

## We're experts now

- PCA is the core of PLS
- PLS generalizes PCA
  - If your two matrices are both the same, e.g.,  $\mathbf{X}$

## We're experts now

- PCA is the core of PLS
- PLS generalizes PCA
  - If your two matrices are both the same, e.g.,  $\mathbf{X}$
  - PLS gives same results as PCA

## We're experts now

- Visualize first



## We're experts now

- Visualize first
- Use what you know to help construct the story from the numbers

## We're experts now

- Visualize first
- Use what you know to help construct the story from the numbers
  - We'll see some additional helpers in next part today

Break!

## We covered a lot!

- And we're not even at the good part yet!

## We covered a lot!

- And we're not even at the good part yet!
- $7 \pm 2$  break

## We covered a lot!

- And we're not even at the good part yet!
- $7 \pm 2$  break
  - Stretch, coffee

## We covered a lot!

- And we're not even at the good part yet!
- $7 \pm 2$  break
  - Stretch, coffee
  - Pressing questions