# Simple & Multiple Correspondence Analyses Contingency, categorical, ordinal, continuous and mixed data

Derek Beaton

Rotman Research Institute

October 28, 2019



#### Our new best friends





via @allison\_horst




via @allison\_horst



What do we do with all of these in a PCA like way?









- ▶ What do we do with all of these in a PCA like way?
- ► Some are *very* difficult and effectively ignored









- ▶ What do we do with all of these in a PCA like way?
- ► Some are *very* difficult and effectively ignored
  - ► We won't do that!

► Not everything is a number

- Not everything is a number
- ► Sometimes numbers aren't numbers!

- Not everything is a number
- Sometimes numbers aren't numbers!
- ▶ We need to recognize when this happens

- Not everything is a number
- Sometimes numbers aren't numbers!
- ▶ We need to recognize when this happens
  - And know what to do

### Typology

► SS Stevens (not a boat!)

### Typology

- ► SS Stevens (not a boat!)
- ► Levels of measurement

### Typology

- SS Stevens (not a boat!)
- Levels of measurement
- Excellent examples: https://en.wikipedia.org/wiki/Level\_of\_measurement

### Where to find everything

► Generally: https://github.com/derekbeaton/workshops

### Where to find everything

- ► Generally: https://github.com/derekbeaton/workshops
- ► Today:

► Revisit PCA

- ► Revisit PCA
- ► Looking at some data

- Revisit PCA
- ► Looking at some data
- ► Simple correspondence analysis

- Revisit PCA
- ► Looking at some data
- ► Simple correspondence analysis
  - and many of its connections

- Revisit PCA
- ► Looking at some data
- ► Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis

- Revisit PCA
- Looking at some data
- Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)

- Revisit PCA
- Looking at some data
- Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)
  - and how to handle various data types

- Revisit PCA
- Looking at some data
- ► Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)
  - and how to handle various data types
- A whole bunch of bonuses

- Revisit PCA
- Looking at some data
- Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)
  - and how to handle various data types
- A whole bunch of bonuses
  - Robustness

- Revisit PCA
- Looking at some data
- Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)
    - and how to handle various data types
- A whole bunch of bonuses
  - Robustness
  - PLS

- Revisit PCA
- Looking at some data
- Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)
  - and how to handle various data types
- A whole bunch of bonuses
  - Robustness
  - PLS
  - Networks

- Revisit PCA
- Looking at some data
- Simple correspondence analysis
  - and many of its connections
- Multiple correspondence analysis
  - generalizes CA (amongst many other things)
  - and how to handle various data types
- A whole bunch of bonuses
  - Robustness
  - PLS
  - Networks
  - Software

## Revisting PCA

▶ When we can compute a covariance or correlation matrix


- ▶ When we can compute a covariance or correlation matrix
- ► Break data into components

- ▶ When we can compute a covariance or correlation matrix
- ► Break data into components
  - Orthogonal


- ▶ When we can compute a covariance or correlation matrix
- Break data into components
  - Orthogonal
  - Rank ordered

- ▶ When we can compute a covariance or correlation matrix
- Break data into components
  - Orthogonal
  - Rank ordered
  - Made of bits & pieces of original measures


### Eigen- and singular value decompositions




### Eigen- and singular value decompositions




## Eigen- and singular value decompositions



## Eigen- and singular value decompositions



# Eigen- and singular value decompositions




Left (row) singular vectors




# Diagnosis and education


|     | CN | Dementia | MCI |
|-----|----|----------|-----|
| ADV | 39 | 7        | 54  |
| В   | 57 | 17       | 75  |
| B+  | 75 | 19       | 113 |
| HS  | 25 | 13       | 46  |
| HS+ | 39 | 9        | 77  |
|     |    |          |     |



| <b>•</b> | Given a table, and asked for a multivariate analysis |
|----------|--|
|          | We do what we know: PCA                              |



\*\*\*



# What did we analyze?


|          | CN    | Dementia | MCI   |
|----------|-------|----------|-------|
| CN       | 1.000 | 0.730    | 0.921 |
| Dementia | 0.730 | 1.000    | 0.652 |
| MCI      | 0.921 | 0.652    | 1.000 |

## What did PCA detect?

|     | CN | Dementia | MCI | Row sums |
|-----|----|----------|-----|----------|
| ADV | 39 | 7        | 54  | 100      |
| В   | 57 | 17       | 75  | 149      |
| B+  | 75 | 19       | 113 | 207      |
| HS  | 25 | 13       | 46  | 84       |
| HS+ | 39 | 9        | 77  | 125      |
|     |    |          |     |          |

# Let's try something different!

|          | ADV | В  | B+  | HS | HS+ |
|----------|-----|----|-----|----|-----|
| CN       | 39  | 57 | 75  | 25 | 39  |
| Dementia | 7   | 17 | 19  | 13 | 9   |
| MCI      | 54  | 75 | 113 | 46 | 77  |
|          |     |    |     |    |     |



Component 1. Explained variance: 97.8%

# What did PCA analyze?

|     | ADV   | В     | B+    | HS    | HS+   |
|-----|-------|-------|-------|-------|-------|
| ADV | 1.000 | 1.000 | 0.995 | 0.935 | 0.963 |
| В   | 1.000 | 1.000 | 0.994 | 0.932 | 0.960 |
| B+  | 0.995 | 0.994 | 1.000 | 0.965 | 0.984 |
| HS  | 0.935 | 0.932 | 0.965 | 1.000 | 0.996 |
| HS+ | 0.963 | 0.960 | 0.984 | 0.996 | 1.000 |

## What did PCA detect?

|          | ADV | В  | В+  | HS | HS+ | Row sums  |
|----------|-----|----|-----|----|-----|-----------|
| CN       | 39  | 57 | 75  | 25 | 39  | 235       |
| Dementia | 7   | 17 | 19  | 13 | 9   | <i>65</i> |
| MCI      | 54  | 75 | 113 | 46 | 77  | 365       |

#### What is PCA for?

► When we can compute a *meaningful* covariance or correlation matrix

### Let's take another look

|                | CN  | Dementia | MCI | Row sums |
|----------------|-----|----------|-----|----------|
| ADV            | 39  | 7        | 54  | 100      |
| В              | 57  | 17       | 75  | 149      |
| B+             | 75  | 19       | 113 | 207      |
| HS             | 25  | 13       | 46  | 84       |
| $\mathit{HS}+$ | 39  | 9        | 77  | 125      |
| Column sums    | 235 | 65       | 365 |          |

► Tell me things about this matrix

#### Let's take another look

|                | CN  | Dementia | MCI | Row sums |
|----------------|-----|----------|-----|----------|
| ADV            | 39  | 7        | 54  | 100      |
| В              | 57  | 17       | 75  | 149      |
| B+             | 75  | 19       | 113 | 207      |
| HS             | 25  | 13       | 46  | 84       |
| $\mathit{HS}+$ | 39  | 9        | 77  | 125      |
| Column sums    | 235 | 65       | 365 |          |

- ► Tell me things about this matrix
- ▶ What kind of problem does this look like?

# Simple correspondence analysis

▶ Initially: visualize contingency tables (a la PCA, factor analyses)

- ▶ Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature

- ▶ Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature
  - From Benzecri (1964) & Escofier (1965)

- Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature
  - From Benzecri (1964) & Escofier (1965)
- ► Fully developed by Escofier (1969)

- Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature
  - From Benzecri (1964) & Escofier (1965)
- ► Fully developed by Escofier (1969)
- Explosion of the technique in France

- Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature
  - From Benzecri (1964) & Escofier (1965)
- ► Fully developed by Escofier (1969)
- Explosion of the technique in France
  - Across virtually every field (except psychology and neuroscience)

- Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature
  - From Benzecri (1964) & Escofier (1965)
- ► Fully developed by Escofier (1969)
- Explosion of the technique in France
  - Across virtually every field (except psychology and neuroscience)
- ▶ The magic of CA relies on the magic of  $\chi^2$

- Initially: visualize contingency tables (a la PCA, factor analyses)
  - ► Text (corpus) of philosphy, biblical passages, literature
  - ► From Benzecri (1964) & Escofier (1965)
- ► Fully developed by Escofier (1969)
- Explosion of the technique in France
  - Across virtually every field (except psychology and neuroscience)
- ▶ The magic of CA relies on the magic of  $\chi^2$ 
  - And there's some crazy magic here

► Hotelling (1933) & Thurstone (1933)

- ► Hotelling (1933) & Thurstone (1933)
- ► Hirschfeld (1935) & Horst (1935)

- ► Hotelling (1933) & Thurstone (1933)
- ► Hirschfeld (1935) & Horst (1935)
- ► Guttman (1941)

- ► Hotelling (1933) & Thurstone (1933)
- ► Hirschfeld (1935) & Horst (1935)
- ► Guttman (1941)
- ▶ Burt (1950)

- ► Hotelling (1933) & Thurstone (1933)
- ► Hirschfeld (1935) & Horst (1935)
- ► Guttman (1941)
- ▶ Burt (1950)
- ► And then Benzecri (1964) & Escofier (1965)

- ► Hotelling (1933) & Thurstone (1933)
- Hirschfeld (1935) & Horst (1935)
- ► Guttman (1941)
- ▶ Burt (1950)
- ► And then Benzecri (1964) & Escofier (1965)
- Many more very important characters to re-discover CA

➤ See Lebart's History & Prehistory of CA

► See Lebart's History & Prehistory of CA

http://www.dtmvic.com/doc/About\_the\_History\_of\_CA.pdf


- ► See Lebart's History & Prehistory of CA http://www.dtmvic.com/doc/About\_the\_History\_of\_CA.pdf
- And Beh & Lombardo's series.


- ► See Lebart's History & Prehistory of CA
  - http://www.dtmvic.com/doc/About\_the\_History\_of\_CA.pdf
- ► And Beh & Lombardo's series
- A geneaology of CA:
  https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-842X.2012.00676.x


- See Lebart's History & Prehistory of CA
- http://www.dtmvic.com/doc/About\_the\_History\_of\_CA.pdf
- - https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-842X.2012.00676.x
  - A geneaology of CA 2: http://sibaese.unisalento.it/index.php/ejasa/article/view/19785


# We're diving in

|     | CN | Dementia | MCI |
|-----|----|----------|-----|
| ADV | 39 | 7        | 54  |
| В   | 57 | 17       | 75  |
| B+  | 75 | 19       | 113 |
| HS  | 25 | 13       | 46  |
| HS+ | 39 | 9        | 77  |









|     | CN | Dementia | MCI |
|-----|----|----------|-----|
| ADV | 39 | 7        | 54  |
| В   | 57 | 17       | 75  |
| B+  | 75 | 19       | 113 |
| HS  | 25 | 13       | 46  |

*HS*+ 39

CN


What if we perform CA on this?

**ADV** 

В B+

HS

HS+



# How did that happen?

Table 1: Data

|     | CN | Dementia | MCI |
|-----|----|----------|-----|
| ADV | 39 | 7        | 54  |
| В   | 57 | 17       | 75  |
| B+  | 75 | 19       | 113 |
| HS  | 25 | 13       | 46  |
| HS+ | 39 | 9        | 77  |

Table 2: Observed probabilites

|     | CN    | Dementia | MCI   |
|-----|-------|----------|-------|
| ADV | 0.059 | 0.011    | 0.081 |
| В   | 0.086 | 0.026    | 0.113 |
| B+  | 0.113 | 0.029    | 0.170 |
| HS  | 0.038 | 0.020    | 0.069 |

0.014

0.116

HS+

0.059

Table 3: Observed probabilites and margins

|                | CN    | Dementia | MCI   | Row sums |
|----------------|-------|----------|-------|----------|
| ADV            | 0.059 | 0.011    | 0.081 | 0.150    |
| В              | 0.086 | 0.026    | 0.113 | 0.224    |
| B+             | 0.113 | 0.029    | 0.170 | 0.311    |
| HS             | 0.038 | 0.020    | 0.069 | 0.126    |
| $\mathit{HS}+$ | 0.059 | 0.014    | 0.116 | 0.188    |
| Column sums    | 0.353 | 0.098    | 0.549 |          |

Table 4: Expected probabilites and margins

|                | CN    | Dementia | MCI   | Row sums |
|----------------|-------|----------|-------|----------|
| ADV            | 0.053 | 0.015    | 0.083 | 0.150    |
| В              | 0.079 | 0.022    | 0.123 | 0.224    |
| B+             | 0.110 | 0.030    | 0.171 | 0.311    |
| HS             | 0.045 | 0.012    | 0.069 | 0.126    |
| $\mathit{HS}+$ | 0.066 | 0.018    | 0.103 | 0.188    |
| Column sums    | 0.353 | 0.098    | 0.549 |          |

Table 5: Deviations: Observed - Expected

|     | CN    | Dementia | MCI    |
|-----|-------|----------|--------|
| ADV | 0.006 | -0.004   | -0.001 |
| В   | 0.007 | 0.004    | -0.010 |
| B+  | 0.003 | -0.002   | -0.001 |

0.007 0.000

-0.005 0.013

HS -0.007

*HS*+ -0.008

Table 6: Row constraints (inverse row margins)

|     | ADV  | В     | В+    | HS    | HS+  |
|-----|------|-------|-------|-------|------|
| ADV | 6 65 | 0.000 | 0.000 | 0.000 | 0.00 |

|     | ADV  | В     | B+    | HS    | HS+  |
|-----|------|-------|-------|-------|------|
| ADV | 6.65 | 0.000 | 0.000 | 0.000 | 0.00 |
| В   | 0.00 | 4.463 | 0.000 | 0.000 | 0.00 |

3.213

0.000

0.000

0.000

7.917

0.000

0.00

0.00

5.32

0.000

0.000

0.000

B+

HS

HS+

0.00


0.00

0.00


Table 7: Column constraints (inverse column margins)

|          | CN   | Dementia | MCI   |
|----------|------|----------|-------|
| CN       | 2.83 | 0.000    | 0.000 |
| Dementia | 0.00 | 10.231   | 0.000 |
| MCI      | 0.00 | 0.000    | 1.822 |


#### What CA needs



**R:** Row constraints (inverse row probabilities)



Z: Deviations



**C:** Column constraints (inverse column probabilities)

## ► GSVD(**R**, **X**, **C**)

- **▶** GSVD(**R**, **X**, **C**)
- ▶ Required for CA and fancier PCA-like techniques & extensions

- **▶** GSVD(**R**, **X**, **C**)
- ▶ Required for CA and fancier PCA-like techniques & extensions
- Uses but generalizes the SVD

- **▶** GSVD(**R**, **X**, **C**)
- ▶ Required for CA and fancier PCA-like techniques & extensions
- Uses but generalizes the SVD
- Uses row & column weights (constraints)

- **▶** GSVD(**R**, **X**, **C**)
- ▶ Required for CA and fancier PCA-like techniques & extensions
- Uses but generalizes the SVDUses row & column weights (constraints)
- Gives back

- ► GSVD(R, X, C)
- ▶ Required for CA and fancier PCA-like techniques & extensions
- Uses but generalizes the SVDUses row & column weights (constraints)
- Gives back
  - Component (factor) scores


- ▶ GSVD(R, X, C)
- ▶ Required for CA and fancier PCA-like techniques & extensions
- Uses but generalizes the SVDUses row & column weights (constraints)
- Gives back
  - Component (factor) scores
  - Component (factor) scores
  - ► Eigenvalues, singular values, & singular vectors

- ▶ GSVD(R, X, C)
- Required for CA and fancier PCA-like techniques & extensions
  - Uses but generalizes the SVD Uses row & column weights (constraints)
- Gives back
  - Component (factor) scores


  - Eigenvalues, singular values, & singular vectors
  - Generalized singular vectors

## What we really decompose

- A rectangle
- · Deviations: Observed Expected
  - Expected from Observed's margins



- Two squares
- Row margins and column margins





# $\frac{\left(\textbf{O}-\textbf{E}\right)}{\textbf{E}^{\frac{1}{2}}}$

$$\chi^2 = \sum \frac{(\mathbf{O} - \mathbf{E})^2}{\mathbf{E}}$$

#### CA's first secrets

▶ Orthogonal slices of  $\chi^2$ 

Maybe here is worth presenting the code and some output?

#### CA's first secrets

- ▶ Orthogonal slices of  $\chi^2$
- Sum of the eigenvalues  $\times$  sum of the table  $=\chi^2$

Maybe here is worth presenting the code and some output?

#### The GSVD

Simple quick magic Then visualize it (as 3 matrices, then 1 over 2 which is just the probs not inverse) Then swing back to Chi2 Then swing to CCA Then expand it & transition to MCA

[[[pick up here and drop most of the stuff below]]]

## The GSVD

► The generalized SVD

#### The GSVD

- ► The generalized SVD
  - ► Constraints (weights) for rows & columns of rectangular table

- ► The generalized SVD
  - ► Constraints (weights) for rows & columns of rectangular table
  - ▶ Required for CA and fancier PCA-like techniques & extensions

- ▶ The generalized SVD
  - Constraints (weights) for rows & columns of rectangular table
  - Required for CA and fancier PCA-like techniques & extensions
- ► For some matrix **X** with *I* rows and *J* columns

- The generalized SVD
  - Constraints (weights) for rows & columns of rectangular table
  - ▶ Required for CA and fancier PCA-like techniques & extensions
- For some matrix **X** with *I* rows and *J* columns
- $\triangleright$  SVD(**X**) vs. GSVD(**W**<sub>I</sub>,**X**,**W**<sub>J</sub>)

- The generalized SVD
  - Constraints (weights) for rows & columns of rectangular table
  - ▶ Required for CA and fancier PCA-like techniques & extensions
- For some matrix **X** with *I* rows and *J* columns
- $\triangleright$  SVD(**X**) vs. GSVD(**W**<sub>I</sub>,**X**,**W**<sub>J</sub>)
- ► GSVD is

- The generalized SVD
  - Constraints (weights) for rows & columns of rectangular table
  - ▶ Required for CA and fancier PCA-like techniques & extensions
- For some matrix **X** with *I* rows and *J* columns
- $\triangleright$  SVD(**X**) vs. GSVD(**W**<sub>I</sub>,**X**,**W**<sub>J</sub>)
- ► GSVD is
  - Matrix multiplication (by constraints on data)

- The generalized SVD
  - Constraints (weights) for rows & columns of rectangular table
  - ▶ Required for CA and fancier PCA-like techniques & extensions
- ▶ For some matrix **X** with *I* rows and *J* columns
- $\triangleright$  SVD(**X**) vs. GSVD(**W**<sub>I</sub>,**X**,**W**<sub>J</sub>)
- ► GSVD is
  - Matrix multiplication (by constraints on data)
  - ► The SVD

- The generalized SVD
  - Constraints (weights) for rows & columns of rectangular table
  - ▶ Required for CA and fancier PCA-like techniques & extensions
- For some matrix **X** with *I* rows and *J* columns
- $\triangleright$  SVD(**X**) vs. GSVD(**W**<sub>I</sub>,**X**,**W**<sub>J</sub>)
- ► GSVD is
  - Matrix multiplication (by constraints on data)
  - ► The SVD
  - More matrix multiplication (by constraints on vectors)

▶ O, wi & wj, E, Z

- ► O, wi & wj, E, Z
- ▶ Oh look that's Chi2

- ► O, wi & wj, E, Z
- ► Oh look that's Chi2
- ► Sum of eigenvalues \* sum of table = Chi2.

- ► O, wi & wj, E, Z
- ► Oh look that's Chi2
- ► Sum of eigenvalues \* sum of table = Chi2.
  - Each component is an additive orthogonal slice of Chi2. WOAH.

- ▶ 0, wi & wj, E, Z
- ▶ Oh look that's Chi2
- ► Sum of eigenvalues \* sum of table = Chi2.
  - Each component is an additive orthogonal slice of Chi2. WOAH.
  - ► The eigenvalues are *magic*

## CA visualized

► Oh look it's CCA-ish

## CA visualized

- ► Oh look it's CCA-ish
- ▶ Oh it really really is CCA-ish!

► It's like PCA

- ► It's like PCA
  - ► Variance (singular values, eigen values)

- ► It's like PCA
  - ► Variance (singular values, eigen values)
  - ▶ Directions & inter-relationships

- ► It's like PCA
  - Variance (singular values, eigen values)
  - ► Directions & inter-relationships
  - Components scores

- ► It's like PCA
  - Variance (singular values, eigen values)
  - ► Directions & inter-relationships
  - Components scores
- ► It's unlike PCA

- It's like PCA
  - Variance (singular values, eigen values)
  - Directions & inter-relationships
  - Components scores
- ► It's unlike PCA
  - ► Relative interpretation *between* sets

- ► It's like PCA
  - Variance (singular values, eigen values)
  - Directions & inter-relationships
  - Components scores
- ► It's unlike PCA
  - ► Relative interpretation *between* sets
  - Generalized singular vectors

- ► It's like PCA
  - Variance (singular values, eigen values)
  - Directions & inter-relationships
  - Components scores
- ► It's unlike PCA
  - ▶ Relative interpretation *between* sets
  - Generalized singular vectors
  - Bifactor


- ► It's like PCA
  - Variance (singular values, eigen values)
  - Directions & inter-relationships
  - Components scores
- ► It's unlike PCA
  - ▶ Relative interpretation *between* sets
  - Generalized singular vectors
  - Bifactor
    - Rows & columns treated the same

- ► It's like PCA
  - Variance (singular values, eigen values)
  - Directions & inter-relationships
  - Components scores
- ► It's unlike PCA
  - ▶ Relative interpretation *between* sets
  - Generalized singular vectors
  - Bifactor
    - Rows & columns treated the same
    - ► Together they help make components, as opposed to PCA







# Chi-squared



# Chi-squared



See here



(Some) References

## See the reference sections of these

▶ Beaton, D., Saporta, G., Abdi, H., & Alzheimer's Disease Neuroimaging Initiative. (2019). A generalization of partial least squares regression and correspondence analysis for categorical and mixed data: An application with the ADNI data. bioRxiv, 598888.

### See the reference sections of these

- ▶ Beaton, D., Saporta, G., Abdi, H., & Alzheimer's Disease Neuroimaging Initiative. (2019). A generalization of partial least squares regression and correspondence analysis for categorical and mixed data: An application with the ADNI data. bioRxiv, 598888.
- Beaton, D., Sunderland, K. M., Levine, B., Mandzia, J., Masellis, M., Swartz, R. H., ... & Strother, S. C. (2019). Generalization of the minimum covariance determinant algorithm for categorical and mixed data types. bioRxiv, 333005.

### And these

Abdi, H., Guillemot, V., Eslami, A., & Beaton, D. (2017). Canonical correlation analysis. Encyclopedia of Social Network Analysis and Mining, 1-16.

#### And these

- Abdi, H., Guillemot, V., Eslami, A., & Beaton, D. (2017). Canonical correlation analysis. Encyclopedia of Social Network Analysis and Mining, 1-16.
- Beaton, D., Dunlop, J., & Abdi, H. (2016). Partial least squares correspondence analysis: A framework to simultaneously analyze behavioral and genetic data. Psychological methods, 21(4), 621.

► Greenacre, M. (2017). Correspondence analysis in practice. CRC press.

- Greenacre, M. (2017). Correspondence analysis in practice. CRC press.
- Greenacre, M. J. (1984). Theory and Applications of Correspondence Analysis. Retrieved from http://books.google.com/books?id=LsPaAAAAMAAJ

▶ Greenacre, M. J. (2010). Correspondence analysis. Wiley Interdisciplinary Reviews: Computational Statistics, 2(5), 613–619. https://doi.org/10.1002/wics.114

- ▶ Greenacre, M. J. (2010). Correspondence analysis. Wiley Interdisciplinary Reviews: Computational Statistics, 2(5), 613–619. https://doi.org/10.1002/wics.114
- Lebart, L., Morineau, A., & Warwick, K. M. (1984). Multivariate descriptive statistical analysis: correspondence analysis and related techniques for large matrices. Wiley.

- ▶ Greenacre, M. J. (2010). Correspondence analysis. Wiley Interdisciplinary Reviews: Computational Statistics, 2(5), 613–619. https://doi.org/10.1002/wics.114
- Lebart, L., Morineau, A., & Warwick, K. M. (1984). Multivariate descriptive statistical analysis: correspondence analysis and related techniques for large matrices. Wiley.
- Nguyen, L. H., & Holmes, S. (2019). Ten quick tips for effective dimensionality reduction. PLOS Computational Biology, 15(6), e1006907.

#### Data

► Escofier, B. (1978). Analyse factorielle et distances répondant au principe d'équivalence distributionnelle. Revue de Statistique Appliquée, 26(4), 29–37.

#### Data

- Escofier, B. (1978). Analyse factorielle et distances répondant au principe d'équivalence distributionnelle. Revue de Statistique Appliquée, 26(4), 29–37.
- Escofier, B. (1979). Traitement simultané de variables qualitatives et quantitatives en analyse factorielle. Cahiers de l'Analyse Des Données, 4(2), 137–146.

#### Data

- ► Escofier, B. (1978). Analyse factorielle et distances répondant au principe d'équivalence distributionnelle. Revue de Statistique Appliquée, 26(4), 29–37.
- Escofier, B. (1979). Traitement simultané de variables qualitatives et quantitatives en analyse factorielle. Cahiers de l'Analyse Des Données, 4(2), 137–146.
- Greenacre, M. (2014). Data Doubling and Fuzzy Coding. In J. Blasius & M. Greenacre (Eds.), Visualization and Verbalization of Data (pp. 239–253). Philadelphia, PA, USA: CRC Press.

# History

▶ Holmes S, Josse J. Discussion of "50 Years of Data Science". Journal of Computational and Graphical Statistics. 2017, V26(4) 768-769. https://www.tandfonline.com/doi/full/10.10 80/10618600.2017.1385471