Mesh Smoothing

Outline

- Differential Geometry
- **Discrete Differential Geometry**
- Mesh Quality Measures

Differential Geometry: Surfaces

- Normals
- First/Second Fundamental forms
- Normal curvature
- Principal curvatures
- mean/Gauss Curvatures

Gauss-Bonnet Theorem

- For **any** closed manifold surface with Euler characteristic $\chi = 2 - 2g$

$$\int_{\Omega \in S} K(u,v)\,du\,dv = 2\pi\chi$$

$\int K($ $) = \int K($ $) = \int K($ $) = 4\pi$

Gauss-Bonnet Theorem

- Sphere

$$\kappa_1 = \kappa_2 = 1/r$$

$$K = \kappa_1 \kappa_2 = 1/r^2$$

$$\int K = 4\pi r^2 \cdot \frac{1}{r^2} = 4\pi$$

– when sphere is deformed new positive and negative curvature cancel out!

Functions on Surfaces

scalar values on surfaces

vector values on surfaces

Differential Operators

用来measure
向量场的源头的汇的大小

- Gradient (vector)

$$\nabla f := \left(\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_n} \right)$$

$$\text{div } g := < \left(\frac{\partial}{\partial x_1}, \ldots \right), g > = \nabla \cdot g$$

  – points in the direction of steepest ascent

Differential Operators

$$\text{div } g := < \left(\frac{\partial}{\partial x_1}, \ldots \right), g > = \nabla \cdot g$$

div 运算

$$\text{div } \mathbf{G} := < \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right), (G_x, G_y, G_z) > = \frac{\partial G_x}{\partial x} + \frac{\partial G_y}{\partial y} + \frac{\partial G_z}{\partial z}$$

e.g. 速度在运动中心分布

指定某点速度 附近速度 是大还是小 流出量

Differential Operators

- Divergence

$$\mathrm{div} F = \nabla \cdot F := \frac{\partial F_1}{\partial x_1} + \ldots + \frac{\partial F_n}{\partial x_n}$$

  - volume density of outward flux of vector field
  - magnitude of source or sink at given point
  - Example: Incompressible fluid
    - velocity field is divergence-free

Differential Operators

- Divergence

$$\mathrm{div} F = \nabla \cdot F := \frac{\partial F_1}{\partial x_1} + \ldots + \frac{\partial F_n}{\partial x_n}$$

high divergence

low divergence

div ⟺ ∇·F ⟺ sum

sum

∇F ≠ vector

∇F ≠ ∇·F

sum

Laplace Operator

$$\text{div}\ \nabla f$$

vector：
所有分量的一阶导

$$\text{div}\ \nabla f = \sum_i \frac{\partial^2 f}{\partial x_i^2}$$

$$\text{div}\ \nabla f = \sum_i \frac{\partial^2 f}{\partial x_i^2} = \Delta f$$

scaler

Laplace Operator

Laplace operator

gradient operator

2nd partial derivatives

$$\Delta f = \mathrm{div} \nabla f = \sum_i \frac{\partial^2 f}{\partial x_i^2}$$

function in Euclidean space

eg. 颜色、温度

divergence operator

Cartesian coordinates

Laplace-Beltrami Operator

coord

Extension of Laplace to **functions on manifolds**

$$p = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

$$\Delta p = -2|H|\vec{n}$$

normal

scalar

vector

mean curvature

Laplace-Beltrami

gradient operator

$n \times n$

$$\Delta_{\mathcal{S}} f = \mathrm{div}_{\mathcal{S}}\, \nabla_{\mathcal{S}} f$$

function on manifold $\mathcal{S}$

divergence operator

Laplace-Beltrami Operator

Extension of Laplace to **functions on manifolds**

Laplace-Beltrami

gradient operator

mean curvature [scalar]

$$\Delta_{\mathcal{S}} \mathbf{x} = \mathrm{div}_{\mathcal{S}} \nabla_{\mathcal{S}} \mathbf{x} = -2H\mathbf{n}$$

n x n

n x 3

x, y, z

n x 3

coordinate function

divergence operator

surface normal

$2\pi - \Sigma \theta_i$

Outline

- Differential Geometry
- **Discrete Differential Geometry**
- Mesh Quality Measures

Discrete Curvatures

- How to discretize curvatures on a mesh?
  - Zero curvature within triangles
  - Infinite curvature at edges / vertices
  - Point-wise definition does not make sense

- Approximate differential properties at point $\mathbf{v}$ as average over local neighborhood $A(\mathbf{v})$
  - $\mathbf{v}$ is a mesh vertex
  - $A(\mathbf{v})$ within one-ring neighborhood

$$K(v) \approx \frac{1}{A(v)} \int_{A(v)} K(\mathbf{x}) \, \mathrm{d}A$$

Area Discretizations

Voronoi Cells

将空间划分为多个region

3d

正方

划分 找出region

Discrete Curvatures

- Which curvatures to discretize?
  - Discretize Laplace-Beltrami operator
  - Laplace-Beltrami gives us mean curvature H
  - Discretize Gaussian curvature K

Laplace Operator on Meshes?

- Extend finite differences to meshes?
  - What weights per vertex / edge?

1D grid 2D grid 2D/3D mesh

Matrix Version of Neighborhood

2D grid

Uniform Laplacian Matrix

neighbor 均匀 & 自己 权重

- Uniform discretization

neighbor

$$\Delta_{\text{uni}} f(v_i) := \frac{1}{|\mathcal{N}_1(v_i)|} \sum_{v_j \in \mathcal{N}_1(v_i)} \left(f(v_j) - f(v_i)\right)$$

随机平滑 过程

$$= \left(\frac{1}{|\ |} \sum_{j} f(v_j)\right) - f(v_i)$$

- Properties
  - depends only on connectivity
  - simple and efficient
  - bad approximation for irregular triangulations
    - can give non-zero *H* for planar meshes
    - tangential drift for mesh smoothing

$$= \begin{bmatrix} \cdots \\ \cdots \end{bmatrix} \begin{bmatrix} f(v_1) \\ \vdots \\ f(v_n) \end{bmatrix}$$

$n \times n$ $\quad n \times 1$

$\Delta u \rightarrow$

Uniform Laplacian Matrix

- Uniform discretization

这是平均

这是理想的，想要这个

$$\Delta_{\text{uni}}\mathbf{x}_i := \frac{1}{|\mathcal{N}_1(v_i)|}\sum_{v_j\in\mathcal{N}_1(v_i)}(\mathbf{x}_j-\mathbf{x}_i) \approx -2H\mathbf{n}$$

- Properties
  - depends only on connectivity
  - simple and efficient
  - bad approximation for irregular triangulations
    - can give non-zero H for planar meshes
    - tangential drift for mesh smoothing

Discrete Laplace-Beltrami

- Cotangent discretization not symmetric

$$\Delta_{\mathcal{S}} f(v_i) := \frac{1}{2A(v_i)} \sum_{v_j \in \mathcal{N}_1(v_i)} (\cot \alpha_{ij} + \cot \beta_{ij}) \left(f(v_j) - f(v_i) \right)$$

symmetric

to solve this

M^{-1}

$$\Delta_S \mathbf{x} = -2H\mathbf{n}$$

$= \begin{bmatrix} 2A_1 & & \\ & 2A_2 & \\ & & \ddots \end{bmatrix}$

#1: Barycentric Cells

- Connect edge midpoints and triangle barycenters
  - Simple to compute
  - Area is 1/3 of triangle areas

#2: Voronoi Cells

- Connect edge midpoints and
  - Circumcenters for all triangles
  - What about obtuse triangles?

Salve

#3: Mixed Cells

- Connect edge midpoints and
  - Circumcenters for non-obtuse triangles
  - Midpoint of opposite edge for obtuse triangles
  - Better approximation, more complex to compute...

Discrete Laplace-Beltrami

- Cotangent discretization

$$\Delta_{\mathcal{S}} f(v) := \frac{1}{2A(v)} \sum_{v_i \in \mathcal{N}_1(v)} (\cot \alpha_i + \cot \beta_i)\,(f(v_i) - f(v))$$

- Problems
  - weights can become negative (when?)
  - depends on triangulation
- Still the most widely used discretization

Discrete Curvatures

- Mean curvature (absolute value)

$$H = \frac{1}{2} \|\Delta_{\mathcal{S}} \mathbf{x}\|$$

$$\Delta_S \mathbf{x} = -2H\mathbf{n}$$

- Gaussian curvature

$$K = (2\pi - \sum_j \theta_j)/A$$

Literature

- Taubin: *A signal processing approach to fair surface design*, SIGGRAPH 1996.
- Desbrun et al.: *Implicit Fairing of Irregular Meshes using Diffusion and Curvature Flow*, SIGGRAPH 1999.
- Meyer et al.: *Discrete Differential-Geometry Operators for Triangulated 2-Manifolds*, VisMath 2002.
- Wardetzky, Mathur, Kaelberer, Grinspun: Discrete Laplace Operators: No free lunch, SGP 2007

Outline

- Differential Geometry
- Discrete Differential Geometry
- **Mesh Quality Measures**

Mesh Quality

- Visual inspection of “sensitive” attributes
  - Specular shading

Mesh Quality

- Visual inspection of “sensitive” attributes
  - Specular shading
  - Reflection lines

Mesh Quality

- Visual inspection of “sensitive” attributes
  - Specular shading
  - Reflection lines
    - differentiability one order lower than surface
    - can be efficiently computed using graphics hardware

C^0 C^1 C^2

Mesh Quality

- Visual inspection of “sensitive” attributes
  - Specular shading
  - Reflection lines
  - Curvature
    - Mean curvature

Mesh Quality

- Visual inspection of “sensitive” attributes
  - Specular shading
  - Reflection lines
  - Curvature
    - Mean curvature
    - Gauss curvature

Mesh Quality Criteria

- Smoothness
  - Low geometric noise

smoother

[grall]

Mesh Quality Criteria

- Smoothness
  - Low geometric noise
- Fairness
  - Simplest shape

Mesh Quality Criteria

- Smoothness
  - Low geometric noise
- Fairness
  - Simplest shape
- Adaptive tessellation
  - Low complexity

Mesh Quality Criteria

- Smoothness
  - Low geometric noise
- Fairness
  - Simplest shape
- Adaptive tessellation
  - Low complexity
- Triangle shape
  - Numerical robustness

Mesh Optimization

- Smoothness
  - ➡ Smoothing
- Fairness
  - ➡ Fairing
- Adaptive tessellation
  - ➡ Decimation
- Triangle shape
  - ➡ Remeshing

#1: Filter Scanning Noise

- Filter out high frequency noise

Desbrun, Meyer, Schroeder, Barr
Implicit Fairing of Irregular Meshes using Diffusion and Curvature Flow
SIGGRAPH 99

Mesh Smoothing *(reduce noise)*

#2: Feature Modification

Original | Low-Pass | Exaggerate

Kim, Rosignac: *Geofilter: Geometric Selection of Mesh Filter Parameters*, Eurographics 05

#3: Fair Surface Design

- Fair Surface Design

Schneider, Kobbelt: *Geometric fairing of irregular meshes for free-form surface design*, CAGD 18(4), 2001

#4: Hole Filling *(energy minimization)*

#5: Mesh Parameterization

$$\mathbf{x}(u, v) \qquad (u, v)$$

Levy et al.: Least squares conformal maps for automatic texture atlas generation, SIGGRAPH 2002.

#6: Surface Deformation

Olga Sorkine and Marc Alexa. As-rigid-as-possible surface modeling, SGP 2007

Botsch, Kobbelt: *An intuitive framework for real-time freeform modeling*, SIGGRAPH 04

Laplace Operator on Surfaces Δ

- Uniform Laplace operator
  - depends on mesh connectivity only
  - approximates second derivative on surfaces

- Cotan operator (Laplace Beltrami)

not symmetric symmetric

$$\Delta = M^{-1} C$$

$$M = M^T \qquad C = C^T$$

$$\Delta^T \neq \Delta$$

$$c_{ij} = \frac{1}{2}\left(\cot(\alpha_{ij}) + \cot(\beta_{ij})\right)$$

$$c_{ii} = -\sum_{j \in N_1(i)} c_{ij} \qquad \sum_j c_{ij} = 0$$

along each row, sum 0 if up

Common Problem Types

- Laplace equation (special case of Poisson equation)

$$\Delta f = 0$$

$$M^{-1}Cf = 0 \rightarrow Cf = 0 \quad \text{subject to boundary conditions}$$

$$f(x) = f_0 \quad x \in \partial B$$

Common Problem Types

- Laplace equation (special case of Poisson equation)

$$\Delta f = 0$$

$$M^{-1}Cf = 0 \rightarrow Cf = 0 \quad \text{subject to boundary conditions}$$

$$f(x) = f_0 \quad x \in \partial B$$

- Poisson equation

$$\Delta f = g$$

$$M^{-1}Cf = g \rightarrow Cf = Mg$$

Common Problem Types

- Laplace equation (special case of Poisson equation)

$$\Delta f = 0$$

$$M^{-1}Cf = 0 \rightarrow Cf = 0 \quad \text{subject to boundary conditions}$$

$$f(x) = f_0 \quad x \in \partial B$$

- Poisson equation

$$\Delta f = g$$

$$M^{-1}Cf = g \rightarrow Cf = Mg$$

- Differential equation

$$f_t = \frac{\partial f}{\partial t} = \Delta f$$

Common Problem Types

- Laplace equation (special case of Poisson equation)

$$\Delta f = 0$$

$$M^{-1}Cf = 0 \rightarrow Cf = 0 \quad \text{subject to boundary conditions}$$

$$f(x) = f_0 \quad x \in \partial B$$

not symmetric

- Poisson equation

$$\Delta f = g$$

$$M^{-1}Cf = g \rightarrow Cf = Mg$$

$\Delta x = \lambda x$

$M^{-1}Cx = \lambda x$

$M^{\frac{1}{2}}M^{-\frac{1}{2}}M^{-\frac{1}{2}}Cx =$

$M^{-\frac{1}{2}}Cx = \lambda M^{\frac{1}{2}}x$

$\vdots$

$y = M^{\frac{1}{2}}x$

$x = M^{-\frac{1}{2}}y$

- Differential equation

$$f_t = \frac{\partial f}{\partial t} = \Delta f$$

- Eigen/spectral analysis

$$\Delta \phi_i = \lambda_i \phi_i$$

$$\left(M^{-\frac{1}{2}}CM^{-\frac{1}{2}}\right) M^{\frac{1}{2}}\phi_i = \lambda_i M^{\frac{1}{2}}\phi_i$$

$$\left(M^{-\frac{1}{2}}CM^{-\frac{1}{2}}\right) \xi_i = \lambda_i \xi_i \qquad \rightarrow \phi_i = M^{-\frac{1}{2}}\xi_i$$

Outline

- **Spectral Analysis**
- Diffusion Flow
- Energy Minimization

Fourier Transform

- Represent a function as a weighted sum of sines and cosines

Joseph Fourier 1768 - 1830

$$f(x) = a_0 + a_1 \cos(x)$$

Fourier Transform

- Represent a function as a weighted sum of sines and cosines

Joseph Fourier 1768 - 1830

$$f(x) = a_0 + a_1 \cos(x) + a_2 \cos(3x) + a_3 \cos(5x) + a_4 \cos(7x) + \ldots$$

Fourier Transform

$$F(\omega) = \int_{-\infty}^{\infty} f(x)\, \mathrm{e}^{-2\pi \mathrm{i} \omega x} \mathrm{d}x$$

$$f(x) = \int_{-\infty}^{\infty} F(\omega)\, \mathrm{e}^{2\pi \mathrm{i} \omega x}\, \mathrm{d}\omega$$

Convolution

- Smooth signal by convolution with a kernel function

$$h(x) \;=\; f * g \;:=\; \int f(y) \cdot g(x-y)\,\mathrm{d}y$$

- Example: Gaussian blurring

Convolution

- Smooth signal by convolution with a kernel function

$$h(x) \;=\; f * g \;:=\; \int f(y) \cdot g(x-y)\,\mathrm{d}y$$

- Convolution in spatial domain ⇔

  Multiplication in frequency domain

$$H(\omega) \;=\; F(\omega) \cdot G(\omega)$$

Fourier Analysis

- Low-pass filter discards high frequencies

spatial domain

frequency domain

LOW PASS

Fourier Transform

- Spatial domain *f(x)* → Frequency domain *F(ω)*

$$F(\omega) = \int_{-\infty}^{\infty} f(x)\, \mathrm{e}^{-2\pi \mathrm{i} \omega x} \mathrm{d}x$$

- Multiply by low-pass filter *G(ω)*

$$F(\omega) \;\leftarrow\; F(\omega) \cdot G(\omega)$$

- Frequency domain *F(ω)* → Spatial domain *f(x)*

$$f(x) = \int_{-\infty}^{\infty} F(\omega)\, \mathrm{e}^{2\pi \mathrm{i} \omega x}\, \mathrm{d}\omega$$

Fourier Transform

- Consider L^2 function space with inner product

$$\langle f, g \rangle := \int_{-\infty}^{\infty} f(x)\, \overline{g(x)}\, \mathrm{d}x$$

- Complex "waves" build an orthonormal basis

$$e_\omega(x) \;:=\; \mathrm{e}^{-2\pi \mathrm{i} \omega x} \;=\; \cos(2\pi\omega x) - \mathrm{i}\sin(2\pi\omega x)$$

- Fourier transform is a change of basis

$$f(x) = \int_{-\infty}^{\infty} \langle f, e_\omega \rangle\; e_\omega\; \mathrm{d}\omega \qquad\qquad f(x) = \sum_{\omega=-\infty}^{\infty} \langle f, e_\omega \rangle\; e_\omega\; \mathrm{d}\omega$$

Fourier Analysis on Meshes?

- Only applicable to parametric patches $f(u, v)$
- Generalize frequency concept to meshes!
- Complex waves are Eigenfunctions of Laplace

$$\Delta\left(e^{2\pi \mathrm{i}\omega x}\right) = \frac{\mathrm{d}^2}{\mathrm{d}x^2} e^{2\pi \mathrm{i}\omega x} = -(2\pi\omega)^2 e^{2\pi \mathrm{i}\omega x}$$

➡ ***Use Eigenfunctions of discrete Laplace-Beltrami***

Discrete Laplace-Beltrami

- Function values sampled at mesh vertices

$$\mathbf{f} = [f_1, f_2, \ldots, f_n] \in \mathbb{R}^n$$

- Discrete Laplace-Beltrami (per vertex)

$$\Delta_{\mathcal{S}} f(v_i) := \frac{1}{2A_i} \sum_{v_j \in \mathcal{N}_1(v_i)} (\cot\alpha_{ij} + \cot\beta_{ij}) \left(f(v_j) - f(v_i)\right)$$

Discrete Laplace-Beltrami

- Discrete Laplace *operator* (per mesh)
  - Sparse matrix

$$\mathbf{L} = \mathbf{M}^{-1}\mathbf{C} \in \mathbb{R}^{n \times n}$$

$$\begin{pmatrix} \vdots \\ \Delta_{\mathcal{S}} f(v_i) \\ \vdots \end{pmatrix} = \mathbf{L} \cdot \begin{pmatrix} \vdots \\ f(v_i) \\ \vdots \end{pmatrix}$$

Discrete Laplace-Beltrami

- Discrete Laplace *operator* (per mesh)

$$\mathbf{L} = \mathbf{M}^{-1}\mathbf{C} \in \mathbb{R}^{n \times n}$$

  – Sparse matrix

$$\mathbf{C}_{ij} = \begin{cases} \cot\alpha_{ij} + \cot\beta_{ij}, & i \neq j\,,\ j \in \mathcal{N}_1(v_i) \\ -\sum_{v_j \in \mathcal{N}_1(v_i)} (\cot\alpha_{ij} + \cot\beta_{ij}) & i = j \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{M}^{-1} = \operatorname{diag}\left(\ldots, \frac{1}{2A_i}, \ldots\right)$$

  – Note fraction 1/2 can either be in C or in M matrix

Discrete Laplace-Beltrami

- Discrete function sampled at mesh vertices

$$\mathbf{f} = [f_1, f_2, \ldots, f_n] \in \mathbb{R}^n$$

- Discrete Laplace-Beltrami (per vertex)

$$\Delta_{\mathcal{S}} f(v_i) := \frac{1}{2A_i} \sum_{v_j \in \mathcal{N}_1(v_i)} (\cot\alpha_{ij} + \cot\beta_{ij}) \left(f(v_j) - f(v_i)\right)$$

- Discrete Laplace-Beltrami matrix **L**
  - Eigenvectors are “natural vibrations”
  - Eigenvalues are “natural frequencies”

$$\Delta \mathbf{e}_i = \lambda_i \mathbf{e}_i$$

Discrete Laplace-Beltrami

$\mathbf{e}_2$ $\mathbf{e}_3$

- Discrete Laplace-Beltrami matrix **L**
  - Eigenvectors are “natural vibrations”
  - Eigenvalues are “natural frequencies”

Spectral Analysis

1. Setup Laplace-Beltrami matrix $\mathbf{L} = \Delta$
2. Compute *k* smallest eigenvectors {$\mathbf{e}_1$,..., $\mathbf{e}_k$}
3. Reconstruct mesh from those (component-wise)

$$\mathbf{x} := [x_1, \ldots, x_n] \qquad \mathbf{y} := [y_1, \ldots, y_n] \qquad \mathbf{z} := [z_1, \ldots, z_n]$$

$$\mathbf{x} \leftarrow \sum_{i=1}^{k} \left(\mathbf{x}^T \mathbf{e}_i\right) \mathbf{e}_i \qquad \mathbf{y} \leftarrow \sum_{i=1}^{k} \left(\mathbf{y}^T \mathbf{e}_i\right) \mathbf{e}_i \qquad \mathbf{z} \leftarrow \sum_{i=1}^{k} \left(\mathbf{z}^T \mathbf{e}_i\right) \mathbf{e}_i$$

Spectral Analysis

1. Setup Laplace-Beltrami matrix **L**
2. Compute *k* smallest eigenvectors {$\mathbf{e}_1$,..., $\mathbf{e}_k$}
3. Reconstruct mesh from those

Levy, Zhang: *Spectral Mesh Processing*, SIGGRAPH Asia Course, 2009

Spectral Analysis

1. Setup Laplace-Beltrami matrix **L**
2. Compute *k* smallest eigenvectors $\{\mathbf{e}_1, \ldots, \mathbf{e}_k\}$
3. Reconstruct mesh from those

Too complex for large meshes!

Bruno Levy: *Laplace-Beltrami Eigenfunctions: Towards an algorithm that understands geometry*, Shape Modeling and Applications, 2006

Review: Curves

Differential Geometry of Curves

- chord length and arc length
- arc-length parameterization
- Frenet frame

$$s = s(t) = \int_a^t \|\mathbf{x}_t\| \, \mathrm{d}t$$

$$\mathbf{t} = \frac{\mathbf{x}_t}{\|\mathbf{x}_t\|}$$

$$\mathbf{n} = \mathbf{b} \times \mathbf{t}$$

$$\mathbf{b} = \frac{\mathbf{x}_t \times \mathbf{x}_{tt}}{\|\mathbf{x}_t \times \mathbf{x}_{tt}\|}$$

Frenet frame

Review: Curves

Differential Geometry of Curves

– ***Frenet-Serret***: Derivatives w.r.t. arc length *s*

$$\begin{aligned} \mathbf{t}_s &= & +\kappa\mathbf{n} & \\ \mathbf{n}_s &= -\kappa\mathbf{t} & & +\tau\mathbf{b} \\ \mathbf{b}_s &= & -\tau\mathbf{n} & \end{aligned}$$

– **Curvature** (*deviation from straight line*)

$$\kappa = \|\mathbf{x}_{ss}\|$$

– **Torsion** (*deviation from planarity*)

$$\tau = \frac{1}{\kappa^2}\det([\mathbf{x}_s, \mathbf{x}_{ss}, \mathbf{x}_{sss}])$$

Review: Surface

Differential Geometry of Surfaces

– **First fundamental form**

$$\mathbf{I} = \begin{pmatrix} E & F \\ F & G \end{pmatrix} := \begin{pmatrix} \mathbf{x}_u^T \mathbf{x}_u & \mathbf{x}_u^T \mathbf{x}_v \\ \mathbf{x}_u^T \mathbf{x}_v & \mathbf{x}_v^T \mathbf{x}_v \end{pmatrix} \qquad \left\langle \begin{pmatrix} \alpha_1 \\ \beta_1 \end{pmatrix}, \begin{pmatrix} \alpha_2 \\ \beta_2 \end{pmatrix} \right\rangle := \begin{pmatrix} \alpha_1 \\ \beta_1 \end{pmatrix}^T \mathbf{I} \begin{pmatrix} \alpha_2 \\ \beta_2 \end{pmatrix}$$

- **Angle**

$$\mathbf{t}_1^T \mathbf{t}_2 = \langle (\alpha_1, \beta_1), (\alpha_1, \beta_1) \rangle$$

- **Length**

$$\mathrm{d}s^2 = E\mathrm{d}u^2 + 2F\mathrm{d}u\mathrm{d}v + G\mathrm{d}v^2$$

- **Area**

$$\mathrm{d}A = \sqrt{EG - F^2}\mathrm{d}u\,\mathrm{d}v$$

Review: Surface

Differential Geometry of Surfaces

– **Second fundamental form**

$$\mathbf{II} = \begin{pmatrix} e & f \\ f & g \end{pmatrix} := \begin{pmatrix} \mathbf{x}_{uu}^T \mathbf{n} & \mathbf{x}_{uv}^T \mathbf{n} \\ \mathbf{x}_{uv}^T \mathbf{n} & \mathbf{x}_{vv}^T \mathbf{n} \end{pmatrix}$$

- **normal curvature**

$$\kappa_n(\bar{\mathbf{t}}) = \frac{\bar{\mathbf{t}}^T \mathbf{II}\, \bar{\mathbf{t}}}{\bar{\mathbf{t}}^T \mathbf{I}\, \bar{\mathbf{t}}} = \frac{ea^2 + 2fab + gb^2}{Ea^2 + 2Fab + Gb^2}$$

- **principal curvatures**

$$\kappa_1 = \max_{\phi} \kappa_n(\phi) \qquad \kappa_2 = \min_{\phi} \kappa_n(\phi)$$

- **mean and Gaussian curvature**

$$H = \frac{\kappa_1 + \kappa_2}{2} \qquad K = \kappa_1 \cdot \kappa_2$$

Outline

smoothing intro:

① $p^{t+1} = p^t + \lambda \Delta p^t$ ⇒ explicit

↓ to adress MM

② $p^{t+1} = p^t + \lambda \Delta p^{t+1}$ ⇒ implicit

$(I - \lambda\Delta)p^{t+1} = p^t$

$p^{t+1} = (I - \lambda\Delta)^{-1} p^t$

$(I - \lambda M^{-1} C)p^{t+1} = p^t$

$M(I - \lambda M^{-1} C)p^{t+1} = M p^t$

$\Rightarrow (M - \lambda C)\, p^{t+1} = M p^t$

smoothing – explicit
– implicit

- Spectral Analysis
- **Diffusion Flow**
- Energy Minimization

Diffusion Flow on Height Fields

diffusion constant

$$\frac{\partial f}{\partial t} = \lambda \Delta f$$

Laplace operator

Diffusion Flow on Meshes

- Iterate $\mathbf{p}_i \leftarrow \mathbf{p}_i + \lambda \Delta \mathbf{p}_i$

0 Iterations

5 Iterations

20 Iterations

Uniform Laplace Discretization

$$\mathbf{p}_i \leftarrow \mathbf{p}_i + \lambda \Delta \mathbf{p}_i$$

- Smoothes geometry and triangulation
- Can be non-zero even for planar triangulations
- Vertex drift can lead to distortions
- Might be desired for mesh regularization

subtraction

ΔS 用 $\cot\alpha + \cot\beta$ 来近似

Desbrun et al., Siggraph 1999

Curvature Flow

- Use diffusion flow with Laplace-Beltrami

$$\frac{\partial \mathbf{p}}{\partial t} = \lambda \Delta_{\mathcal{S}} \mathbf{p}$$

- Laplace-Beltrami is parallel to surface normal

$$\frac{\partial \mathbf{p}}{\partial t} = -2\lambda H \mathbf{n}$$

➡ Avoids vertex drift on surface

Comparison

niloy 说这了
只在地心坐

Original | Uniform Laplace | Laplace-Beltrami

Numerical Integration

- Write update $\mathbf{p}_i^{(t+1)} = \mathbf{p}_i^{(t)} + \lambda \Delta \mathbf{p}_i^{(t)}$ in matrix notation

$$\mathbf{P}^{(t)} = \left(\mathbf{p}_1^{(t)}, \ldots, \mathbf{p}_n^{(t)}\right)^T \in \mathbb{R}^{n \times 3}$$

千万别用这个

$$\mathbf{p}_i \leftarrow \mathbf{p}_i + \lambda \Delta \mathbf{p}_i$$

- Corresponds to **explicit** integration

用这个

$$\mathbf{P}^{(t+1)} = (\mathbf{I} + \lambda \mathbf{L})\, \mathbf{P}^{(t)}$$

Requires small λ for stability!

- **Implicit** integration is *unconditionally stable*

$$(\mathbf{I} - \lambda \mathbf{L})\, \mathbf{P}^{(t+1)} = \mathbf{P}^{(t)}$$

Implementation

- Solve linear system each iteration

$$(\mathbf{I} - \lambda \mathbf{L})\mathbf{P}^{(t+1)} = \mathbf{P}^{(t)}$$

- Matrix $L = M^{-1}L_w$ is not symmetric because of M
  - ➡ Symmetrize by multiplying M from left

$$(\mathbf{M} - \lambda \mathbf{L}_w)\mathbf{P}^{(t+1)} = \mathbf{M}\mathbf{P}^{(t)}$$

- Solve sparse symmetric positive definite system
  - ➡ Iterative conjugate gradients, sparse Cholesky

Outline

- Spectral Analysis
- Diffusion Flow [smoothing]
- **Energy Minimization**

Fairness

- Idea: Penalize “unaesthetic behavior”
- **Measure** fairness
  - Principle of the simplest shape
  - Physical interpretation
- **Minimize** some fairness functional
  - Surface area, curvature
  - Membrane energy, thin plate energy

Non-Linear Energies

- Membrane energy (surface area)

$$\int_{\mathcal{S}} \mathrm{d}A \;\rightarrow\; \min \quad \text{with} \quad \delta\mathcal{S} = \mathbf{c}$$

surface

- Thin-plate surface (curvature)

$$\int_{\mathcal{S}} \kappa_1^2 + \kappa_2^2 \,\mathrm{d}A \;\rightarrow\; \min \quad \text{with} \quad \delta\mathcal{S} = \mathbf{c}, \;\; \mathbf{n}(\delta\mathcal{S}) = \mathbf{d}$$

normal

- Too complex ... simplify energies

Membrane Surfaces

- Surface parameterization [eg. texture].

$$\mathbf{x} : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$$

- Membrane energy (surface area)

$$\min_{\mathbf{x}} \int_{\Omega} \|\mathbf{x}_u\|^2 + \|\mathbf{x}_v\|^2 \mathrm{d}u\mathrm{d}v$$

Variational Calculus in 1D

- 1D membrane energy

$$L(f) = \int_a^b f'^2(x)\,\mathrm{d}x \;\rightarrow\; \min$$

- Add test function *u* with *u*(*a*) = *u*(*b*) = 0

$$L(f+\lambda u) = \int_a^b (f'+\lambda u')^2 = \int_a^b f'^2 + 2\lambda f' u' + \lambda^2 u'^2$$

- If *f* minimizes *L*, the following has to vanish

$$\left.\frac{\partial L(f+\lambda u)}{\partial \lambda}\right|_{\lambda=0} = \int_a^b 2 f' u' \;\stackrel{!}{=}\; 0$$

Variational Calculus in 1D

- Has to vanish for any u with $u(a) = u(b) = 0$

$$\int_a^b f'u' = \underbrace{[f'u]_a^b}_{=0} - \int_a^b f''u \overset{!}{=} 0 \quad \forall u$$

$$\int_0^1 x'y = [xy]_0^1 - \int_0^1 xy'$$

- Only possible if

$$f'' = \Delta f = 0$$

➡ ***Euler-Lagrange equation***

Bivariate Variational Calculus

- Find minimum of functional

$$\underset{f}{\operatorname{argmin}} \int_{\Omega} L\left(f_{uu}, f_{vv}, f_u, f_v, f, u, v\right)$$

- Euler-Lagrange PDE defines the minimizer *f*

$$\frac{\partial L}{\partial f} - \frac{\partial}{\partial u}\frac{\partial L}{\partial f_u} - \frac{\partial}{\partial v}\frac{\partial L}{\partial f_v} + \frac{\partial^2}{\partial u^2}\frac{\partial L}{\partial f_{uu}} + \frac{\partial^2}{\partial u \partial v}\frac{\partial L}{\partial f_{uv}} + \frac{\partial^2}{\partial v^2}\frac{\partial L}{\partial f_{vv}} = 0$$

  – Again, subject to suitable boundary constraints

Membrane Surfaces

- Surface parameterization

$$\mathbf{x} : \Omega \subset \mathbb{R}^2 \to \mathbb{R}^3$$

- Membrane energy (surface area)

$$\min_{\mathbf{x}} \int_{\Omega} \|\mathbf{x}_u\|^2 + \|\mathbf{x}_v\|^2 \mathrm{d}u\mathrm{d}v \quad ①$$

$$\Updownarrow$$

解出 ②

就是什么 ①

- Variational calculus

$$\Delta \mathbf{x} = 0 \quad ②$$

Laplace-Beltrami Operator

- Extension of Laplace to functions on manifolds

Laplace-Beltrami

gradient operator

mean curvature

$$\Delta_{\mathcal{S}}\mathbf{x} = \mathrm{div}_{\mathcal{S}}\, \nabla_{\mathcal{S}}\mathbf{x} = -2H\mathbf{n}$$

$$\nabla\mathbf{x} = [\partial\mathbf{x}/\partial u \quad \partial\mathbf{x}/\partial v]$$

$$\mathrm{div}\mathbf{y} := \nabla \cdot \mathbf{y} = \partial\mathbf{y}_u/\partial u + \partial\mathbf{y}_v/\partial v$$

$$\mathrm{div}(\nabla\mathbf{x}) = \Delta\mathbf{x}$$

Thin-Plate Surfaces

- Surface parameterization

$$\mathbf{p} : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$$

曲率

- Thin-plate energy (curvature)

$$\int_\Omega \|\mathbf{p}_{uu}\|^2 + 2\,\|\mathbf{p}_{uv}\|^2 + \|\mathbf{p}_{vv}\|^2 \,\mathrm{d}u\mathrm{d}v \;\rightarrow\; \min$$

找出曲率最小

- Variational calculus

$$\Delta^2 \mathbf{p} = 0$$

Energy Functionals

[这章在讲 边界问题]

given 边界条件

Membrane

$$\Delta_{\mathcal{S}}\mathbf{p} = 0$$

minimize 表面积 曲面光滑 [但是没提到]

Thin Plate

$$\Delta_{\mathcal{S}}^{2}\mathbf{p} = 0$$

$$\Delta_{\mathcal{S}}^{3}\mathbf{p} = 0$$

Analysis

- Minimizer surfaces satisfy Euler-Lagrange PDE

$$\Delta_{\mathcal{S}}^{k}\mathbf{p} = 0$$

- They are ***stationary surfaces of Laplacian flows***

$$\frac{\partial \mathbf{p}}{\partial t} = \Delta_{\mathcal{S}}^{k}\mathbf{p}$$

- Explicit flow integration corresponds to iterative solution of linear system

References

- Book: Chapter 4 (Botsch et al.),
  Chapter 5 (CohenOr et al.)
- Levy: *Laplace-Beltrami Eigenfunctions: Towards an algorithm that understands geometry*, Shape Modeling and Applications, 2006
- Taubin: *A signal processing approach to fair surface design*, SIGGRAPH 1996
- Desbrun, Meyer, Schroeder, Barr: *Implicit Fairing of Irregular Meshes using Diffusion and Curvature Flow*, SIGGRAPH 99