

# 1 Kepler's Second Law Derivation

## 1.1 Review of Calculus

### 1.1.1 Limits and Derivatives

Limits describe the behavior of a function near a point. For example,  $\lim_{x \rightarrow 2} x + 4 = 6$  since  $2 + 4 = 6$ . Special limits like

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

represent the derivative, or instantaneous slope, of a function at a point. Similarly, you may take for granted that an expression like

$$\lim_{t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}.$$

### 1.1.2 Integration

The power rule of integration for any function  $x$  is as follows:


$$\int x^n dx = \frac{x^{n+1}}{n+1} + C.$$

## 1.2 Definition

A line joining a planet and the Sun sweeps out equal areas during equal intervals of time.

## 1.3 Derivation

For this line to sweep out equal area during equal time, the rate at which the area changes,  $\frac{dA}{dt}$  must stay constant. One small slice of the area of a planet's elliptical orbit may be represented as a triangle



with base  $r$ , height  $r\Delta\theta$ , and area  $\Delta A$  (recall that arc length is  $r\Delta\theta$ ). Recall the equation for the area of a triangle,

$$A = \frac{1}{2}bh.$$

Substituting the givens from our slice above, we get

$$\Delta A = \frac{1}{2}(r)(r\Delta\theta).$$

Divide both sides by  $\Delta t$

$$\frac{\Delta A}{\Delta t} = \frac{1}{2}(r)(r\frac{\Delta\theta}{\Delta t}).$$

Take the limit of both sides as  $\Delta t \rightarrow 0$  to find an “infinitely” small change of area over an “infinitely” small change of time (keep in mind that  $r$  is constant for our slice and that area  $A$  is a function of time).

$$\lim_{\Delta t \rightarrow 0} (\frac{\Delta A}{\Delta t}) = \lim_{\Delta t \rightarrow 0} (\frac{1}{2}(r)(r\frac{\Delta\theta}{\Delta t})),$$

which simplifies to

$$\frac{dA}{dt} = \frac{1}{2}(r)(r\frac{d\theta}{dt})$$

and further simplifies to

$$\frac{dA}{dt} = \frac{1}{2}rv_t.$$

Recall that angular momentum is

$$L = mrv_t$$

and is a conserved quantity. Substituting  $L$  into our area gets

$$\frac{dA}{dt} = \frac{L}{2m}.$$

We can then rearrange and solve the differential equation

$$\int dA = \int \frac{L}{2m} dt,$$

which shows us that

$$A = \frac{L}{2m}T.$$

This shows us a directly proportional relationship between the area  $A$  and time  $T$ .