



Digestive tract mycobiota and microbiota and the effects on the immune system

Nina Gouba^{a,*}, Yeri Esther Hien^b, Marie Laure Guissou^c, Maxime Descartes Mbogning Fonkou^d, Yves Traoré^b, Zekiba Tarnagda^e

^a Université Nazi Boni de Bobo-Dioulasso, 01 BP 1091 Bobo-Dioulasso 01, Burkina Faso

^b Unité de Formation et de Recherche en Science de la Vie et de la Terre, Laboratoire de Biochimie et d'Immunologie Appliquée (LaBIA), Université Ouaga 1 Pr Joseph Ki-Zerbo, 03 B. P 7021 Ouagadougou 03, Burkina Faso

^c Université Norbert ZONGO, BP 376 Koudougou, Burkina Faso

^d Aix Marseille University, IRD, AP-HM, MEPHI, IHU Méditerranée-Infection, Marseille, France

^e Institut de Recherche en Science de la Santé Direction Régionale de Bobo-Dioulasso, IRSS-DRO, 399 avenue de la liberté 01, BP 545, Burkina Faso

ARTICLE INFO

Keywords:

Human digestive tract
Microbiota
Mycobiota
Immune system
Dysbiosis

ABSTRACT

The human gastrointestinal tract exists as a complex ecosystem and contains a mycobiome and a microbiome that play central roles in host health, disease and immune system regulation. Here, we reviewed the traditional culture-dependent methods, the culturomics methods and the molecular methods used to study the gut mycobiome and microbiome. With the development of next-generation sequencing techniques, these last two methods have greatly broadened the understanding of the roles of gut bacteria in health and disease. Thus, dysbiosis of the gut microbiota and mycobiota has been found to be associated with some diseases; disruptions to the fungal and bacterial commensal communities influence the immune response and impact disease status.

1. Introduction

The human digestive tract microbiome is a complex ecosystem that harbours a diversity of microorganisms, including Bacteria, Archaea, viruses and Eukarya [1]. This microbial ecosystem plays a central role in host development, including immune system development, physiology, digestion, vitamin and nutrient production and protection against pathogen colonization [2]. The gut microbiota (bacterial community) composition is influenced by interactions among community members, the genetics of the host, dietary habits and the environment [3]. Bacteria are the most dominant group in the community and have the four following major phyla residents: *Firmicutes*, *Bacteroidetes*, *Actinobacteria* and *Proteobacteria* [1]. Therefore, the bacterial community has been well characterized and has been the focus of human microbiome research in health and disease. The mycobiota (fungal community) represents the most significant group of eukaryotes in the human gut microbiota and has been found to be associated with a wide range of diseases, such as hepatitis, inflammatory bowel diseases, and systemic mycosis [4]. In recent years, the fungal community has been gaining attention, but the field of mycobiome research is still in its infancy. The mycobiome has been traditionally studied using culture-dependent methods. To date, to investigate the role of the fungal

community in both health and disease, researchers have begun to use next-generation sequencing (NGS) technology [5]. We review here the recent knowledge from the study of digestive tract mycobiota and microbiota in healthy and diseased individuals and its interaction with the host's immune system.

2. Digestive tract mycobiota and microbiota characterization

2.1. Culture-dependent methods

The preceding investigations of digestive tract fungal and bacterial diversity were based on culture methods using classic selective media to isolate fungi and bacteria [5]. Yeast and filamentous fungi are cultured in different culture media, including on Sabouraud dextrose agar, malt agar, potato dextrose agar, CZAPEK, Columbia agar [4]. A mix of broad-spectrum antibiotics are then added into the culture media to limit bacterial growth [4]. The isolated colonies are identified by classic phenotypic identification methods based on microscopic and biochemical tests. Thus, Gram, Ziehl-Neelsen, and Giemsa stains are used in bacterial microscopic identifications, while lactophenol staining is used for fungal macro- and microscopic identifications. Based on the traditional culture methods, approximately 688 bacterial species, 2

* Corresponding author at: Université Nazi Boni de Bobo-Dioulasso, 01 BP 1091 Bobo-Dioulasso 01, Burkina Faso.

E-mail address: goubanina@yahoo.fr (N. Gouba).

<https://doi.org/10.1016/j.humic.2019.100056>

Archaea and 88 fungal species have been isolated from the human gut [4,6]. This may reflect the approximative gut composition of bacteria at 10^{11-12} CFU per gram of faeces and of fungi at 0 up to 10^6 CFU per gram of faeces, as reported in the literature [7,8]. The fungal species commonly isolated belong to genera of *Candida*, *Aspergillus*, and *Penicillium*, and the nine bacterial genera belong to four phyla (*Actinobacteria*, *Proteobacteria*, *Firmicutes*, *Bacteroidetes*) [1,6]. However, traditional methods present some disadvantages, such as misidentification and time requirements. The major bias associated with traditional culture methods is the fast growth of the major population in classic nutrient growth media; thus, the minor populations, including some potential pathogenic populations, such as *Salmonella typhi* [6], are missed. A recent alternative tool, such as Matrix Assisted Laser Desorption Ionisation – Time of Flight – Mass Spectrometry (MALDI-TOF-MS), is used for both fungal and bacterial colony identifications [9,10]. The MALDI-TOF MS-based identification technique analyses the protein content from treated or intact microorganisms in the form of a spectrum that is considered to be a protein fingerprint that is specific to a microorganism. An unknown bacterial or fungal colony is identified by comparing its spectrum with the spectra in the reference database [9]. MALDI-TOF MS-based identification is simple, fast, accurate and allows for the rapid screening of large numbers of colonies isolated in culture [10]. Culturomics methods, which consist of the application of high-throughput culture conditions followed by MALDI-TOF MS and/or 16S rRNA identification, allow the known diversity of the gut microbiota to be expanded [10]. This approach has recently added 531 bacterial species to the human gut bacterial repertoire [11]. The first study that used a culturomics approach to characterize the gut mycobiota isolated 41 diverse fungal species from 14 faecal samples, of which 10 species have been reported for the first time in the human gut [12]. The advantages of culturomics are that it allows the growth of minor populations under various culture conditions and that it allows the culture of uncultured organisms that correspond to sequences not previously assigned [11]. However, there are no standard culture conditions published for mycobiome studies as there are for bacterial culturomics conditions. The bias associated with culture conditions is the lack of some ‘non-cultivable’ microorganisms. Additionally, the culturomics technique does not directly provide information on the enzymatic abilities of the microorganisms isolated and requires a major workload [13].

2.2. Culture-independent methods

Molecular tools have emerged since 1990 for digestive tract mycobiota and microbiota investigations. PCR-based methods used primers that targeted 16S rRNA in microbiota studies and primers that targeted 18S, 5.8S, 26S, and 28S rRNA and the ITS (Internal Transcribed Spacer) regions in mycobiome studies. The variability of the ITS region enables the classification of fungi into the genera and species levels. Then, computational algorithms are used to align the operational taxonomic unit (OTUs) with those from the fungal database. Other molecular techniques, such as cloning libraries as rDNA-based denaturing gradient gel electrophoresis (DGGE), restriction fragment length polymorphism (RFLP) analysis, in situ hybridization techniques, and Sanger sequencing, have been used on stool samples or digestive mucosa biopsies to investigate the fungal and bacterial diversity [1,14,15]. All these molecular techniques helped to identify more than 30 bacterial phyla (major phyla of *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Fusobacteria*, *Proteobacteria*, *Verrucomicrobia*, and *Cyanobacteria*), whereas only three fungal phyla (*Ascomycota*, *Basidiomycota*, and *Zygomycota*) are commonly reported [1,15]. Since 2006, metagenomic studies have been used to explore gut bacterial diversity. However, this approach was not widely applied in gut mycobiota analysis. The variation and relative abundance of bacterial phyla in several studies depend significantly on the 16S hypervariable region and on the DNA extraction kit used [6].

More recently, NGS (next-generation sequencing) technologies

including Roche 454, Illumina HiSeq and MiSeq, and Ion Torrent have been used to characterize the digestive tract mycobiome and microbiome in patients and healthy persons in a number of studies [12,16,17]. The high variability of reported fungi in these different studies seems to depend on host status, environment, diet and choices of primers and DNA extraction techniques. Despite the progress in technology, there are limited databases available for NGS data generated for mycobiome analyses compared to the number of bacterial databases [18].

The combination of culturomics and next-generation sequencing approaches is helpful for human digestive tract mycobiome analysis and can contribute to reducing the gap in gut mycobiota knowledge.

3. Gut mycobiota and microbiota interactions in healthy and diseased participants

3.1. In healthy participants or healthy volunteers

There are no specific fungi described in the literature that are only specific to healthy people. Several studies of the gut mycobiome in healthy participants or in healthy volunteers have reported fungal genera belonging to *Ascomycota*, *Basidiomycota* and *Zygomycota* [4] with the predominance of *Candida*, *Saccharomyces*, *Trichosporon* and *Cladosporium* [19]. *Candida* species are usually described as commensal yeasts in individuals with healthy gastrointestinal tracts. The first investigation combined the culture and molecular methods and reported a low diversity of fungi, approximately 1 to 3 fungal species per individual, including *Candida parapsilosis*, *Paecilomyces fumosoroseus*, and *Gloeotinia temulenta* [8]. Further, a large diversity and high variability of fungal species have been reported in healthy individuals in different studies [16,17,20,21]. Some fungi have been described as more prevalent in non-obese patients, such as *Mucor*, *Candida* and *Penicillium*, whereas *Candida*, *Nakaeseomyces*, *Penicillium* and *Pichia* (teleomorph of *Candida* spp.) are found predominantly in obese patients [19,22]. Most of the fungal species reported in these works were from the environment (*Cladosporium* sp., *Puccinia poarum*, *Aspergillus restrictus*), diet (*Penicillium roqueforti*, *Saccharomyces cerevisiae*) or opportunist pathogens (*C. albicans*, *Candida glabrata*, and *Trichosporon asahii*) [20,23]. The mycobiome reported in healthy humans revealed a high inter-individual variability that was much less persistent in individuals and was instable over time [16]. The diversity and variability of fungal species observed in healthy individuals could be related to differences in geographical location and lifestyle, as has been observed in bacterial studies. However, further studies are needed to clarify the transmission of residential gut fungal species. Knowledge is limited on specific role of fungal species found in healthy gut compared to bacteria. Bacterial species that belong to the phyla *Actinobacteria*, *Bacteroidetes*, *Firmicutes* and *Proteobacteria* are more diverse and abundant in healthy guts. Different compositions of gut microbiota have been reported according to different locations, ages, sexes, races, and diets of the host [24]. The bacterial community has a low diversity during the first year of life and develops gradually during the period of growth. In contrast to the gut mycobiota, the bacterial community is relatively stable in adults over time [25]. Diet can impact and shape the composition of the gut microbiota. For example, the *Bacteroides* genus has been found to be associated with the consumption of animal proteins, while *Prevotella* are found to be associated with diets rich in carbohydrates [26]. The dominance of *Bacteroides* has been described in the microbiota from European patients, whereas *Prevotella* was found to be dominant in those from Africa [27]. Additionally, regarding the interaction between the bacterial and fungal communities in healthy participants, Hoffmann et al. reported a significant correlation between both *Ascomycota* and *Basidiomycota* with some bacterial lineages [17]. Indeed, this author observed a positive association between *Candida* and *Methanobrevibacter* with the *Prevotella* group, and each of these was found to be correlated with a diet rich in carbohydrates. Now, we know more about

some of the functions and roles of the gut microbiome, and further studies are needed to elucidate the roles of the gut mycobiome in healthy participants.

3.2. In diseased participants

Candida species have been primarily described in various diseases, such as inflammatory bowel disease (IBD), Crohn's disease (CD), ulcerative colitis (UC), and gut inflammation [4]. Opportunist pathogenic yeasts are commonly described in immunodeficient patients (HIV, cancer, and transplanted patients) [4]. The non-albicans fungal species of *Candida*, *Trichosporon*, *Rhodotorula*, *Cryptococcus neoformans*, and *Geotrichum* are increasingly emerging in the frequent diagnosis of immunodeficient patients [28]. Additionally, an increased diversity and richness of the gut mycobiome have been described in IBD, CD, UC, and hepatitis B patients compared to those in controls [14,15]. In IBD patients, Sokol et al. [29] observed a dysbiosis of the gut mycobiome with a decreased proportion of *Saccharomyces cerevisiae* and an increased proportion of *C. albicans* compared with those in the healthy controls. Fungal microbiota dysbiosis in IBD is characterized by alterations in biodiversity and composition. This observation suggested that the mycobiome might play a role in inflammatory bowel disease. Another investigation on the gut mycobiome and bacterial dysbiosis in infants reported that there was a positive association between the development of atopic wheeze, an increase in the fungal species *Pichia kudriavzevii* and a decrease in *Saccharomycetales* [30]. Similar to fungi, bacterial dysbiosis has been found to be associated with a number of diseases, including IBD, diabetes, allergy, obesity, central nervous diseases, colorectal cancer and infectious diseases [31]. Previous studies have observed a decrease in the abundance and diversity of Firmicutes in IBD patients [25]. Antibiotic treatments in infectious diseases alter the gut microbiota composition and reduce the resistance to pathogen colonization. An increase in susceptibility to *Clostridium difficile* colonization has been reported in patients with antibiotic treatment compared to that in healthy controls [25]. In diabetic patients, an alteration and a low diversity of gut microbiota have been reported [31]. A decrease in the diversity of *Bacteroides* has been reported in patients suffering from allergies. Additionally, another study observed a high abundance of some bacterial species of *Streptococcus* sp. and *Bacteroides* sp. As well as a decrease in *Ruminococcus gnavus* and *Bifidobacterium* sp. in atopic wheeze patients compared to the abundance in healthy controls [30]. The gut microbiota role is described in a number of diseases, but the mycobiome role remains unclear; therefore, further investigations are needed to elucidate the role of fungi in disease.

4. Microbiota, mycobiota and host immunity

4.1. The impact of microbiota on the immune system

Technological advances are shedding new light on the sophisticated ways in which microbes influence human health; one of the major impacts of the mammalian microbiota is its effect on the development and function of the immune system. Communities of bacteria and immune cells are closely linked, especially those residing in the intestinal tract [32]. This interaction is likely to be a key factor in the maturation of both T helper 1 (Th1) cells [33] and regulatory T cell (Treg) functions [34] and in the functional suppression of the normal neonatal propensity for T helper 2 (Th2) cell responses, which are still dominant in the perinatal period. While early hypotheses have suggested that these host-microbial interactions were largely initiated at the beginning of the postnatal period, microbes have now been detected in the placental and foetal tissues during normal pregnancies [35]. This is likely to suggest that maternal microbial transfer to the offspring begins during pregnancy. Therefore, variations in the maternal environment appear to modulate foetal immune responses, with measurable differences in both effector and Treg cell responses in the cord blood of newborns whose

mothers inhabit more diverse microbial environments compared to the responses in those who do not [36]. Among mothers living on farms, Treg cells were mainly at higher levels and were more efficient in offspring after specific stimulation compared with those of nonfarming mothers. This was associated with decreased Th2 cytokine levels and less cell proliferation in the offspring of farming mothers. Exposure to several animal species also has an effect on Treg cell markers and interferon gamma (IFN- γ) secretion [37]. These well-balanced regulatory responses constrain proinflammatory Th1, Th2 and Th17 clones and curb the risk of chronic inflammatory diseases, such as allergies and autoimmune disease [38]. This has been well illustrated in rural Austria, where children in settings with rich microbial exposure have a reduced risk of allergies [37], and antenatal exposure largely accounts for this protective effect [39]. Experimental animal models also demonstrate that maternal exposure to commensal bacteria, such as *Lactobacillus rhamnosus* [40], during pregnancy can attenuate allergic sensitization and inflammation in the offspring. Thus, microbial contact during prenatal life may imprint the offspring microbiota and immune system in preparation for the much larger inoculum transferred during vaginal delivery and breastfeeding.

4.2. Fungal interactions with the human immune system

Commensal fungi are part of the intestinal microbial community in many species, such as rats, guinea pigs, rabbits, pigs, dogs, and humans, with the highest densities in the terminal colon [41]. Although few studies have suggested the presence of commensal fungi in the gut [14,41], their interaction with the immune system of the mucosa or their involvement in diseases is poorly understood.

Various specific immune mechanisms have been implicated in the clearance of fungi at mucosal surfaces, and many innate immune receptors have been shown to interact with fungal pathogens [42,43]. Among the mechanisms implicated in the clearance of fungi, C-type lectin domain family 7, member A (Dectin-1), caspase recruitment domain-containing protein 9 (CARD9), interleukin-17 (IL-17) and IL-22 have emerged as molecules that are responsible for host defence. For example, a genetic mutation in each of these molecules is associated with susceptibility to fungal infections in humans. A Dectin-1-deficient mouse model showed increased susceptibility to colitis that was characterized by increased mucosal erosion, inflammatory cell infiltration, TNF- α production and the augmented production of IFN- γ and IL-17. The more severe colitis cases in mice were accompanied by a fungal invasion of the colonic mucosa, as well as a general expansion of opportunistic fungi, such as *Candida* and *Trichosporon* species, and a decrease in the non-pathogenic *Saccharomyces* species. A Dectin-1 deficiency leads to altered immune responses by the commensal fungi in the gut; this inability to mount effective immune responses to specific intestinal fungi creates conditions that promote inflammation.

Intestinal fungi can influence immunity through both direct interactions with immune cells and the production of metabolites, such as prostaglandin E2 (PGE2), a potent immunomodulator produced by immune cells. It has been proposed that *Candida*-produced PGE2 can reach the lungs through the bloodstream, act on lung macrophages and promote allergic inflammation [44]. In addition, *Candida* can activate host cells to produce prostaglandins [45], suggesting a possibility that the overgrowth of the gut fungal microbiota may alter immune responses via PGE2. *Candida* also produces ligands for pattern recognition receptors (PRRs), including β -glucans, chitin, mannans, β -(1,2)-linked oligomannosides and fungal nucleic acids [46], which stimulate the innate immune response.

Moreover, fungi detected by the intestinal immune system can cause inflammation or tolerant immune responses and can modulate immune cell trafficking. Thus, phagocytes that digest fungi can release fungal-derived molecules into the circulation and may have immunoregulatory effects [47].

4.3. Dysbiosis and host immunity

The composition of the gut microbiota is more stable throughout adult life than the mycobiota is but can be altered as a result of bacterial infections, antibiotic treatment, lifestyle, surgery, and a long-term change in diet. Changes in the gut bacteria are initially described as “dysbiosis”, a disruption of the healthy microbial communities that allows the overgrowth of fungi [48]; this indicates that, in the steady state, the bacterial communities keep fungi in check. Fungal overgrowth in the gut can induce macrophage polarization that increases allergic airway inflammatory cell infiltration [44]. In particular, *Candida* spp. infection can induce the production of inflammatory mediators by host cells. In patients with IBD, the disease has been reported to be associated with enrichment for *Candida* species (specifically *Candida dubliniensis* and *Candida parapsilosis*), suggesting that fungal dysbiosis may accompany disease. Previous results have shown that patients with diseases, such as ulcerative colitis, Crohn’s disease, and gastric ulcers, were more heavily colonized with the fungus *C. albicans* [49]. However, although the study of the fungal microbiota is a rapidly emerging field, the mechanisms by which gut dysbiosis drives fungal overgrowth in the gut and affects the host immune responses remain poorly understood.

Similarly, treating animals with oral antifungal drugs has been reported to alter the bacterial microbiota. Antifungal treatment affects specific bacterial taxa with a decrease in the relative detection levels of *Bacteroides*, *Allobaculum*, *Clostridium*, *Desulfovibrio*, and *Lactobacillus* spp., while the relative detection of *Anaerostipes*, *Coprococcus*, and *Streptococcus* is increased [50]; this results in the alteration of the local and peripheral immune homeostasis. Matthew L. Wheeler et al. [50] showed that the treatment of mice with antifungal drugs increased the disease severity in chemically induced and T cell transfer-mediated models of experimental colitis and exacerbated the development of house dust mite-induced allergic airway disease. This provides compelling evidence to support a functional role of mycobiota in the modulation of immune function and the development of inflammatory disease.

5. Conclusion

Here, we have reviewed and highlighted the methods used to investigate the gut mycobiome and microbiome. We also reported the fungal and bacterial communities found in human health and disease and their interactions with the immune response. Compared to the roles of bacteria, the roles of the gut fungal community remain poorly understood in health and disease and require further investigation.

Declaration of Competing Interest

The authors declare no competing interests.

References

- Rajilić-Stojanović M, Smidt H, De Vos Willem M. Diversity of the human gastrointestinal tract microbiota revisited. *Environ Microbiol* 2007;9(9):2125–36.
- Klein Tsai M, Coyle Walter J. The microbiome and obesity: is obesity linked to our gut flora? *Curr Gastroenterol Rep* 2009;11:307–13.
- Hoarau GMP, Gower-Rousseau C, Hager C, Chandra J, Retuerto MA, Neut C, Vermeire S, Clemente J, Colombel JF, Fujioka H, Poulain D, Sendid B, Ghannoum MA. Bacteriome and mycobiome interactions underscore microbial dysbiosis in familial Crohn’s disease. *mBio* 2016;7(5). p. e01250-16.
- Gouba N, Drancourt M. Digestive tract mycobiota: a source of infection. *Méd Maladies Infect* 2015;45(1–2):9–16.
- Huseyin CE, O’Toole PW, Cotter PD, Scanlan PD. Forgotten fungi—the gut mycobiome in human health and disease. *FEMS Microbiol Rev* 2017;41(4):479–511.
- Lagier J-C, Million M, Hugon P, Armougom F, Raoult D. Human gut microbiota: repertoire and variations. *Front Cell Infect Microbiol* 2012;2:136.
- Ley RE, Peterson DA, Gordon JI. Ecological and evolutionary forces shaping microbial diversity in the human intestine. *Cell* 2006;124(4):837–48.
- Scanlan PDM, Julian R. Micro-eukaryotic diversity of the human distal gut microbiota: qualitative assessment using culture-dependent and-independent analysis of faeces. *ISME J* 2008;2(12):1183.
- Cassagne C, Ranque S, Normand A-C, Fourquet P, Thiebault S, Planard C, et al. Mould routine identification in the clinical laboratory by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *PLoS One* 2011;6(12):e28425.
- Lagier J-C, Armougom F, Million M, Hugon P, Pagnier I, Robert C, et al. Microbial culturomics: paradigm shift in the human gut microbiome study. *Clin Microbiol Infect* 2012;18(12):1185–93.
- Lagier J-C, Khelaifia S, Alou MT, Ndongo S, Dione N, Hugon P, et al. Culture of previously uncultured members of the human gut microbiota by culturomics. *Nat Microbiol* 2016;1(12):16203.
- Hamad I, Ranque S, Azhar EI, Yasir M, Jiman-Fatani AA, Tissot-Dupont H, et al. Culturomics and amplicon-based metagenomic approaches for the study of fungal population in human gut microbiota. *Sci Rep* 2017;7(1):16788.
- Greub G. Culturomics: a new approach to study the human microbiome. *Clin Microbiol Infect* 2012;18(12):1157–9.
- Ott SJ, Kühbacher T, Musfeldt M, Rosenstiel P, Hellmig S, Rehman A, et al. Fungi and inflammatory bowel diseases: alterations of composition and diversity. *Scand J Gastroenterol* 2008;43(7):831–41.
- Yu Chen ZC, Guo Renyong, Chen Nan, Haifeng Lu, Huang Shuai, Wang Jie, Li Lanjuan. Correlation between gastrointestinal fungi and varying degrees of chronic hepatitis B virus infection. *Diagn Microbiol Infect Dis* 2011;2011(70):492–8.
- Hallen-Adams HE, Kachman SD, Kim J, Legge RM, Martinez I. Fungi inhabiting the healthy human gastrointestinal tract: a diverse and dynamic community. *Fungal Ecol* 2015;15:9–17.
- Hoffmann C, Dollive S, Grunberg S, Chen J, Li H, Wu GD, et al. Archaea and fungi of the human gut microbiome: correlations with diet and bacterial residents. *PLoS One* 2013;8(6):e66019.
- Huseyin CE, Rubio RC, O’Sullivan O, Cotter PD, Scanlan PD. The fungal frontier: a comparative analysis of methods used in the study of the human gut mycobiome. *Front Microbiol* 2017;8:1432.
- Witherden EA, Shoaie S, Hall RA, Moyes DL. The human mucosal mycobiome and fungal community interactions. *J Fungi* 2017;3(4):56.
- Gouba N, Raoult D, Drancourt M. Eukaryote culturomics of the gut reveals new species. *PLoS One* 2014;9(9):e106994.
- Hamad I, Sokhna C, Raoult D, Bittar F. Molecular detection of eukaryotes in a single human stool sample from senegal. *PLoS One* 2012;7(7):e40888.
- Gouba N, Raoult D, Drancourt M. Plant and fungal diversity in gut microbiota as revealed by molecular and culture investigations. *PLoS One* 2013;8(3):e59474.
- Hallen-Adams HE, Suhr MJ. Fungi in the healthy human gastrointestinal tract. *Virulence* 2017;8(3):352–8.
- Hollister EB, Gao C, Versalovic J. Compositional and functional features of the gastrointestinal microbiome and their effects on human health. *Gastroenterology* 2014;146(6):1449–58.
- Clemente JC, Ursell LK, Parfrey LW, Knight R. The impact of the gut microbiota on human health: an integrative view. *Cell* 2012;148(6):1258–70.
- Wang H, Wei C-X, Min L, Zhu L-Y. Good or bad: gut bacteria in human health and diseases. *Biotechnol Bioinform Equip* 2018:1–6.
- De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, et al. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci* 2010;107(33):14691–6.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infect Dis* 2011;11(2):142–51.
- Sokol H, Leducq V, Aschard H, Pham H-P, Jegou S, Landman C, et al. Fungal microbiota dysbiosis in IBD. *Gut* 2016. p. gutjnl-2015-310746.
- Arrieta M-C, Arévalo A, Stiemsma L, Dimitriu P, Chico ME, Loo S, et al. Associations between infant fungal and bacterial dysbiosis and childhood atopic wheeze in a nonindustrialized setting. *J Allergy Clin Immunol* 2018;142(2). pp. 424–434. e10.
- Wang B, Yao M, Lv L, Ling Z, Li L. The human microbiota in health and disease. *Engineering* 2017;3(1):71–82.
- Mathis D, Benoist C. Microbiota and autoimmune disease: the hosted self. *Cell Host Microbe* 2011;10(4):297–301.
- Prescott SL, Macaubas C, Smallacombe T, Holt BJ, Sly PD, Holt PG. Development of allergen-specific T-cell memory in atopic and normal children. *Lancet* 1999;353(9148):196–200.
- Tulic MK, Andrews D, Crook ML, Charles A, Tourigny MR, Moqbel R, et al. Changes in thymic regulatory T-cell maturation from birth to puberty: differences in atopic children. *J Allergy Clin Immunol* 2012;129(1). pp. 199–206. e4.
- Logan AC, Jacka FN, Prescott SL. Immune-microbiota interactions: dysbiosis as a global health issue. *Curr Allergy Asthma Rep* 2016;16(2):13.
- Schaub B, Liu J, Hoppeler S, Schleich I, Huehn J, Olek S, et al. Maternal farm exposure modulates neonatal immune mechanisms through regulatory T cells. *J Allergy Clin Immunol* 2009;123(4). pp. 774–782. e5.
- Riedler J, Eder W, Oberfeld G, Schreuer M. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin Exp Allergy* 2000;30:194–200.
- Garn H, Neves JF, Blumberg RS, Renz H. Effect of barrier microbes on organ-based inflammation. *J Allergy Clin Immunol* 2013;131(6):1465–78.
- Ege MJ, Bieli C, Frei R, van Strien RT, Riedler J, Üblagger E, et al. Prenatal farm exposure is related to the expression of receptors of the innate immunity and to atopic sensitization in school-age children. *J Allergy Clin Immunol* 2006;117(4):817–23.
- Blümer N, Sel S, Virna S, Patrascu C, Zimmermann S, Herz U, et al. Perinatal maternal application of *Lactobacillus rhamnosus* GG suppresses allergic airway inflammation in mouse offspring. *Clin Exp Allergy* 2007;37(3):348–57.
- Iliev ID, Funari VA, Taylor KD, Nguyen Q, Reyes CN, Strom SP, et al. Interactions between commensal fungi and the C-type lectin receptor Dectin-1 influence colitis.

- Science 2012;1221789.
- [42] Brown GD. Innate antifungal immunity: the key role of phagocytes. *Annu Rev Immunol* 2011;29:1–21.
 - [43] Underhill DM, Iliiev ID. The mycobiota: interactions between commensal fungi and the host immune system. *Nat Rev Immunol* 2014;14(6):405.
 - [44] Kim Y-G, Udayanga KGS, Totsuka N, Weinberg JB, Núñez G, Shibuya A. Gut dysbiosis promotes M2 macrophage polarization and allergic airway inflammation via fungi-induced PGE2. *Cell Host Microbe* 2014;15(1):95–102.
 - [45] Lee YK, Mazmanian SK. Has the microbiota played a critical role in the evolution of the adaptive immune system? *Science* 2010;330(6012):1768–73.
 - [46] Romani L. Immunity to fungal infections. *Nat Rev Immunol* 2011;11(4):275.
 - [47] Limon JJ, Skalski JH, Underhill DM. Commensal fungi in health and disease. *Cell Host Microbe* 2017;22(2):156–65.
 - [48] Iliiev ID, Leonardi I. Fungal dysbiosis: immunity and interactions at mucosal barriers. *Nat Rev Immunol* 2017;17(10):635.
 - [49] Kumamoto CA. Inflammation and gastrointestinal Candida colonization. *Curr Opin Microbiol* 2011;14(4):386–91.
 - [50] Wheeler ML, Limon JJ, Bar AS, Leal CA, Gargus M, Tang J, et al. Immunological consequences of intestinal fungal dysbiosis. *Cell Host Microbe* 2016;19(6):865–73.