

# Verilog Size Checking

## Expressions

- $\mathcal{A}$  is the set of atoms in an expression. In the AST, they correspond to leaves. In System-Verilog, they refer to what the standard calls an “operand” (variables, integers, function calls, slices of a variable, etc.).
- $\mathcal{E}$  is the set of System-Verilog expressions. We have  $\mathcal{A} \subset \mathcal{E}$ . An expression either contains an expression or is an atom.

## Contexts

- Set of contexts identifiers:  $\mathcal{C} \simeq \mathbb{N}$ .
- A context  $q \subseteq \mathcal{Q}$  with:

$$\mathcal{Q} := \underbrace{\{\text{Size } s \mid s \in \mathbb{N}\}}_{\text{Atomic size}} \sqcup \underbrace{\{\text{Id } c \mid c \in \mathcal{C}\}}_{\text{Identity dependencies}} \sqcup \underbrace{\{\text{Add } (c_1, c_2) \mid c_1 \in \mathcal{C}, c_2 \in \mathcal{C}\}}_{\text{Additive dependencies}} \sqcup \underbrace{\{\text{Mul } (s, c) \mid c \in \mathcal{C}, s \in \mathbb{N}\}}_{\text{Multiplicative dependencies}}$$

A context represent all the information needed to compute the size of associated expressions.

- $\Pi : \mathcal{C} \rightarrow 2^{\mathcal{Q}}$  a partial mapping from context identifiers to the smallest possible context set. This means that:

$$\Pi(c) \subset \Pi'(c) \implies \Pi(c) = \Pi'(c)$$

## Rules

The statement  $\Pi \vdash e : c$  means:

In the context environment  $\Pi$ , all the information needed to compute the size of the expression  $e$  is in the context-set identified by  $c$  in  $\Pi$ .

We use the following notations:

- $\Gamma$  compute the size of an atom.
- $\Phi$  compute the size of a lvalue.

### Base case

$$\frac{s = \Gamma(e) \quad \Pi = \{c \mapsto q\} \quad \text{Size } s \in q \quad e \in \mathcal{A}}{\Pi \vdash e : c}$$

### Operators

- $\oplus \in \{+, -, *, /, \%, \&, |, \wedge, \sim, \sim\sim\}$ :

$$\frac{\Pi \vdash a : c \quad \Pi \vdash b : c}{\Pi \vdash a \oplus b : c}$$

- $\oplus \in \{+, -, ++, --, \sim\}$ :

$$\frac{\Pi \vdash e : c}{\Pi \vdash \oplus e : c}$$

- $\oplus \in \{\$signed, \$unsigned\}$ :

$$\frac{\Pi' \vdash e : c' \quad c \notin \text{dom } \Pi' \quad \Pi = \Pi' \sqcup \{c \mapsto q\} \quad \text{Id } c' \in q}{\Pi \vdash \oplus(e) : c}$$

- $\oplus \in \{==, !=, ==?, !=?, ==, !=, >, >=, <, <= \}$ :

$$\frac{\Pi' \vdash a : c' \quad \Pi' \vdash b : c' \quad c \notin \text{dom } \Pi' \quad \Pi = \Pi' \sqcup \{c \mapsto q\} \quad \text{Size } 1 \in q}{\Pi \vdash a \oplus b : c}$$

- $\oplus \in \{\&\&, ||, -, <->\}$ :

$$\frac{\Pi_1 \vdash a : c_1 \quad \Pi_2 \vdash b : c_2 \quad \text{dom } \Pi_1 \cap \text{dom } \Pi_2 = \emptyset \quad c \notin \text{dom } \Pi_1 \sqcup \text{dom } \Pi_2 \quad \Pi = \Pi_1 \sqcup \Pi_2 \sqcup \{c \mapsto q\} \quad \text{Size } 1 \in q}{\Pi \vdash a \oplus b : c}$$

- $\oplus \in \{\&, \sim\&, |, \sim|, \hat{\phantom{x}}, \sim\hat{\phantom{x}}, \hat{\sim}, !\}$ :

$$\frac{\Pi' \vdash a : c' \quad c \notin \text{dom } \Pi' \quad \Pi = \Pi' \sqcup \{c \mapsto q\} \quad \text{Size } 1 \in q}{\Pi \vdash \oplus a : c}$$

- $\oplus \in \{>>, <<, **, >>>, <<<\}$ :

$$\frac{\Pi_1 \vdash a : c \quad \Pi_2 \vdash b : c' \quad \Pi = \Pi_1 \sqcup \Pi_2 \quad \text{dom } \Pi_1 \cap \text{dom } \Pi_2 = \emptyset}{\Pi \vdash a \oplus b : c}$$

- $\oplus \in \{=, +=, -=, *=, /=, \%=, \&=, |=, \hat{=}\}$ :

$$\frac{s = \Phi(l) \quad \Pi' \vdash b : c' \quad \text{Size } s \in \Pi'(c') \quad c \notin \text{dom } \Pi' \quad \Pi = \Pi' \sqcup \{c \mapsto q\} \quad \text{Size } s \in q}{\Pi \vdash l \oplus b : c}$$

- $\oplus \in \{<<=, >>=, <<<=, >>>=\}$ :

$$\frac{s = \Phi(l) \quad \Pi' \vdash b : c' \quad c \notin \text{dom } \Pi' \quad \Pi = \Pi' \sqcup \{c \mapsto q\} \quad \text{Size } s \in q}{\Pi \vdash l \oplus b : c}$$

- If expression:

$$\frac{\Pi_1 \vdash e : c' \quad \Pi_2 \vdash a : c \quad \Pi_2 \vdash b : c \quad \Pi = \Pi_1 \sqcup \Pi_2 \quad \text{dom } \Pi_1 \cap \text{dom } \Pi_2 = \emptyset}{\Pi \vdash e?a:b : c}$$

- Concatenation:

$$\frac{\Pi_1 \vdash e_1 : c_1 \quad \Pi_2 \vdash e_2 : c_2 \quad \text{dom } \Pi_1 \cap \text{dom } \Pi_2 = \emptyset \quad c \notin \text{dom } \Pi_1 \sqcup \text{dom } \Pi_2 \quad \Pi = \Pi_1 \sqcup \Pi_2 \sqcup \{c \mapsto q\} \quad \text{Add } (c_1, c_2) \in q}{\Pi \vdash \{e_1, e_2\} : c}$$

- Replication:

$$\frac{i \in \mathbb{N} \quad \Pi' \vdash e : c' \quad c \notin \text{dom } \Pi' \quad \Pi = \Pi' \sqcup \{c \mapsto q\} \quad \text{Mul } (i, c') \in q}{\Pi \vdash \{i\} e : c}$$